

Review

Short Overview on the Involvement of Lipid Peroxidation Product 4-Hydroxynonenal in Diverse Pathways of Cell Death

Morana Jaganjac¹ , Anita Stojanovic Markovic¹ , Nicola Deiana², Neven Zarkovic^{1,*} 

¹Laboratory for Oxidative Stress (LabOS), Division of Molecular Medicine, Rudjer Boskovic Institute, HR-10000 Zagreb, Croatia

²Uniss PhD School, School of the University of Sassari, 07100 Sassari, Italy

*Correspondence: zarkovic@irb.hr (Neven Zarkovic)

Academic Editor: Guohui Sun

Submitted: 14 January 2025 Revised: 24 March 2025 Accepted: 31 March 2025 Published: 26 September 2025

Abstract

The oxidation of lipids, notably of polyunsaturated fatty acids (PUFAs), under oxidative stress is a self-catalyzed chain reaction that generates reactive aldehydes, among which 4-hydroxynonenal (4-HNE) is considered to act as a second messenger of free radicals. The pleiotropic effects of 4-HNE, which include the regulation of cellular antioxidant capacities, proliferation, differentiation, and apoptosis, are concentration-dependent as they depend on cell type. Therefore, 4-HNE has important roles in various pathophysiological processes and the pathogenesis of acute and chronic diseases, especially degenerative and malignant diseases. Before 4-HNE was recognized as a signaling molecule, it was known to be the cytotoxic mediator of oxidative stress, acting even if lipid peroxidation was not present, because it remains bound to proteins, changing their structure and function. Research in this field has revealed several novel modes of activities of 4-HNE associated with cell death, including not only apoptosis/programmed cell death and necrosis but also ferroptosis, autophagy, pyroptosis, necroptosis, parthanatos, oxeiptosis and cuproptosis. This review shortly summarizes these findings, aiming to encourage further research in the field that might open new ways to use 4-HNE as the bioactive factor for targeted cell death, in particular cancer cells.

Keywords: oxidative stress; cell death; apoptosis; necrosis; ferroptosis

1. Introduction—From Redox Homeostasis to Oxidative Stress, Lipid Peroxidation and Cell Death

Initially thought to be detrimental by-products of metabolism, reactive oxygen species (ROS) are now recognized as important regulators of diverse cellular functions and processes via redox signaling [1,2]. The imbalance in the production of ROS and antioxidants, which favors prooxidants, causes oxidative stress. Antioxidant defenses include both enzymatic and non-enzymatic antioxidants. The thioredoxin system, the glutathione (GSH) system, peroxiredoxin, superoxide dismutase, and catalase are the main endogenous antioxidant defenses [3]. The nuclear factor, erythroid-derived 2-like 2 (Nrf2), which is endogenously inhibited by Kelch-like ECH-associated protein 1 (Keap1), regulates the expression of a multitude of antioxidants and is essential for maintaining redox homeostasis. Under low oxidative stress, Keap1 dissociates from Nrf2, promoting the expression of antioxidant genes [3]. Excessive ROS damage macromolecules, including lipids, thus triggering lipid peroxidation of polyunsaturated fatty acids (PUFA) [4]. Lipid peroxidation of n-6 PUFAs produces the bioactive aldehyde 4-hydroxynonenal (4-HNE), which can have either direct or indirect effects on cellular signaling pathways and functions. The 4-HNE directly interferes with normal cellular processes, enzymatic functions, and functions of organelles [5–7]. The presence of a carbonyl

group at the C1 position, a hydroxyl group, and an α , β -double bond makes this aldehyde extremely reactive with macromolecules containing thiol and amino groups, consequently leading to the formation of Schiff bases or Michael adducts [8]. Michael adducts predominate because the kinetics of Schiff base formation are inherently slow and reversible. Biomolecules such as proteins (especially histidine, cysteine, and lysine residues), lipids (containing an amino group), and nucleic acids (primarily the guanosine portion of DNA) can all readily interact with 4-HNE [9]. When proteins are modified by 4-HNE, their structure and function may change, affecting a variety of physiological and pathological cellular processes [10,11]. For example, 4-HNE adduction to proteins is implicated in the regulation of insulin signaling, the activity of various enzymes, mitochondrial metabolism, fatty acid metabolism, and tubulin polymerization [11]. Moreover, 4-HNE is much more stable than ROS and can therefore reach targets far away from the original site of its origin [12]. 4-HNE concentrations in human blood and serum have been estimated at 0.05 to 1.4 μ M [13,14], while it may be several folds higher in cells and tissues. Under pathological conditions and around the core of the lipid peroxidation site, 4-HNE concentration can increase substantially, even above 100 μ M [1,15]. Cellular 4-HNE concentrations below 5 μ M are essential for normal cellular processes and function, including cell metabolism, mitochondrial function, cell dif-



Copyright: © 2025 The Author(s). Published by IMR Press.
This is an open access article under the CC BY 4.0 license.

Publisher's Note: IMR Press stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

ferentiation, proliferation, kinase activity, and upregulation of antioxidant defense mechanisms [16–19]. On the other side, exposure to higher concentrations of 4-HNE may have detrimental effects on cells and tissues [1]. Diseases associated with redox imbalance, such as neurodegenerative diseases [20], macular degeneration [21], obesity [22], psoriasis [23], cardiovascular disease, atherosclerosis [24], metabolic syndrome [25–28] and malignancies [29,30], are associated with elevated levels of 4-HNE, which is both a causative factor and a biomarker of oxidative stress in many pathologies. The wide-ranging effects of 4-HNE on biological systems make it a key player in oxidative stress-related disorders, while its roles as a signaling molecule, as a modulator of apoptosis, and as a biomarker of disease progression make 4-HNE attractive for research in various acute and chronic diseases and respective therapies.

The most important biological process that occurs in response to harmful events is death, which may be regulated or uncontrolled. Except for necrosis, which is an acute, irreversible, and unspecific process of cellular death, there are several distinct cell death mechanisms, besides apoptosis, all of which are tightly regulated by intracellular signaling pathways. Some scientists subdivide programmed cell death into non-lytic forms, such as apoptosis, in which membrane integrity is maintained, and lytic forms, such as pyroptosis, necroptosis, and ferroptosis, in which membrane integrity is not maintained and which lead to cell disintegration. Apoptosis is a homeostatic process that keeps cell populations in tissues stable and occurs naturally during development and aging. In immunological responses or when cells are damaged by different noxious pathogens, diseases, or harmful substances, apoptosis also takes place as a protective mechanism [31]. To date, 4-HNE is known to be involved in a variety of cell death mechanisms (Fig. 1).

Although the focus of this review will be on the roles of 4-HNE in apoptosis, its involvement in other mechanisms of cell death will also be discussed.

2. Mechanisms of Regulated Cell Death—The Roles of 4-HNE

Various mechanisms and phenotypes of regulated cell death are now known, while apoptosis was the first type of regulated cell death discovered. 4-HNE has been shown to play a role in apoptosis as well as in other forms of regulated cell death. The following is, therefore, an overview of the known mechanisms and involvement of 4-HNE in apoptosis, ferroptosis, autophagy, pyroptosis, necroptosis, parthanatos, oxeiptosis, and cuproptosis.

2.1 The Involvement of 4-HNE in Oxidative Stress-Induced Apoptosis

Apoptosis, a programmed cell death, is a regulated process of intracellular, biochemical, and morphological changes that lead to cell death [32]. Apoptosis is tightly regulated by pro-apoptotic and anti-apoptotic proteins, and

the genes required for appropriate apoptosis are evolutionarily conserved from *nematodes* to *Homo sapiens* [33]. Two major signaling pathways can trigger apoptosis: the extrinsic, often referred to as the “death receptor pathway”, and the intrinsic, also known as the “mitochondrial pathway”.

Transmembrane receptor-mediated interactions are part of the death receptor pathway, also called the extrinsic mechanism of apoptosis. Ligand-receptor complexes are formed when different chemicals bind to their respective death receptors. Tumor necrosis factor (TNF)/TNF-receptor 1 (TNF-R1), Fas ligand (FasL)/Fas receptor (FasR), and TNF-related inducing apoptotic ligand (TRAIL)/TRAIL-receptor 1 are some of these complexes. The adaptor molecule FAS-associated death domain protein (FADD) identifies ligand-receptor complexes and binds to the receptors via death domains. When caspase-8 is activated, it separates from the death-inducing signaling complex (DISC) formed from the adaptor and bound caspase-8. This triggers a series of caspases that further promote apoptosis [34].

2.1.1 Regulation of Fas Death Receptor Signaling by 4-HNE

The binding of FasL to its receptor FasR is an example of 4-HNE-mediated apoptosis. The 4-HNE-induced apoptosis is accompanied by the activation of c-Jun N-terminal kinase (JNK) and caspase-3 [35]. While it was shown that 4-HNE activates JNK, caspase-3, and apoptosis in human lens epithelial cells HLE B-3 and induces Fas expression in a concentration- and time-dependent manner, there are findings indicating that 4-HNE-mediated regulation of Fas is a generalized phenomenon, not strictly limited to the specific cell type [36]. It appears that 4-HNE activates the Fas signaling pathway primarily through a DISC-independent mechanism that does not involve FADD or caspase-8. This is supported by the discovery that 4-HNE can induce Fas-dependent apoptosis in Jurkat cells lacking pro-caspase-8. *In vitro*, the death receptor Fas, which has an extracellular domain with many cysteines, has been shown to form adducts with 4-HNE [37]. In the absence of DISC involvement, 4-HNE-induced Fas activation triggers downstream signaling via apoptosis signal-regulating kinase 1 (ASK1), JNK, and caspase-3. After JNK phosphorylation, the c-jun transcription factor translocates to the nucleus and regulates the expression of pro-apoptotic genes. 4-HNE also activates a negative feedback loop via death domain-associated protein (Daxx), a nuclear protein involved in stress responses. 4-HNE also triggers the induction of Daxx and facilitates its release from the nucleus to the cytosol, where it cooperates with Fas to inhibit downstream pro-apoptotic signaling and modulate apoptosis. Depletion of Daxx further amplifies the activation of JNK and caspase-3, intensifying 4-HNE-induced apoptosis [37]. 4-HNE promotes the expression of both Daxx and Fas [38] and can promote the export of Daxx from the nucleus to the cytoplasm by




Fig. 1. Elevated ROS levels trigger lipid peroxidation, resulting in 4-HNE formation that may, depending on its concentration and macromolecular targets, be implicated in the diverse types of cell death. The figures are drawn in PowerPoint (Microsoft Office 2019, Redmond, WA, USA). Abbreviations: ROS, reactive oxygen species; 4-HNE, 4-hydroxynonenal; PGAM5, phosphoglycerate mutase 5; AIFM1, apoptosis inducing factor mitochondria associated 1; PARP1, poly(ADP-ribose) polymerase 1.

forming covalent bonds with it through histidine residues. The ASK1-JNK signaling pathway and activation of apoptosis can be affected by the Fas-Daxx binding. While the pathway regulated by Fas-ASK1 and JNK is important and causes major damage under oxidative stress, Daxx-Fas coupling restricts the activation of apoptosis under moderate stress. Since 4-HNE is diffusible, as shown *in vivo*, this negative loop can maintain tissue integrity. Since Daxx acts in both the cytosol and the nucleus, its activity as a pro- or anti-apoptotic agent may vary. Differences between types of cells and the relative significance of the JNK and Fas pathways further indicate the complexity of apoptosis regulation by 4-HNE. A schematic representation of the involvement of 4-HNE in apoptosis is presented in Fig. 2.

2.1.2 The Intrinsic Apoptotic Pathway as a Target of 4-HNE

The intrinsic, i.e., mitochondrial pathway of apoptosis is the most common mechanism of apoptosis in vertebrates. Activation of this pathway can be triggered by toxic substances, viral infections, DNA damage, radiation exposure, or growth factor deprivation [31,39]. 4-HNE, a product of lipid peroxidation, damages mitochondrial DNA and activates the intrinsic apoptotic pathway. This damage caused by 4-HNE results in respiratory enzyme defects, lowering the capacity of reducing potential and energy production. 4-HNE can be produced via cytochrome c (cyt-c) oxidation of

cardiolipin, which is an essential phospholipid found in mitochondrial membranes. Cardiolipin oxidation is required for further steps of intrinsic apoptosis [40]. It activates mitochondrial outer membrane permeabilization (MOMP) and causes the activation of proteins of the B-cell lymphoma 2 (Bcl-2) family, which contain pro-apoptotic and anti-apoptotic factors and regulate the mitochondrial pathway of apoptosis. Under physiological cellular conditions, these proteins form heterodimers with Bcl-2-associated X protein (Bax) and Bcl-2 homologous antagonist/killer (Bak) proteins from pro-apoptotic groups, blocking their action and maintaining the integrity of the mitochondrial membrane. 4-HNE can induce mitochondria-dependent apoptosis as shown in mouse leukemic macrophage cell line, with common signs of intrinsic apoptosis, such as DNA fragmentation, pro-apoptotic protein upregulation, and anti-apoptotic protein downregulation [41,42]. When a cell is threatened by DNA damage, BH3-only proteins are expressed, initiating apoptosis and binding anti-apoptotic factors, releasing Bax and Bak proteins. Anti-apoptotic factors, especially myeloid cell leukemia-1, are ubiquitinated and degraded to induce apoptosis. Uninhibited pro-apoptotic factors Bax and Bak form oligomers that insert into the outer mitochondrial membrane, leading to pores and permeabilization. Hence, soluble proteins from the intramembrane space of mitochondria spill into the cytoplasm [31,43], so cyt-c also leaks from the mitochondria into the




Fig. 2. The involvement of 4-HNE in apoptosis. 4-HNE, a by-product of lipid peroxidation, induces apoptosis by activating FAS and the TRAIL pathways. The ASK1/JNK pathway triggers AP-1-dependent transcription through the DISC-independent route. After JNK phosphorylation, the c-jun transcription factor translocates to the nucleus and regulates the expression of pro-apoptotic genes. Exported Daxx is involved in the regulation of the FAS pathway. The intrinsic apoptotic pathway is induced by oxidative stress, p53, and mitochondrial membrane disturbances. The figures are drawn in PowerPoint (Microsoft Office 2019, Redmond, WA, USA) and Inkscape (Software Freedom Conservancy, Brooklyn, NY, USA). Abbreviations: 4-HNE, 4-hydroxynonenal; AP-1, activator protein 1; ASK1, apoptosis signal-regulating kinase 1; Daxx, death domain-associated protein; FADD, Fas-associated protein with death domain; FasL, Fas ligand; JNK, c-Jun N-terminal kinase; cyt-c, cytochrome c; MOMP, mitochondrial outer membrane permeabilization; TRAIL, TNF-related apoptosis-inducing ligand; Bak, Bcl-2 homologous antagonist/killer; Bax, Bcl-2-associated X protein; Bcl-2, B-cell lymphoma 2.

cytoplasm and binds to the apoptotic protease activating factor 1 (Apafl). This leads to the activation of Apafl, which is activated by a circular homoheptameric complex denoted apoptosome. The main role of the apoptosome is to activate the inactive pro-caspase-9. For the proper activity of caspase-9, proteolytic cleavage is not as important as binding to the apoptosome. The holoenzyme of apoptosome and caspase-9 activates executioner caspases (3 and 7) [31,43]. It has been shown that caspase-3 and caspase-7 are crucial for apoptosis. There is a functional redundancy between caspase-3 and caspase-7, but they do not overlap completely [44]. However, caspase-3 is the main executor of the process, while caspase-7 has more of a supporting role by causing an accumulation of ROS [45]. Research on sympathetic and hippocampal neurons showed that the downregulation of caspase-3 and caspase-7 protects from 4-HNE-induced death [46]. In age-related sarcopenia, the increase in 4-HNE generation during muscle cell apoptosis results in the inactivation of anti-apoptotic protein and activation of caspase-9 and JNK [47], with the same observation made in PC-12 pheochromocytoma cells [48], and RKO colon cancer cells [49].

2.1.3 Heat Shock Proteins in the Regulation of 4-HNE-Induced Apoptosis

Apoptotic proteins often interact with heat shock proteins (HSP), which are molecular chaperones that enhance a cellular ability to withstand various types of stress. HSPs can regulate apoptosis at multiple levels [50]. HSP70 plays a crucial role in regulating the intrinsic apoptosis pathway by regulating apoptosis at the level of mitochondrial permeabilization and preventing the recruitment of caspases and the formation of apoptosome [51]. One study showed that 4-HNE did not alter the expression of HSP70 protein, but it caused late apoptosis in retinal pigment epithelial cells ARPE-19 cells along with increased levels of 4-HNE-modified HSP70. While the expression of the X-linked inhibitor of apoptosis protein (XIAP) is downregulated by 4-HNE-modified HSP70, the 4-HNE-mediated apoptosis of ARPE-19 is reduced by overexpression of HSP70 or XIAP [52]. Interestingly, in addition to its pro-apoptotic effects, 4-HNE was also found to activate stress response pathways via heat shock factor 1 (HSF1), which stimulates the expression of HSP40 and HSP70-1, most likely due to 4-HNE disrupting the inhibitory relationship between HSP70-1 and HSF1 [53]. In addition, HSPB1 regulates the intrinsic path-

way by activating the protein kinase B (Akt), preventing the activation of Bax, and inactivating the apoptosis initiator Bcl-2-associated death promoter (Bad). They indirectly reduce the activity of the intrinsic pathway by stabilizing the actin cytoskeleton and maintaining connections between cells. Exposure of cells to 4-HNE reduced the activity of Akt via 4-HNE-mediated Ser473 dephosphorylation [54], therefore, two possible pathways for 4-HNE-mediated downregulation of Akt were proposed, one dependent on protein phosphatase 2A and one independent of protein phosphatase 2A [35]. Tumor protein 53 (p53) is a transcription factor and tumor suppressor that regulates the expression of genes involved in apoptosis, DNA repair, and the cell cycle. The p53 is activated to protect and repair the cell when oxidative stress and DNA damage occur. However, p53 triggers cell death when the stress or damage reaches a certain threshold. By entering the mitochondria and binding to the anti-apoptotic factors Bcl-x and Bcl-2, p53 helps to activate the intrinsic pathway of apoptosis and initiate the transcription of pro-apoptotic genes of the Bcl-2 family. This enables the permeabilization of the mitochondrial membrane. It also promotes the activation of the extrinsic pathway of apoptosis by inducing the transcription of death receptor genes [39,55]. The effect that 4-HNE has on HSPs is shown in Fig. 3, while the effect on p53 has been shown in Fig. 2.

2.1.4 The Possible Relevance of 4-HNE-Induced Apoptosis Signaling in the Progression of Diseases

In Jurkat, human T-cell leukemia cells treated with 4-HNE, glutathione depletion was found to directly activate caspases-8, -9, and -3. Instead of Fas-mediated glutathione release across the plasma membrane, oxidation by 4-HNE was the cause of the underlying glutathione depletion [56]. In HepG2 cells, 4-HNE was shown to trigger apoptotic signaling through both the p53-mediated intrinsic and Fas-mediated extrinsic pathways. 4-HNE activates Bax, cyclin-dependent kinase inhibitor 1A (p21), JNK, and caspase-3, which in turn triggers apoptosis via the p53 pathway in HepG2 cells [38]. DNA fragmentation, cytochrome c and apoptosis-induced factor release from mitochondria, poly (ADP-ribose) polymerase cleavage, Bcl-2 downregulation, Bax overexpression, and caspase activation were among the conventional hallmarks of apoptosis that were demonstrated in 4-HNE-induced chondrocyte death by increased expression of p53 and Fas while suppressing pro-survival Akt kinase activity [57]. Moreover, the LNCaP hormone-sensitive prostate cancer cells showed enhanced anti-proliferative and pro-apoptotic actions of 4-HNE in a time- and dose-dependent manner that corresponded with the activation of p53-mediated intrinsic apoptotic signaling. On the other hand, hormone-insensitive DU145 prostate cancer cells displayed reduced apoptosis and enhanced proliferation, while their JNK was continuously activated, but p53, p21, Bax, and caspase-3 were not [58].

In human umbilical vein endothelial cells subjected to heat stress, the accumulation of 4-HNE was induced by silencing aldehyde dehydrogenase 2 (ALDH2). An ALDH2 agonist (Alda-1) reduced the activation of inflammatory pathways, senescence, and apoptosis induced by heat stress. In an *in vivo* study in lung homogenates of mice subjected to whole-body heating, Alda-1 was shown to significantly reduce the accumulation of 4-HNE and the activation of p65 and p38 [59]. Morphological changes and DNA damage were observed in ARPE-19 retinal pigment epithelial cells after 4-HNE application, but the damage was able to be reversed with Fucoxanthin treatment [60].

Moreover, 4-HNE was shown to cause neuronal cell death by inducing aberrant expression of apoptotic markers, such as caspase-3, Bax, and p53. In neuroblastoma SH-SY5Y cells, increased expressions of Bax and p53 were associated with HNE-induced oxidative stress. Increased caspase-3 expression and activity confirmed mitochondrial-mediated apoptosis [61]. 4-HNE was found to be associated with mortality and oxidative destruction of neurons after subarachnoid hemorrhage, especially in the cerebral cortex and hippocampus on the third day after injury. Administration of edaravone dextrose strongly inhibited the increase of 4-HNE, reduced neuronal death, and improved long-term neurological outcomes by preventing the oxidative stress response associated with 4-HNE [25]. 4-HNE is widely viewed as a central factor in oxidative stress-mediated neuronal apoptosis. For example, researchers found 4-HNE adducts to neurofilaments in Alzheimer's disease [62], while in Parkinson's disease, it can cause neuronal cell death via adducts with proteins connected to the proteasome system [63]. Under conditions marked by elevated ROS, 4-HNE induces neuron death by binding to and altering transmembrane receptors, cytosolic enzymes, and even DNA, ultimately triggering apoptotic signaling cascades.

Furthermore, 4-HNE can impair mitochondrial function or reduce cellular antioxidant capacity, leading to additional ROS generation [64-67]. Metabolic syndrome encompasses several interrelated metabolic disturbances, such as impaired glucose tolerance, insulin resistance, dyslipidemia, obesity, and liver disease. In the context of diabetes, pancreatic β cells are notably vulnerable to ROS, while 4-HNE can induce β cell apoptosis and exacerbate glucose intolerance, thus contributing to the onset or progression of diabetes [68].

Therefore, it is certain that 4-HNE is a key mediator of oxidative stress-induced apoptosis and acts through various mechanisms. Its pathological significance underscores the importance of understanding its role in apoptosis and exploring targeted interventions to mitigate its detrimental effects in various diseases.




Fig. 3. Heat shock proteins and 4-HNE in apoptosis. 4-HNE modifies HSP70 without altering its overall expression levels. Modified HSP70 downregulates the apoptosis-inhibitory protein XIAP, thereby indirectly enhancing apoptosis. Experimental overexpression of either HSP70 or XIAP effectively counteracts 4-HNE-induced apoptosis by providing stronger apoptotic inhibition. 4-HNE activates HSF1, leading to elevated expression of HSP40 and HSP70-1, which enhances the cellular stress response, providing protective, anti-apoptotic effects. HSPB1 stimulates Akt kinase, which inhibits pro-apoptotic proteins Bad and Bax, thereby preventing apoptosis under normal conditions. 4-HNE-mediated inhibition of Akt kinase removes this protective inhibition, indirectly activating Bad/Bax and promoting apoptosis. The figures are drawn in PowerPoint (Microsoft Office 2019, Redmond, WA, USA). Abbreviations: 4-HNE, 4-hydroxynonenal; HSP, heat shock protein; XIAP, X-linked inhibitor of apoptosis protein; HSF1, heat shock factor 1; Akt, protein kinase B; Bad, Bcl-2-associated death promoter; Bax, Bcl-2-associated X protein.

2.2 4-HNE is a Critical Mediator and Biomarker of Ferroptosis

Lipid peroxidation is a hallmark of ferroptosis. In the presence of free intracellular iron, ferroptosis can be triggered by excessive accumulation of ROS, including lipid ROS. Lipid ROS may be reduced by the action of glutathione peroxidase 4 (GPX4) and thus prevent ferroptosis [69]. On the other side, direct inhibition or inactivation of GPX4, inhibition of GSH biosynthesis, or excessive intracellular iron uptake are all conditions that favor ferroptosis-induced cell death [70]. The cystine/glutamate antiporter system (X_c^-) is essential for the intracellular production of GSH by providing cystine. In recent years, GSH-independent ferroptosis defense mechanisms have been identified based on ferroptosis suppressor protein 1 (FSP1) and dihydroorotate dehydrogenase [71–73]. Furthermore, glutathione S-transferase P1 (GSTP1) detoxifies lipid ROS by catalyzing 4-HNE-GSH conjugation independent from GPX4/FSP1 acting as an unconventional

ferroptosis regulator [74]. Excessive iron was shown to accumulate in mitochondria, causing mitochondrial dysfunction in lipopolysaccharide-induced acute lung injury via ferroptosis [75]. Administration of an iron chelator such as deferoxamine or a ferroptosis inhibitor such as ferrostatin-1 (Fer-1) can reduce the amount of available iron, attenuating lipid peroxidation and eventually ferroptosis [75].

Changes in extracellular pH can also trigger ferroptosis. Namely, G protein-coupled receptor 4 (GPR4), an acid-sensing receptor, is implicated in neuronal ferroptosis in early brain injury. Administration of a GPR4 antagonist reduces 4-HNE formation, mitochondrial damage, and neuronal ferroptosis achieved through the Ras homolog family member A/Yes-associated Protein pathway [76]. Furthermore, elevated intracellular pH decreases ALDH2 activity and inhibits the Extracellular Signal-Regulated Kinase/cAMP Response Element-Binding Protein 1/GPX4 axis, preventing its role in inhibiting ferroptosis via 4-HNE detoxification [77]. In addition, 4-HNE reduces GPX4 ex-




Fig. 4. The involvement of 4-HNE in ferroptosis and autophagy. Excessive ROS promotes peroxidation of membrane-bound PUFAs, triggering a chain reaction of lipid peroxidation and formation of lipid ROS and eventually 4-HNE. GPX4 has a major role in lipid ROS detoxification. Inhibition of the cystine/glutamate antiporter X_c^- system deprives cellular cysteine, affecting GSH biosynthesis that consequentially impacts GPX4 activity, resulting in elevated lipid ROS. Intracellular labile iron accumulation further promotes the accumulation of lipid ROS, eventually resulting in ferroptotic cell death. Alternatively, 4-HNE can inhibit autophagy by inhibiting AMPK activity but also promote autophagy via a modified JNK pathway. Moreover, 4-HNE was shown to activate the mammalian target of rapamycin (mTOR) pathway to trigger light chain 3 (LC3) formation and induce endoplasmic reticulum stress, all of which promote autophagy. The figures are drawn in PowerPoint (Microsoft Office 2019, Redmond, WA, USA). Abbreviations: 4-HNE, 4-hydroxynonenal; AMPK, AMP-activated protein kinase; GSH, glutathione; GPX4, glutathione peroxidase 4; JNK, c-Jun N-terminal kinase; X_c^- , the cystine/glutamate antiporter system.

pression and modifies GPX4 function by its carbonylation [78]. A schematic presentation of the roles of 4-HNE in ferroptosis is presented in Fig. 4.

NADPH-oxidase 4 activation was found to promote ferroptosis by impacting myeloperoxidase and osteopontin, causing mitochondrial dysfunction and excessive 4-HNE formation in astrocytes, which ultimately promotes ferroptosis [79]. On the other hand, the release of corticosteroids increased under stress induces activation of the protein kinase C pathway, which facilitates 15-lipoxygenase-1 localization in the vicinity of plasma membranes and promotes lipid peroxidation triggering ferroptosis in dopaminergic neurons [80].

A transmembrane glycoprotein, fibronectin type III domain containing 5 (FNDC5) has cardioprotective effects. When cells are exposed to oxidized low-density lipoprotein (ox-LDL), the expression of FNDC5 and the expression of antioxidants are reduced, while it promotes the release of inflammatory cytokines, lipid peroxidation, and the formation of 4-HNE, ultimately leading to ferroptosis [81]. A recent study identified the important role

of adipose tissue-derived protein asprosin in ferroptosis. Through integrin $\beta 3$ /bone morphogenetic protein/hepcidin-mediated iron overload and lipid peroxidation, asprosin was shown to influence ferroptosis [82]. Because osteoporosis is also associated with ferroptosis, the activation of the vitamin D receptor attenuates ferroptosis via activation of the Nrf2/GPX4 signaling pathway in osteoblasts [83].

Therefore, it is evident that 4-HNE is not only the hallmark of ferroptosis but also a critical factor of ferroptosis in various diseases. Accordingly, a better understanding of the roles 4-HNE has in ferroptotic cell death in different diseases may provide novel potential therapeutic options targeting the pathways that regulate 4-HNE formation, lipid peroxidation, and iron homeostasis.

2.3 Dual Regulation of Autophagy by 4-HNE

Autophagy is a strictly regulated process in which cells degrade cellular components such as proteins and organelles. While stress-induced autophagy can promote cell survival, it is also an essential process alongside other mechanisms of regulated cell death or autophagy-

dependent cell death [34]. Key proteins in autophagy include microtubule-associated protein 1 light chain 3 (LC3) and autophagy-related proteins (ATG), which orchestrate the formation and elongation of the phagophore and, finally, the formation of a mature autophagosome that, following the fusion with the lysosome, forms an autolysosome [84].

Autophagy is closely linked to the physiological response to oxidative stress. Intracellular accumulation of 4-HNE promotes the activation of kinases, including p38 mitogen-activated protein kinase, JNK, and protein kinase R-like endoplasmic reticulum kinase. It also increases the expression of heme oxygenase-1 and LC3-II formation, enabling cell survival, while inhibition of JNK may ameliorate these effects, resulting in cell death [85]. The protective effect of autophagy was also observed in retinal pigment epithelial (RPE) cells, where ethanol exposure damaged mitochondria and caused accumulation of 4-HNE, while autophagy was activated as a defense mechanism of degradation of impacted mitochondria and 4-HNE aggresomes [86]. On the other hand, dysfunctional autophagic machinery promotes the accumulation of oxidized proteins, protein aggregates, damaged lipids, and damaged organelles, thus promoting diseases such as Down syndrome, and age-related macular degeneration [87,88]. The p62, a selective autophagy receptor [89] is another target of 4-HNE. Exposure to 4-HNE upregulates the p62 expression, promoting autophagy of damaged macromolecules and organelles [90,91].

Dysregulated autophagy can also have deleterious effects. Depending on the concentration, 4-HNE exhibits a dual effect on autophagy. 4-HNE can inhibit autophagy via inhibition of AMP-activated protein kinase (AMPK) activity but may also promote autophagy via a modified JNK pathway [85,92]. Additionally, 4-HNE can activate the mTOR pathway [93], induce endoplasmic reticulum stress [85] and trigger LC3 formation [94] promoting autophagy.

Moreover, 4-HNE was shown to activate the mTOR pathway, to trigger LC3 formation, and to induce endoplasmic reticulum stress, all of which promote autophagy. Furthermore, low concentrations of 4-HNE have been shown to promote autophagy in neuronal cells, while at higher concentrations, 4-HNE causes mitochondrial dysfunction and inhibits autophagy, leading to cellular dysfunction [95]. The altered metabolism of cells, such as the inhibition of glycolysis in neuroblastoma cells, can enhance cellular sensitivity to 4-HNE by attenuating autophagy while enhancing apoptosis, highlighting an important role of glucose metabolism in the regulation of autophagy activated by 4-HNE [96]. In the recent review, the current evidence on the involvement of 4-HNE from the initiation of autophagy and the formation of phagophores to the final degradation stage of autophagy has been summarized [97].

The findings regarding the role of 4-HNE in autophagy may represent the basis for the development of innovative strategies to either enhance protective autophagy

or prevent its dysfunction and thus offer promising avenues for treating a wide range of diseases, particularly diseases associated with oxidative stress and autophagic dysfunction.

2.4 4-HNE Involvement in NLRP3 Inflammasome-Mediated Pyroptosis

Inflammatory cell death, known as pyroptosis, is characterized by inflammasome-dependent caspase-1 activity [98]. Nucleotide-binding oligomerization domain-like receptor protein 3 (NLRP3) is an example of an intracellular sensor that can trigger pyroptosis and attract the apoptosis-associated speck-like protein containing a caspase recruitment domain (ASC), yielding inflammasome formation. This promotes caspase-1 activation, which cleaves gasdermin D (GSDMD). It was recently found that degradation of X_c^- via the ubiquitin-proteasome system promotes lipid peroxidation, as evidenced by increased 4-HNE production, which in turn leads to gasdermin D-mediated pyroptosis in chronic obstructive pulmonary disease [99]. Similarly, pyroptosis may be induced by cytosolic lipopolysaccharide, which activates caspase-4/5/11, leading to GSDMD cleavage. Released gasdermin-N then promotes plasma membrane pore formation, leading to ion fluxes, cell swelling, the release of inflammatory mediators, and, ultimately, the lysis of the cell [100].

The ox-LDL-induced activation of the NLRP3 inflammasome is important for the onset and progression of atherosclerosis. Namely, ox-LDL alters cellular redox homeostasis, promotes the formation of 4-HNE, reduces ALDH2 activity, activates the NLRP3 inflammasome, and may cause DNA damage, leading to pyroptosis as observed in RAW264.7 cells. One of ALDH2's primary substrates is 4-HNE, and the reduced enzyme activity could be at least partially due to ox-LDL-induced 4-HNE. Hence, the cells were partially rescued from pyroptosis and excessive 4-HNE by administration of the ALDH2 activator Alda-1 [101]. The protective role of ALDH2 by inhibiting pyroptosis has also been documented in cardiac injury induced by high glucose, which could be attributed at least in part to NLRP3 inflammasome action, excessive mitochondrial ROS production, and lipid peroxidation. Overexpression of ALDH2 in H9C2 cardiac cells exposed to high glucose inhibited mitochondrial ROS production, decreased 4-HNE levels, and decreased key components of pyroptosis, such as caspase-1 activity, NLRP3 inflammasome-related proteins, and ASC, thereby inhibiting pyroptosis [102]. In addition, increased levels of 4-HNE, NLRP3, and gasdermin are induced in the heart and the brain after ischemia/reperfusion injury, while Alda-1 ameliorated oxidative stress, 4-HNE formation, and NLRP3 inflammasome activation, resulting in the inhibition of pyroptosis and improved recovery of the heart and brain in pigs after eight minutes of untreated ventricular fibrillation [103]. The interaction between mitochondrial ALDH2, ferroptosis, and

pyroptosis has also been evidenced in the murine sepsis-induced lung injury model. Namely lung injury was accompanied by excessive ROS, elevated levels of 4-HNE-modified proteins, increased prostaglandin-endoperoxide synthase 2, and decreased GPX4, while administration of Alda1, Fer-1, and the NLRP3 inflammasome inhibitor MCC950 alleviated those effects [104]. On the other hand, in the mouse models of acute lung injury and sepsis, 4-HNE attenuated the NLRP3 inflammasome, acting even as an endogenous inhibitor of pyroptosis [105].

In a mouse model of metabolic dysfunction-associated fatty liver disease, caspase-1 activity, level of 4-HNE, and the expression of GSDMD mRNA are increased [106]. Interestingly, the ferroptosis inhibitor lipoxystatin-1 alleviated these alterations. In addition, lipoxystatin-1 also inhibited apoptosis and necroptosis, suggesting its novel role as an inhibitor of PANoptosis, a type of cell death that involves an interplay of pyroptosis, apoptosis, and necroptosis, sharing their characteristics [107]. This process is regulated by a large protein complex called the PANoptosome, which is made up of sensors, adaptors, and catalytic effectors [108]. A recent study demonstrated that hypoxia-induced cell death involves crosstalk between ferroptosis and PANoptosis and that both processes can be mitigated with the scavenger of lipid peroxidation [109].

Therefore, understanding the intricate interplay between oxidative stress, 4-HNE, inflammasome activation, and pyroptosis is of high relevance for developing targeted therapies to manage diseases linked to pyroptosis and PANoptosis.

2.5 Evidence on the Involvement of 4-HNE in Necroptosis, Parthanatos, Oxeiptosis, and Cuproptosis

The activation of death receptors like TNF-R1 or pathogen recognition receptors like toll-like receptor 4 triggers a regulated form of cell death with a necrotic phenotype known as necroptosis [34,110]. Necroptosis requires activation of both mixed lineage kinase domain like pseudokinase (MLKL) and receptor interacting protein kinase 1 (RIPK1)-induced RIPK3 activation [34]. Recent studies have observed the involvement of lipid peroxidation in necroptosis, too. Hence, linoleic acid and α -linolenic acid treatment of pancreatic cancer cells resulted in elevated lipid peroxidation, which eventually induced the ferroptosis pathway, which in turn triggered the RIPK/MLKL-dependent necroptosis pathway [111]. Furthermore, the study using an animal model of age-related macular degeneration using GPx4 cKO has shown that lipid peroxidation-derived reactive aldehydes, including 4-HNE, can trigger RPE degeneration, while administration of the necroptosis inhibitor necrostatin-1s or the ferroptosis inhibitor Fer-1 ameliorated cellular degeneration. The same results were not observed when apoptosis was inhibited, suggesting the coexistence of necroptosis and ferroptosis in RPE degeneration [112]. Another comparative study of different ox-

idative stress-induced cell death mechanisms of RPE revealed that the accumulation of lipid ROS and activation of RIPK1/RIPK3 are common in both ferroptosis and necroptosis [113]. The same study also showed that although in necroptosis there is a mild accumulation of lipid ROS, 4-HNE-mediated RPE cell death most likely occurred via the necroptotic pathway, although this might depend on the amount of 4-HNE. Necroptosis of cardiomyocytes was induced by 4-HNE after ischemia/reperfusion injury both *in vivo* and *in vitro* [114]. 4-HNE also enhanced necroptosis of H9c2 cells, obtained from rat heart tissue, in a time- and concentration-dependent manner, likely due to 4-HNE-mediated carbonylation of RIPK1, which prevents polyubiquitination of RIPK1 at K48 and subsequent degradation [114]. Furthermore, excessive alcohol consumption aggravates hepatic encephalopathy, and related apoptosis and necroptosis of neurons in the cerebellum is marked by decreased neuronal antioxidant capacity and elevated 4-HNE, pointing to an important role of oxidative stress, lipid peroxidation, and 4-HNE in both apoptotic and necroptotic neuronal decay [115].

The involvement of oxidative stress and lipid peroxidation in parthanatos has not been extensively explored. Hyperactivation of poly(ADP-ribose) polymerase 1 (PARP1) drives the parthanatos mechanism of programmed cell death [34], while one of the factors that triggers PARP1 activation is oxidative stress-induced DNA damage. Namely, a recent study showed a positive correlation of oxidative stress parameters, including 4-HNE, with PARP1 activation and parthanatos in human leukocytes derived from chronic heart failure patients [116].

In the last few years, two new forms of programmed cell death closely associated with oxidative stress were discovered and termed oxeiptosis and cuproptosis [117,118]. Oxeiptosis is a form of cell death triggered by ROS that resembles apoptosis but is independent of caspases [118]. Oxeiptosis is mediated through the interaction of Keap1, mitochondrial serine-threonine phosphatase phosphoglycerate mutase 5 (PGAM5), and apoptosis inducing factor mitochondria associated 1 (AIFM1). The Keap1 and Nrf2 complex is normally attached to mitochondria via PGAM5 [119], while under mitochondrial stress, PGAM5 dissociates from Nrf2, inducing the translocation of Nrf2 to the nucleus and the expression of the mitochondrial unfolded protein response [120]. At high levels of oxidative stress, PGAM5 dissociates from Keap1 and interacts with AIFM1, ultimately resulting in oxeiptotic, non-inflammatory cell death. Although the direct role of 4-HNE in oxeiptosis has not been extensively studied, 4-HNE may influence oxeiptosis, though further research is needed to confirm this potential interaction.

Cuproptosis depends on copper imbalance and mitochondrial respiration. Excessive copper targets lipoylated components of the tricarboxylic acid cycle, which results in protein aggregation, proteo-toxic stress, and cell death




Fig. 5. Necrosis induced by 4-HNE. 4-HNE added at 100 μ M to human carcinoma cells (HeLa; ATCC, Manassas, VA, USA) *in vitro* induces necrosis in some cells in less than five minutes, which is associated with the shrinking of the cells into a rounded, pyknotic shape associated with lipid peroxidation of the cellular membrane manifested by blebbing (left photo, no staining, magnification 650 \times). The remaining cells undergo necrosis in an additional period of 20–30 minutes at the latest. A similar process can be seen *in vivo* in the malignant tissue of human adenocarcinoma, where 4-HNE was detected by monoclonal antibodies specific for the 4-HNE-histidine adducts (brown color), while the absence of 4-HNE-protein adducts can be observed by contrast hematoxylin blue color contrast staining (right photo, magnification 450 \times). Lipid peroxidation generated 4-HNE-protein adducts that caused necrosis and disintegration of cancer tissue. Scale bar = 50 μ m.

[117]. Moreover, excessive copper, acting as a transition metal resembling iron, induces ROS production, reduces antioxidant defenses, and promotes lipid peroxidation [121]. Although 4-HNE and copper induce oxidative stress and peroxidation of lipids, which further generates 4-HNE, the direct involvement of 4-HNE in cuproptosis is not well-established. Thus, further research is needed to determine whether 4-HNE interacts with copper-induced cell death mechanisms.

3. Necrosis

Because 4-HNE has a strong affinity for binding to the proteins, its regulatory as well as toxic effects depend on the function of the proteins modified by 4-HNE and their overall intracellular concentration [2,11,122–125]. Generally speaking, relatively low levels of 4-HNE are regulatory; higher levels induce cellular decay, while very high

levels can induce rapid cell death caused by an irreversible process of autolysis known as necrosis (Fig. 5).

Necrosis is an unspecific, uncontrolled, and irreversible process of cellular death, usually developing after acute and severe damage followed by the rapid degradation of the cells and the release of the cellular contents in the affected tissue, thus spreading the damage and the death of the cells [126]. The involvement of lipid peroxidation as a crucial pathogenic factor of liver necrosis induced by carbon tetrachloride (CCl₄), was known more than fifty years ago [127]. However, at that time, 4-HNE was not known to be an important factor of liver necrosis, as was found afterward to be even in the case of liver inflammation, i.e., hepatitis, and damage of other organs caused by various diseases [128].

Induction of cell proliferation, apoptosis, and necrosis by 4-HNE also depends on the type of cells, among which

cancer cells are more sensitive to the cytotoxic effects of 4-HNE than are non-malignant normal cells [129–131]. The sensitivity of the cells to 4-HNE also depends on the level of differentiation, suggesting that reduced proliferation of cancer cells associated with induced differentiation also increases the sensitivity of cancer cells to the pro-apoptotic and necrotic effects of 4-HNE [132]. However, in case of extremely intense production of 4-HNE and accumulation of its protein adducts, as in the case of COVID-19, 4-HNE can induce local and systemic, unselective ferroptosis and necrosis of normal cells, thus having lethal consequences for the entire organism [133–135].

4. Conclusions and Perspectives

Taking together the complex regulatory and cytotoxic effects of 4-HNE *in vitro* and *in vivo*, we can conclude that oxidative stress, lipid peroxidation, and thus generated 4-HNE have fundamental roles in the death of cells included by different pathways under different pathophysiological and pathological processes. Due to the presence of 4-HNE-modified proteins even after the end of oxidative stress or even in the absence of manifested lipid peroxidation, we can assume that 4-HNE can cause not only acute but also chronic consequences associated with the decay of cells and the entire organism. However, accumulation of the 4-HNE-modified proteins can be reversible in vital cells, if not associated with necrosis, while 4-HNE targets mostly aged and malignant cells. Therefore, options for targeted cytotoxic effects of 4-HNE might be an interesting option for future treatments of degenerative and malignant processes.

Abbreviations

4-HNE, 4-hydroxynonenal; AIFM1, apoptosis inducing factor mitochondria associated 1; Akt, protein kinase B; Alda-1, ALDH2 agonist; ALDH2, aldehyde dehydrogenase 2; AMPK, AMP-activated protein kinase; AP-1, activator protein 1; Apaf1, apoptotic protease activating factor 1; ASC, apoptosis-associated speck-like protein containing a CARD; ASK1, apoptosis signal-regulating kinase 1; ATG, autophagy-related proteins; Bad, Bcl-2-associated death promoter; Bak, Bcl-2 homologous antagonist/killer; Bax, Bcl-2-associated X protein; Bcl-2, B-cell lymphoma 2; cyt-c, cytochrome c; Daxx, death domain-associated protein; DISC, death signaling complex; FADD, FAS-associated death domain protein; FasL, Fas ligand; FasR, Fas receptor; Fer-1, ferrostatin-1; FNDC5, Fibronectin Type III Domain Containing 5; FSP1, ferroptosis suppressor protein 1; GPX4, glutathione peroxidase 4; GSDMD, gasdermin D; GSH, glutathione; GSTP1, glutathione S-transferase P1; HSF1, heat shock factor 1; HSP, heat shock proteins; JNK, c-Jun N-terminal kinase; Keap1, Kelch-like ECH-associated protein 1; LC3, light chain 3; MLKL, mixed lineage kinase domain like pseudokinase; MOMP, mitochondrial outer membrane permeabilization; mTOR, mammalian target of rapamycin; NLRP3, nucleotide binding

oligomerization domain-like receptor protein 3; Nrf2, nuclear factor, erythroid 2 like 2; ox-LDL, oxidized low-density lipoprotein; p21, Cyclin-dependent kinase inhibitor 1A; p53, tumor protein 53; PARP1, poly(ADP-ribose) polymerase 1; PGAM5, phosphoglycerate mutase 5; PUFA, poly unsaturated fatty acids; RIPK1, receptor interacting protein kinase-1; ROS, reactive oxygen species; RPE, retinal pigment epithelial cells; TNF, tumor necrosis factor; TNF-R1, TNF-receptor 1; TRAIL, TNF-related inducing apoptotic ligand; X_c[−], the cystine/glutamate antiporter system; XIAP, X-linked inhibitor of apoptosis protein.

Author Contributions

Conceptualization, NZ; ND, acquisition of references; writing—original draft preparation, MJ, ASM, ND and NZ; writing—review and editing, MJ, ASM and NZ; visualization, MJ and ASM; supervision, NZ and MJ. All authors read and approved the final version of the manuscript. All authors read and approved the final manuscript. All authors have participated sufficiently in the work and agreed to be accountable for all aspects of the work.

Ethics Approval and Consent to Participate

Not applicable.

Acknowledgment

We would like to express our gratitude to all those who helped us during the writing of this manuscript and its revision.

Funding

This research received no external funding.

Conflict of Interest

The authors declare no conflict of interest. Neven Zarkovic is serving as the Guest Editor and Editorial Board member of this journal. We declare that Neven Zarkovic was not involved in the peer review of this article and has no access to information regarding its peer review. Full responsibility for the editorial process for this article was delegated to Guohui Sun.

References

- [1] Jaganjac M, Milkovic L, Gegotek A, Cindric M, Zarkovic K, Skrzyliewska E, *et al.* The relevance of pathophysiological alterations in redox signaling of 4-hydroxynonenal for pharmacological therapies of major stress-associated diseases. *Free Radical Biology & Medicine*. 2020; 157: 128–153. <https://doi.org/10.1016/j.freeradbiomed.2019.11.023>.
- [2] Jaganjac M, Milkovic L, Zarkovic N, Zarkovic K. Oxidative stress and regeneration. *Free Radical Biology & Medicine*. 2022; 181: 154–165. <https://doi.org/10.1016/j.freeradbiomed.2022.02.004>.
- [3] Jaganjac M, Milkovic L, Sunjic SB, Zarkovic N. The Nrf2, Thioredoxin, and Glutathione System in Tumorigenesis and An-

ticancer Therapies. *Antioxidants*. 2020; 9: 1151. <https://doi.org/10.3390/antiox9111151>.

[4] Jaganjac M, Cindrić M, Jakovčević A, Žarković K, Žarković N. Lipid peroxidation in brain tumors. *Neurochemistry International*. 2021; 149: 105118. <https://doi.org/10.1016/j.neuint.2021.105118>.

[5] Csala M, Kardon T, Legeza B, Lizák B, Mandl J, Margittai É, et al. On the role of 4-hydroxynonenal in health and disease. *Biochimica et Biophysica Acta*. 2015; 1852: 826–838. <https://doi.org/10.1016/j.bbacta.2015.01.015>.

[6] Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxidative Medicine and Cellular Longevity*. 2014; 2014: 360438. <https://doi.org/10.1155/2014/360438>.

[7] Zarkovic N. 4-hydroxynonenal as a bioactive marker of pathophysiological processes. *Molecular Aspects of Medicine*. 2003; 24: 281–291. [https://doi.org/10.1016/s0098-2997\(03\)00023-2](https://doi.org/10.1016/s0098-2997(03)00023-2).

[8] Živković M, Žarković K, Škrinjar L, Waeg G, Poljak-Blaži M, Borović Šunjić S, et al. A new method for detection of HNE-histidine conjugates in rat inflammatory cells. *Croatica Chemica Acta*. 2005; 78: 91–98.

[9] Schaur RJ. Basic aspects of the biochemical reactivity of 4-hydroxynonenal. *Molecular Aspects of Medicine*. 2003; 24: 149–159. [https://doi.org/10.1016/s0098-2997\(03\)00009-8](https://doi.org/10.1016/s0098-2997(03)00009-8).

[10] Schaur RJ, Siems W, Bresgen N, Eckl PM. 4-Hydroxy-nonenal-A Bioactive Lipid Peroxidation Product. *Biomolecules*. 2015; 5: 2247–2337. <https://doi.org/10.3390/biom5042247>.

[11] Milkovic L, Zarkovic N, Marusic Z, Zarkovic K, Jaganjac M. The 4-Hydroxynonenal-Protein Adducts and Their Biological Relevance: Are Some Proteins Preferred Targets? *Antioxidants*. 2023; 12: 856. <https://doi.org/10.3390/antiox12040856>.

[12] Jaganjac M, Cipak A, Schaur RJ, Zarkovic N. Pathophysiology of neutrophil-mediated extracellular redox reactions. *Frontiers in Bioscience (Landmark Edition)*. 2016; 21: 839–855. <https://doi.org/10.2741/4423>.

[13] Strohmaier H, Hinghofer-Szalkay H, Schaur RJ. Detection of 4-hydroxynonenal (HNE) as a physiological component in human plasma. *Journal of Lipid Mediators and Cell Signalling*. 1995; 11: 51–61. [https://doi.org/10.1016/0929-7855\(94\)00027-a](https://doi.org/10.1016/0929-7855(94)00027-a).

[14] Gil L, Siems W, Mazurek B, Gross J, Schroeder P, Voss P, et al. Age-associated analysis of oxidative stress parameters in human plasma and erythrocytes. *Free Radical Research*. 2006; 40: 495–505. <https://doi.org/10.1080/10715760600592962>.

[15] Smathers RL, Fritz KS, Galligan JJ, Shearn CT, Reigan P, Marks MJ, et al. Characterization of 4-HNE modified L-FABP reveals alterations in structural and functional dynamics. *PLoS ONE*. 2012; 7: e38459. <https://doi.org/10.1371/journal.pone.0038459>.

[16] Chapple SJ, Cheng X, Mann GE. Effects of 4-hydroxynonenal on vascular endothelial and smooth muscle cell redox signaling and function in health and disease. *Redox Biology*. 2013; 1: 319–331. <https://doi.org/10.1016/j.redox.2013.04.001>.

[17] Al-Menhal AS, Banu S, Angelova PR, Barcaru A, Horvatovich P, Abramov AY, et al. Lipid peroxidation is involved in calcium dependent upregulation of mitochondrial metabolism in skeletal muscle. *Biochimica et Biophysica Acta. General Subjects*. 2020; 1864: 129487. <https://doi.org/10.1016/j.bbagen.2019.129487>.

[18] Al-Menhal AS, Anderson C, Gourine AV, Abramov AY, D'Souza A, Jaganjac M. Proteomic Analysis of Cardiac Adaptation to Exercise by High Resolution Mass Spectrometry. *Frontiers in Molecular Biosciences*. 2021; 8: 723858. <https://doi.org/10.3389/fmolb.2021.723858>.

[19] Ishikado A, Nishio Y, Morino K, Ugi S, Kondo H, Makino T, et al. Low concentration of 4-hydroxy hexenal increases heme oxygenase-1 expression through activation of Nrf2 and antioxidant activity in vascular endothelial cells. *Biochemical and Biophysical Research Communications*. 2010; 402: 99–104. <https://doi.org/10.1016/j.bbrc.2010.09.124>.

[20] Zarkovic K. 4-hydroxynonenal and neurodegenerative diseases. *Molecular Aspects of Medicine*. 2003; 24: 293–303. [https://doi.org/10.1016/s0098-2997\(03\)00024-4](https://doi.org/10.1016/s0098-2997(03)00024-4).

[21] Choudhary S, Xiao T, Srivastava S, Zhang W, Chan LL, Vergara LA, et al. Toxicity and detoxification of lipid-derived aldehydes in cultured retinal pigmented epithelial cells. *Toxicology and Applied Pharmacology*. 2005; 204: 122–134. <https://doi.org/10.1016/j.taap.2004.08.023>.

[22] Halasz M, Łuczaj W, Jarocka-Karpowicz I, Stasiewicz A, Soldo AM, Soldo I, et al. Relationship between systemic biomarker of lipid peroxidation 4-hydroxynonenal and lipidomic profile of morbidly obese patients undergoing bariatric surgery. *Free Radical Biology & Medicine*. 2024; 224: 564–573. <https://doi.org/10.1016/j.freeradbiomed.2024.09.018>.

[23] Wójcik P, Biernacki M, Wroński A, Łuczaj W, Waeg G, Žarković N, et al. Altered Lipid Metabolism in Blood Mononuclear Cells of Psoriatic Patients Indicates Differential Changes in Psoriasis Vulgaris and Psoriatic Arthritis. *International Journal of Molecular Sciences*. 2019; 20: 4249. <https://doi.org/10.3390/ijms20174249>.

[24] Negre-Salvayre A, Vieira O, Escargueil-Blanc I, Salvayre R. Oxidized LDL and 4-hydroxynonenal modulate tyrosine kinase receptor activity. *Molecular Aspects of Medicine*. 2003; 24: 251–261. [https://doi.org/10.1016/s0098-2997\(03\)00020-7](https://doi.org/10.1016/s0098-2997(03)00020-7).

[25] Mattson MP. Roles of the lipid peroxidation product 4-hydroxynonenal in obesity, the metabolic syndrome, and associated vascular and neurodegenerative disorders. *Experimental Gerontology*. 2009; 44: 625–633. <https://doi.org/10.1016/j.exger.2009.07.003>.

[26] Elrayess MA, Almuraikhy S, Kafienah W, Al-Menhal A, Al-Khelaifi F, Bashah M, et al. 4-hydroxynonenal causes impairment of human subcutaneous adipogenesis and induction of adipocyte insulin resistance. *Free Radical Biology & Medicine*. 2017; 104: 129–137. <https://doi.org/10.1016/j.freeradbiomed.2017.01.015>.

[27] Jaganjac M, Almuraikhy S, Al-Khelaifi F, Al-Jaber M, Bashah M, Mazloum NA, et al. Combined metformin and insulin treatment reverses metabolically impaired omental adipogenesis and accumulation of 4-hydroxynonenal in obese diabetic patients. *Redox Biology*. 2017; 12: 483–490. <https://doi.org/10.1016/j.redox.2017.03.012>.

[28] Jaganjac M, Zarkovic N. Lipid Peroxidation Linking Diabetes and Cancer: The Importance of 4-Hydroxynonenal. *Antioxidants & Redox Signaling*. 2022; 37: 1222–1233. <https://doi.org/10.1089/ars.2022.0146>.

[29] Karihtala P, Kauppila S, Puistola U, Jukkola-Vuorinen A. Divergent behaviour of oxidative stress markers 8-hydroxydeoxyguanosine (8-OHDG) and 4-hydroxy-2-nonenal (HNE) in breast carcinogenesis. *Histopathology*. 2011; 58: 854–862. <https://doi.org/10.1111/j.1365-2559.2011.03835.x>.

[30] Perkovic MN, Jaganjac M, Milkovic L, Horvat T, Rojo D, Zarkovic K, et al. Relationship between 4-Hydroxynonenal (4-HNE) as Systemic Biomarker of Lipid Peroxidation and Metabolomic Profiling of Patients with Prostate Cancer. *Biomolecules*. 2023; 13: 145. <https://doi.org/10.3390/biom13010145>.

[31] Elmore S. Apoptosis: a review of programmed cell death. *Toxicologic Pathology*. 2007; 35: 495–516. <https://doi.org/10.1080/01926230701320337>.

[32] Fulda S, Gorman AM, Hori O, Samali A. Cellular stress responses: cell survival and cell death. *International Journal of Cell Biology*. 2010; 2010: 214074. <https://doi.org/10.1155/2010/214074>.

[33] Arya R, White K. Cell death in development: Signaling path-

ways and core mechanisms. *Seminars in Cell & Developmental Biology*. 2015; 39: 12–19. <https://doi.org/10.1016/j.semcd.2015.02.001>.

[34] Galluzzi L, Vitale I, Aaronson SA, Abrams JM, Adam D, Agostinis P, *et al.* Molecular mechanisms of cell death: recommendations of the Nomenclature Committee on Cell Death 2018. *Cell Death and Differentiation*. 2018; 25: 486–541. <https://doi.org/10.1038/s41418-017-0012-4>.

[35] Awasthi YC, Sharma R, Cheng JZ, Yang Y, Sharma A, Singhal SS, *et al.* Role of 4-hydroxynonenal in stress-mediated apoptosis signaling. *Molecular Aspects of Medicine*. 2003; 24: 219–230. [https://doi.org/10.1016/s0098-2997\(03\)00017-7](https://doi.org/10.1016/s0098-2997(03)00017-7).

[36] Li J, Sharma R, Patrick B, Sharma A, Jeyabal PVS, Reddy PMRV, *et al.* Regulation of CD95 (Fas) expression and Fas-mediated apoptotic signaling in HLE B-3 cells by 4-hydroxynonenal. *Biochemistry*. 2006; 45: 12253–12264. <https://doi.org/10.1021/bi060780t>.

[37] Sharma R, Sharma A, Dwivedi S, Zimniak P, Awasthi S, Awasthi YC. 4-Hydroxynonenal self-limits fas-mediated DISC-independent apoptosis by promoting export of Daxx from the nucleus to the cytosol and its binding to Fas. *Biochemistry*. 2008; 47: 143–156. <https://doi.org/10.1021/bi701559f>.

[38] Chaudhary P, Sharma R, Sharma A, Vatsyayan R, Yadav S, Singhal SS, *et al.* Mechanisms of 4-hydroxy-2-nonenal induced pro- and anti-apoptotic signaling. *Biochemistry*. 2010; 49: 6263–6275. <https://doi.org/10.1021/bi100517x>.

[39] Kiraz Y, Adan A, Kartal Yandim M, Baran Y. Major apoptotic mechanisms and genes involved in apoptosis. *Tumour Biology*. 2016; 37: 8471–8486. <https://doi.org/10.1007/s13277-016-5035-9>.

[40] Liu W, Porter NA, Schneider C, Brash AR, Yin H. Formation of 4-hydroxynonenal from cardiolipin oxidation: Intramolecular peroxy radical addition and decomposition. *Free Radical Biology & Medicine*. 2011; 50: 166–178. <https://doi.org/10.1016/j.freeradbiomed.2010.10.709>.

[41] Haynes RL, Brune B, Townsend AJ. Apoptosis in RAW 264.7 cells exposed to 4-hydroxy-2-nonenal: dependence on cytochrome C release but not p53 accumulation. *Free Radical Biology & Medicine*. 2001; 30: 884–894. [https://doi.org/10.1016/s0891-5849\(01\)00476-2](https://doi.org/10.1016/s0891-5849(01)00476-2).

[42] Raza H, John A. 4-hydroxynonenal induces mitochondrial oxidative stress, apoptosis and expression of glutathione S-transferase A4-4 and cytochrome P450 2E1 in PC12 cells. *Toxicology and Applied Pharmacology*. 2006; 216: 309–318. <https://doi.org/10.1016/j.taap.2006.06.001>.

[43] Xiong S, Mu T, Wang G, Jiang X. Mitochondria-mediated apoptosis in mammals. *Protein & Cell*. 2014; 5: 737–749. <https://doi.org/10.1007/s13238-014-0089-1>.

[44] Lakhani SA, Masud A, Kuida K, Porter GA, Jr, Booth CJ, Mehal WZ, *et al.* Caspases 3 and 7: key mediators of mitochondrial events of apoptosis. *Science*. 2006; 311: 847–851. <https://doi.org/10.1126/science.1115035>.

[45] Brentnall M, Rodriguez-Menocal L, De Guevara RL, Cepero E, Boise LH. Caspase-9, caspase-3 and caspase-7 have distinct roles during intrinsic apoptosis. *BMC Cell Biology*. 2013; 14: 32. <https://doi.org/10.1186/1471-2121-14-32>.

[46] Rabacchi SA, Friedman WJ, Shelanski ML, Troy CM. Divergence of the apoptotic pathways induced by 4-hydroxynonenal and amyloid beta-protein. *Neurobiology of Aging*. 2004; 25: 1057–1066. <https://doi.org/10.1016/j.neurobiolaging.2003.10.003>.

[47] Braga M, Sinha Hikim AP, Datta S, Ferrini MG, Brown D, Kovacheva EL, *et al.* Involvement of oxidative stress and caspase 2-mediated intrinsic pathway signaling in age-related increase in muscle cell apoptosis in mice. *Apoptosis*. 2008; 13: 822–832. <https://doi.org/10.1007/s10495-008-0216-7>.

[48] Siddiqui MA, Kumar V, Kashyap MP, Agarwal M, Singh AK, Khanna VK, *et al.* Short-term exposure of 4-hydroxynonenal induces mitochondria-mediated apoptosis in PC12 cells. *Human & Experimental Toxicology*. 2012; 31: 336–345. <https://doi.org/10.1177/0960327111432500>.

[49] Ji C, Amarnath V, Pietenpol JA, Marnett LJ. 4-hydroxynonenal induces apoptosis via caspase-3 activation and cytochrome c release. *Chemical Research in Toxicology*. 2001; 14: 1090–1096. <https://doi.org/10.1021/tx000186f>.

[50] Regimbeau M, Abrey J, Vautrot V, Causse S, Gobbo J, Garrido C. Heat shock proteins and exosomes in cancer therapeutics. *Seminars in Cancer Biology*. 2022; 86: 46–57. <https://doi.org/10.1016/j.semcan.2021.07.014>.

[51] Vostakolaei MA, Hatami-Baroogh L, Babaei G, Molavi O, Kordi S, Abdolalizadeh J. Hsp70 in cancer: A double agent in the battle between survival and death. *Journal of Cellular Physiology*. 2021; 236: 3420–3444. <https://doi.org/10.1002/jcp.30132>.

[52] Yang LL, Chen H, Wang J, Xia T, Sun H, Yuan CH, *et al.* 4-HNE Induces Apoptosis of Human Retinal Pigment Epithelial Cells by Modifying HSP70. *Current Medical Science*. 2019; 39: 442–448. <https://doi.org/10.1007/s11596-019-2057-8>.

[53] Jacobs AT, Marnett LJ. Heat shock factor 1 attenuates 4-Hydroxynonenal-mediated apoptosis: critical role for heat shock protein 70 induction and stabilization of Bcl-XL. *The Journal of Biological Chemistry*. 2007; 282: 33412–33420. <https://doi.org/10.1074/jbc.M706799200>.

[54] Liu W, Akhand AA, Takeda K, Kawamoto Y, Itoigawa M, Kato M, *et al.* Protein phosphatase 2A-linked and -unlinked caspase-dependent pathways for downregulation of Akt kinase triggered by 4-hydroxynonenal. *Cell Death and Differentiation*. 2003; 10: 772–781. <https://doi.org/10.1038/sj.cdd.4401238>.

[55] Chaabane W, User SD, El-Gazzah M, Jaksik R, Sajjadi E, Rzeszowska-Wolny J, *et al.* Autophagy, apoptosis, mitoptosis and necrosis: interdependence between those pathways and effects on cancer. *Archivum Immunologiae et Therapiae Experimentalis*. 2013; 61: 43–58. <https://doi.org/10.1007/s00005-012-0205-y>.

[56] Liu W, Kato M, Akhand AA, Hayakawa A, Suzuki H, Miyata T, *et al.* 4-hydroxynonenal induces a cellular redox status-related activation of the caspase cascade for apoptotic cell death. *Journal of Cell Science*. 2000; 113: 635–641. <https://doi.org/10.1242/jcs.113.4.635>.

[57] Vaillancourt F, Fahmi H, Shi Q, Lavigne P, Ranger P, Fernandes JC, *et al.* 4-Hydroxynonenal induces apoptosis in human osteoarthritic chondrocytes: the protective role of glutathione-S-transferase. *Arthritis Research & Therapy*. 2008; 10: R107. <https://doi.org/10.1186/ar2503>.

[58] Cao ZG, Xu X, Xue YM, Zhao SL. Comparison of 4-hydroxynonenal-induced p53-mediated apoptosis in prostate cancer cells LNCaP and DU145. *Contemporary Oncology*. 2014; 18: 22–28. <https://doi.org/10.5114/wo.2014.40456>.

[59] Tsai HY, Hsu YJ, Lu CY, Tsai MC, Hung WC, Chen PC, *et al.* Pharmacological Activation Of Aldehyde Dehydrogenase 2 Protects Against Heatstroke-Induced Acute Lung Injury by Modulating Oxidative Stress and Endothelial Dysfunction. *Frontiers in Immunology*. 2021; 12: 740562. <https://doi.org/10.3389/fimmu.2021.740562>.

[60] Chiang YF, Chen HY, Chang YJ, Shih YH, Shieh TM, Wang KL, *et al.* Protective Effects of Fucoxanthin on High Glucose- and 4-Hydroxynonenal (4-HNE)-Induced Injury in Human Retinal Pigment Epithelial Cells. *Antioxidants*. 2020; 9: 1176. <https://doi.org/10.3390/antiox9121176>.

[61] Abarikwu SO, Pant AB, Farombi EO. 4-Hydroxynonenal induces mitochondrial-mediated apoptosis and oxidative stress in SH-SY5Y human neuronal cells. *Basic & Clinical Pharmacology & Toxicology*. 2012; 110: 441–448. <https://doi.org/10.1111/j.1365-2710.2012.01912.x>.

1111/j.1742-7843.2011.00834.x.

[62] Perry EA, Castellani RJ, Moreira PI, Nunomura A, Lui Q, Harris PLR, *et al.* Neurofilaments are the major neuronal target of hydroxynonenal-mediated protein cross-links. *Free Radical Research*. 2013; 47: 507–510. <https://doi.org/10.3109/10715762.2013.794265>.

[63] Jenner P. Oxidative stress in Parkinson's disease. *Annals of Neurology*. 2003; 53: S26–S36; discussion S36–S38. <https://doi.org/10.1002/ana.10483>.

[64] Dalleau S, Baradat M, Guéraud F, Huc L. Cell death and diseases related to oxidative stress: 4-hydroxynonenal (HNE) in the balance. *Cell Death and Differentiation*. 2013; 20: 1615–1630. <https://doi.org/10.1038/cdd.2013.138>.

[65] Perluigi M, Coccia R, Butterfield DA. 4-Hydroxy-2-nonenal, a reactive product of lipid peroxidation, and neurodegenerative diseases: a toxic combination illuminated by redox proteomics studies. *Antioxidants & Redox Signaling*. 2012; 17: 1590–1609. <https://doi.org/10.1089/ars.2011.4406>.

[66] Lee JY, Jung GY, Heo HJ, Yun MR, Park JY, Bae SS, *et al.* 4-Hydroxynonenal induces vascular smooth muscle cell apoptosis through mitochondrial generation of reactive oxygen species. *Toxicology Letters*. 2006; 166: 212–221. <https://doi.org/10.1016/j.toxlet.2006.07.305>.

[67] Poli G, Schaur RJ, Siems WG, Leonarduzzi G. 4-hydroxynonenal: a membrane lipid oxidation product of medicinal interest. *Medicinal Research Reviews*. 2008; 28: 569–631. <https://doi.org/10.1002/med.20117>.

[68] Mattson MP. Roles of the lipid peroxidation product 4-hydroxynonenal in obesity, the metabolic syndrome, and associated vascular and neurodegenerative disorders. *Experimental Gerontology*. 2009; 44: 625–633. <https://doi.org/10.1016/j.exger.2009.07.003>.

[69] Yang WS, SriRamaratnam R, Welsch ME, Shimada K, Skouta R, Viswanathan VS, *et al.* Regulation of ferroptotic cancer cell death by GPX4. *Cell*. 2014; 156: 317–331. <https://doi.org/10.1016/j.cell.2013.12.010>.

[70] Ulker OC, Panieri E, Suzen S, Jaganjac M, Zarkovic N, Saso L. Short overview on the relevance of microRNA-reactive oxygen species (ROS) interactions and lipid peroxidation for modulation of oxidative stress-mediated signalling pathways in cancer treatment. *The Journal of Pharmacy and Pharmacology*. 2022; 74: 503–515. <https://doi.org/10.1093/jpp/rbab045>.

[71] Doll S, Freitas FP, Shah R, Aldrovandi M, da Silva MC, Ingold I, *et al.* FSP1 is a glutathione-independent ferroptosis suppressor. *Nature*. 2019; 575: 693–698. <https://doi.org/10.1038/s41586-019-1707-0>.

[72] Bersuker K, Hendricks JM, Li Z, Magtanong L, Ford B, Tang PH, *et al.* The CoQ oxidoreductase FSP1 acts parallel to GPX4 to inhibit ferroptosis. *Nature*. 2019; 575: 688–692. <https://doi.org/10.1038/s41586-019-1705-2>.

[73] Mao C, Liu X, Zhang Y, Lei G, Yan Y, Lee H, *et al.* DHODH-mediated ferroptosis defence is a targetable vulnerability in cancer. *Nature*. 2021; 593: 586–590. <https://doi.org/10.1038/s41586-021-03539-7>.

[74] Zhang W, Dai J, Hou G, Liu H, Zheng S, Wang X, *et al.* SMURF2 predisposes cancer cell toward ferroptosis in GPX4-independent manners by promoting GSTP1 degradation. *Molecular Cell*. 2023; 83: 4352–4369.e8. <https://doi.org/10.1016/j.molcel.2023.10.042>.

[75] Wang X, Wei T, Luo J, Lang K, Song Y, Ning X, *et al.* Iron Overload-Dependent Ferroptosis Aggravates LPS-Induced Acute Lung Injury by Impairing Mitochondrial Function. *Inflammation*. 2024; 47: 2013–2026. <https://doi.org/10.1007/s10753-024-02022-5>.

[76] He Q, Zhou Y, Wu L, Huang L, Yuan Y, Flores JJ, *et al.* Inhibition of acid-sensing receptor GPR4 attenuates neuronal ferroptosis via RhoA/YAP signaling in a rat model of subarachnoid hemorrhage. *Free Radical Biology & Medicine*. 2024; 225: 333–345. <https://doi.org/10.1016/j.freeradbiomed.2024.10.273>.

[77] Shan G, Bian Y, Yao G, Liang J, Shi H, Hu Z, *et al.* Targeting ALDH2 to augment platinum-based chemosensitivity through ferroptosis in lung adenocarcinoma. *Free Radical Biology & Medicine*. 2024; 224: 310–324. <https://doi.org/10.1016/j.freeradbiomed.2024.08.026>.

[78] Liu L, Pang J, Qin D, Li R, Zou D, Chi K, *et al.* Deubiquitinase OTUD5 as a Novel Protector against 4-HNE-Triggered Ferroptosis in Myocardial Ischemia/Reperfusion Injury. *Advanced Science*. 2023; 10: e2301852. <https://doi.org/10.1002/advs.202301852>.

[79] Boonpraman N, Yoon S, Kim CY, Moon JS, Yi SS. NOX4 as a critical effector mediating neuroinflammatory cytokines, myeloperoxidase and osteopontin, specifically in astrocytes in the hippocampus in Parkinson's disease. *Redox Biology*. 2023; 62: 102698. <https://doi.org/10.1016/j.redox.2023.102698>.

[80] Lin XM, Pan MH, Sun J, Wang M, Huang ZH, Wang G, *et al.* Membrane phospholipid peroxidation promotes loss of dopaminergic neurons in psychological stress-induced Parkinson's disease susceptibility. *Aging Cell*. 2023; 22: e13970. <https://doi.org/10.1111/acel.13970>.

[81] Du Y, Ouyang H, Wang H, Shi J, Li Y, Xu L, *et al.* FNDC5 alleviates Ox-LDL-induced HUVECs dysfunction by inhibiting ferroptosis through AMPK/Nrf2 signaling. *Tissue and Cell*. 2025; 93: 102696. <https://doi.org/10.1016/j.tice.2024.102696>.

[82] Zheng F, Lei JZ, Wang JX, Xu XY, Zhou B, Ge R, *et al.* Crucial roles of asprosin in cisplatin-induced ferroptosis and acute kidney injury. *Free Radical Biology & Medicine*. 2025; 227: 296–311. <https://doi.org/10.1016/j.freeradbiomed.2024.12.024>.

[83] Xu P, Lin B, Deng X, Huang K, Zhang Y, Wang N. VDR activation attenuates osteoblastic ferroptosis and senescence by stimulating the Nrf2/GPX4 pathway in age-related osteoporosis. *Free Radical Biology & Medicine*. 2022; 193: 720–735. <https://doi.org/10.1016/j.freeradbiomed.2022.11.013>.

[84] Dikic I, Elazar Z. Mechanism and medical implications of mammalian autophagy. *Nature Reviews Molecular Cell Biology*. 2018; 19: 349–364. <https://doi.org/10.1038/s41580-018-0003-4>.

[85] Haberzettl P, Hill BG. Oxidized lipids activate autophagy in a JNK-dependent manner by stimulating the endoplasmic reticulum stress response. *Redox Biology*. 2013; 1: 56–64. <https://doi.org/10.1016/j.redox.2012.10.003>.

[86] Flores-Bellver M, Bonet-Ponce L, Barcia JM, Garcia-Verdugo JM, Martinez-Gil N, Saez-Atienzar S, *et al.* Autophagy and mitochondrial alterations in human retinal pigment epithelial cells induced by ethanol: implications of 4-hydroxy-nonenal. *Cell Death & Disease*. 2014; 5: e1328. <https://doi.org/10.1038/cddis.2014.288>.

[87] Di Domenico F, Tramutola A, Barone E, Lanzillotta C, Defever O, Arena A, *et al.* Restoration of aberrant mTOR signaling by intranasal rapamycin reduces oxidative damage: Focus on HNE-modified proteins in a mouse model of down syndrome. *Redox Biology*. 2019; 23: 101162. <https://doi.org/10.1016/j.redox.2019.101162>.

[88] Wen C, Yu X, Zhu J, Zeng J, Kuang X, Zhang Y, *et al.* Gastrodin ameliorates oxidative stress-induced RPE damage by facilitating autophagy and phagocytosis through PPAR α -TFEB/CD36 signal pathway. *Free Radical Biology & Medicine*. 2024; 224: 103–116. <https://doi.org/10.1016/j.freeradbiomed.2024.08.023>.

[89] Ning S, Wang L. The Multifunctional Protein p62 and Its Mechanistic Roles in Cancers. *Current Cancer Drug Targets*. 2019; 19: 468–478. <https://doi.org/10.2174/1568009618666181016164920>.

[90] Zhang S, Eitan E, Mattson MP. Early involvement of lyso-

some dysfunction in the degeneration of cerebral cortical neurons caused by the lipid peroxidation product 4-hydroxynonenal. *Journal of Neurochemistry*. 2017; 140: 941–954. <https://doi.org/10.1111/jnc.13957>.

[91] Baba SP, Zhang D, Singh M, Dassanayaka S, Xie Z, Jagathesan G, *et al.* Deficiency of aldose reductase exacerbates early pressure overload-induced cardiac dysfunction and autophagy in mice. *Journal of Molecular and Cellular Cardiology*. 2018; 118: 183–192. <https://doi.org/10.1016/j.yjmcc.2018.04.002>.

[92] Qin QF, Li XJ, Li YS, Zhang WK, Tian GH, Shang HC, *et al.* AMPK-ERK/CARM1 Signaling Pathways Affect Autophagy of Hepatic Cells in Samples of Liver Cancer Patients. *Frontiers in Oncology*. 2019; 9: 1247. <https://doi.org/10.3389/fonc.2019.01247>.

[93] Calamaras TD, Lee C, Lan F, Ido Y, Siwik DA, Colucci WS. The lipid peroxidation product 4-hydroxy-trans-2-nonenal causes protein synthesis in cardiac myocytes via activated mTORC1-p70S6K-RPS6 signaling. *Free Radical Biology & Medicine*. 2015; 82: 137–146. <https://doi.org/10.1016/j.freeradbiomed.2015.01.007>.

[94] Hill BG, Haberzettl P, Ahmed Y, Srivastava S, Bhatnagar A. Unsaturated lipid peroxidation-derived aldehydes activate autophagy in vascular smooth-muscle cells. *The Biochemical Journal*. 2008; 410: 525–534. <https://doi.org/10.1042/BJ20071063>.

[95] Dodson M, Wani WY, Redmann M, Benavides GA, Johnson MS, Ouyang X, *et al.* Regulation of autophagy, mitochondrial dynamics, and cellular bioenergetics by 4-hydroxynonenal in primary neurons. *Autophagy*. 2017; 13: 1828–1840. <https://doi.org/10.1080/15548627.2017.1356948>.

[96] Dodson M, Liang Q, Johnson MS, Redmann M, Fineberg N, Darley-Usmar VM, *et al.* Inhibition of glycolysis attenuates 4-hydroxynonenal-dependent autophagy and exacerbates apoptosis in differentiated SH-SY5Y neuroblastoma cells. *Autophagy*. 2013; 9: 1996–2008. <https://doi.org/10.4161/auto.26094>.

[97] Gęgotek A, Skrzypieńska E. Lipid peroxidation products' role in autophagy regulation. *Free Radical Biology & Medicine*. 2024; 212: 375–383. <https://doi.org/10.1016/j.freeradbiomed.2024.01.001>.

[98] Bergsbaken T, Fink SL, Cookson BT. Pyroptosis: host cell death and inflammation. *Nature Reviews Microbiology*. 2009; 7: 99–109. <https://doi.org/10.1038/nrmicro2070>.

[99] Hou T, Zhu L, Zhang Y, Tang Y, Gao Y, Hua S, *et al.* Lipid peroxidation triggered by the degradation of xCT contributes to gasdermin D-mediated pyroptosis in COPD. *Redox Biology*. 2024; 77: 103388. <https://doi.org/10.1016/j.redox.2024.103388>.

[100] Shi J, Gao W, Shao F. Pyroptosis: Gasdermin-Mediated Programmed Necrotic Cell Death. *Trends in Biochemical Sciences*. 2017; 42: 245–254. <https://doi.org/10.1016/j.tibs.2016.10.004>.

[101] Xu Y, Yuan Q, Cao S, Cui S, Xue L, Song X, *et al.* Aldehyde dehydrogenase 2 inhibited oxidized LDL-induced NLRP3 inflammasome priming and activation via attenuating oxidative stress. *Biochemical and Biophysical Research Communications*. 2020; 529: 998–1004. <https://doi.org/10.1016/j.bbrc.2020.06.075>.

[102] Cao R, Fang D, Wang J, Yu Y, Ye H, Kang P, *et al.* ALDH2 Overexpression Alleviates High Glucose-Induced Cardiotoxicity by Inhibiting NLRP3 Inflammasome Activation. *Journal of Diabetes Research*. 2019; 2019: 4857921. <https://doi.org/10.1155/2019/4857921>.

[103] Diao M, Xu J, Wang J, Zhang M, Wu C, Hu X, *et al.* Alda-1, an Activator of ALDH2, Improves Postresuscitation Cardiac and Neurological Outcomes by Inhibiting Pyroptosis in Swine. *Neurochemical Research*. 2022; 47: 1097–1109. <https://doi.org/10.1007/s11064-021-03511-x>.

[104] Cao Z, Qin H, Huang Y, Zhao Y, Chen Z, Hu J, *et al.* Crosstalk of pyroptosis, ferroptosis, and mitochondrial aldehyde dehydrogenase 2-related mechanisms in sepsis-induced lung injury in a mouse model. *Bioengineered*. 2022; 13: 4810–4820. <https://doi.org/10.1080/21655979.2022.2033381>.

[105] Hsu CG, Chávez CL, Zhang C, Sowden M, Yan C, Berk BC. The lipid peroxidation product 4-hydroxynonenal inhibits NLRP3 inflammasome activation and macrophage pyroptosis. *Cell Death and Differentiation*. 2022; 29: 1790–1803. <https://doi.org/10.1038/s41418-022-00966-5>.

[106] Tong J, Lan XT, Zhang Z, Liu Y, Sun DY, Wang XJ, *et al.* Ferroptosis inhibitor liproxstatin-1 alleviates metabolic dysfunction-associated fatty liver disease in mice: potential involvement of PANoptosis. *Acta Pharmacologica Sinica*. 2023; 44: 1014–1028. <https://doi.org/10.1038/s41401-022-01010-5>.

[107] Malireddi RKS, Kesavardhana S, Kanneganti TD. ZBP1 and TAK1: Master Regulators of NLRP3 Inflammasome/Pyroptosis, Apoptosis, and Necroptosis (PAN-optosis). *Frontiers in Cellular and Infection Microbiology*. 2019; 9: 406. <https://doi.org/10.3389/fcimb.2019.00406>.

[108] Christgen S, Zheng M, Kesavardhana S, Karki R, Malireddi RKS, Banoth B, *et al.* Identification of the PANoptosome: A Molecular Platform Triggering Pyroptosis, Apoptosis, and Necroptosis (PANoptosis). *Frontiers in Cellular and Infection Microbiology*. 2020; 10: 237. <https://doi.org/10.3389/fcimb.2020.00237>.

[109] Wu MF, Peng X, Zhang MC, Guo H, Xie HT. Ferroptosis and PANoptosis under hypoxia pivoting on the crosstalk between DHODH and GPX4 in corneal epithelium. *Free Radical Biology & Medicine*. 2025; 228: 173–182. <https://doi.org/10.1016/j.freeradbiomed.2024.12.050>.

[110] Vandenabeele P, Galluzzi L, Vanden Berghe T, Kroemer G. Molecular mechanisms of necroptosis: an ordered cellular explosion. *Nature Reviews Molecular Cell Biology*. 2010; 11: 700–714. <https://doi.org/10.1038/nrm2970>.

[111] Suda A, Umaru BA, Yamamoto Y, Shima H, Saiki Y, Pan Y, *et al.* Polyunsaturated fatty acids-induced ferroptosis suppresses pancreatic cancer growth. *Scientific Reports*. 2024; 14: 4409. <https://doi.org/10.1038/s41598-024-55050-4>.

[112] Azuma K, Suzuki T, Kobayashi K, Nagahara M, Imai H, Suga A, *et al.* Retinal pigment epithelium-specific ablation of GPx4 in adult mice recapitulates key features of geographic atrophy in age-related macular degeneration. *Cell Death & Disease*. 2024; 15: 763. <https://doi.org/10.1038/s41419-024-07150-2>.

[113] Tong Y, Wu Y, Ma J, Ikeda M, Ide T, Griffin CT, *et al.* Comparative mechanistic study of RPE cell death induced by different oxidative stresses. *Redox Biology*. 2023; 65: 102840. <https://doi.org/10.1016/j.redox.2023.102840>.

[114] Zhai X, Wang W, Sun S, Han Y, Li J, Cao S, *et al.* 4-Hydroxy-2-Nonenal Promotes Cardiomyocyte Necroptosis via Stabilizing Receptor-Interacting Serine/Threonine-Protein Kinase 1. *Frontiers in Cell and Developmental Biology*. 2021; 9: 721795. <https://doi.org/10.3389/fcell.2021.721795>.

[115] Tamnanloo F, Chen X, Oliveira MM, Tremblay M, Rose CF. Excessive intragastric alcohol administration exacerbates hepatic encephalopathy and provokes neuronal cell death in male rats with chronic liver disease. *Journal of Neuroscience Research*. 2024; 102: e25337. <https://doi.org/10.1002/jnr.25337>.

[116] Bárán T, Simon A, Szabó G, Benkő R, Mezei Z, Molnár L, *et al.* Oxidative Stress-Related Parthanatos of Circulating Mononuclear Leukocytes in Heart Failure. *Oxidative Medicine and Cellular Longevity*. 2017; 2017: 1249614. <https://doi.org/10.1155/2017/1249614>.

[117] Tsvetkov P, Coy S, Petrova B, Dreishpoon M, Verma A, Abdusamad M, *et al.* Copper induces cell death by targeting lipoylated TCA cycle proteins. *Science*. 2022; 375: 1254–1261. <https://doi.org/10.1126/science.abf0529>.

[118] Holze C, Michaudel C, Mackowiak C, Haas DA, Benda C, Hubel P, *et al.* Oxeiptosis, a ROS-induced caspase-independent

apoptosis-like cell-death pathway. *Nature Immunology*. 2018; 19: 130–140. <https://doi.org/10.1038/s41590-017-0013-y>.

[119] Lo SC, Hannink M. PGAM5 tethers a ternary complex containing Keap1 and Nrf2 to mitochondria. *Experimental Cell Research*. 2008; 314: 1789–1803. <https://doi.org/10.1016/j.yexcr.2008.02.014>.

[120] Safakli RN, Gray SP, Bernardi N, Smyrnias I. Unveiling a novel signalling pathway involving NRF2 and PGAM5 in regulating the mitochondrial unfolded protein response in stressed cardiomyocytes. *The International Journal of Biochemistry & Cell Biology*. 2025; 178: 106704. <https://doi.org/10.1016/j.biocel.2024.106704>.

[121] Lu Q, Zhang Y, Zhao C, Zhang H, Pu Y, Yin L. Copper induces oxidative stress and apoptosis of hippocampal neuron via pCREB/BDNF/ and Nrf2/HO-1/NQO1 pathway. *Journal of Applied Toxicology*. 2022; 42: 694–705. <https://doi.org/10.1002/jat.4252>.

[122] Zarkovic N, Ilic Z, Jurin M, Schaur RJ, Puhl H, Esterbauer H. Stimulation of HeLa cell growth by physiological concentrations of 4-hydroxynonenal. *Cell Biochemistry and Function*. 1993; 11: 279–286. <https://doi.org/10.1002/cbf.290110409>.

[123] Sovic A, Borovic S, Loncaric I, Kreuzer T, Zarkovic K, Vukovic T, *et al.* The carcinostatic and proapoptotic potential of 4-hydroxynonenal in HeLa cells is associated with its conjugation to cellular proteins. *Anticancer Research*. 2001; 21: 1997–2004.

[124] Borovic S, Rabuzin F, Waeg G, Zarkovic N. Enzyme-linked immunosorbent assay for 4-hydroxynonenal-histidine conjugates. *Free Radical Research*. 2006; 40: 809–820. <https://doi.org/10.1080/10715760600693422>.

[125] Gasparovic AC, Milkovic L, Sunjic SB, Zarkovic N. Cancer growth regulation by 4-hydroxynonenal. *Free Radical Biology & Medicine*. 2017; 111: 226–234. <https://doi.org/10.1016/j.freeradbiomed.2017.01.030>.

[126] D'Arcy MS. Cell death: a review of the major forms of apoptosis, necrosis and autophagy. *Cell Biology International*. 2019; 43: 582–592. <https://doi.org/10.1002/cbin.11137>.

[127] Cignoli EV, Castro JA. Lipid peroxidation, necrosis and the in vivo depression of liver glucose 6-phosphatase by carbon tetrachloride. *Experimental and Molecular Pathology*. 1971; 14: 43–56. [https://doi.org/10.1016/0014-4800\(71\)90051-7](https://doi.org/10.1016/0014-4800(71)90051-7).

[128] Zarkovic K, Jakovcevic A, Zarkovic N. Contribution of the HNE-immunohistochemistry to modern pathological concepts of major human diseases. *Free Radical Biology & Medicine*. 2017; 111: 110–126. <https://doi.org/10.1016/j.freeradbiomed.2016.12.009>.

[129] Borovic S, Cipak A, Meinitzer A, Kejla Z, Perovic D, Waeg G, *et al.* Differential sensitivity to 4-hydroxynonenal for normal and malignant mesenchymal cells. *Redox Report*. 2007; 12: 50–54. <https://doi.org/10.1179/135100007X162194>.

[130] Semlitsch T, Tillian HM, Zarkovic N, Borovic S, Purtscher M, Hohenwarter O, *et al.* Differential influence of the lipid peroxidation product 4-hydroxynonenal on the growth of human lymphatic leukaemia cells and human peripheral blood lymphocytes. *Anticancer Research*. 2002; 22: 1689–1697.

[131] Bauer G, Zarkovic N. Revealing mechanisms of selective, concentration-dependent potentials of 4-hydroxy-2-nonenal to induce apoptosis in cancer cells through inactivation of membrane-associated catalase. *Free Radical Biology & Medicine*. 2015; 81: 128–144. <https://doi.org/10.1016/j.freeradbiomed.2015.01.010>.

[132] Sunjic SB, Gasparovic AC, Jaganjac M, Rechberger G, Meinitzer A, Grune T, *et al.* Sensitivity of Osteosarcoma Cells to Concentration-Dependent Bioactivities of Lipid Peroxidation Product 4-Hydroxynonenal Depend on Their Level of Differentiation. *Cells*. 2021; 10: 269. <https://doi.org/10.3390/cells10020269>.

[133] Jacobs W, Lammens M, Kerckhofs A, Voets E, Van San E, Van Coillie S, *et al.* Fatal lymphocytic cardiac damage in coronavirus disease 2019 (COVID-19): autopsy reveals a ferroptosis signature. *ESC Heart Failure*. 2020; 7: 3772–3781. <https://doi.org/10.1002/ehf2.12958>.

[134] Žarković N, Orebovec B, Milković L, Baršić B, Tatzber F, Wonisch W, *et al.* Preliminary Findings on the Association of the Lipid Peroxidation Product 4-Hydroxynonenal with the Lethal Outcome of Aggressive COVID-19. *Antioxidants*. 2021; 10: 1341. <https://doi.org/10.3390/antiox10091341>.

[135] Zarkovic N, Jakovcevic A, Mataic A, Jaganjac M, Vukovic T, Waeg G, *et al.* Post-mortem Findings of Inflammatory Cells and the Association of 4-Hydroxynonenal with Systemic Vascular and Oxidative Stress in Lethal COVID-19. *Cells*. 2022; 11: 444. <https://doi.org/10.3390/cells11030444>.