CERN-EP-2025-067
2025/09/16

CMS-HIG-23-014

Measurements of inclusive and differential Higgs boson production cross sections at $\sqrt{s} = 13.6$ TeV in the $H \rightarrow \gamma\gamma$ decay channel

The CMS Collaboration*

Abstract

Inclusive and differential cross sections for Higgs boson production in proton-proton collisions at a centre-of-mass energy of 13.6 TeV are measured using data collected with the CMS detector at the LHC in 2022, corresponding to an integrated luminosity of 34.7 fb^{-1} . Events with the diphoton final state are selected, and the measured inclusive fiducial cross section is $\sigma_{\text{fid}} = 74 \pm 11 \text{ (stat)}^{+5}_{-4} \text{ (syst) fb}$, in agreement with the standard model prediction of $67.8 \pm 3.8 \text{ fb}$. Differential cross sections are measured as functions of several observables: the Higgs boson transverse momentum and rapidity, the number of associated jets, and the transverse momentum of the leading jet in the event. Within the uncertainties, the differential cross sections agree with the standard model predictions.

Published in the Journal of High Energy Physics as doi:10.1007/JHEP09(2025)070.

1 Introduction

The Higgs boson (H) was discovered in 2012 by the ATLAS [1] and CMS [2, 3] Collaborations using proton-proton (pp) collisions at the CERN LHC [4]. Since then, many properties of the Higgs boson have been examined [5, 6]. The large amount of data collected since the observation permits detailed testing of the standard model (SM) with differential measurements. Whereas measurements in the simplified template cross section scheme target the different production modes and their differential properties, the fiducial inclusive and differential cross sections account for the sum of all production mode contributions to the fiducial phase space and are hence less model dependent [7].

The ATLAS and CMS Collaborations measured fiducial inclusive and differential cross sections at the centre-of-mass energies of 7, 8, and 13 TeV in the $H \rightarrow \gamma\gamma$ [8–11], $H \rightarrow ZZ \rightarrow 4\ell$ [12–15], $H \rightarrow WW \rightarrow \ell\ell'\nu\nu'$ [16–20], and $H \rightarrow \tau\tau$ [21, 22] decay channels. The measurements in the diphoton, four-lepton, and WW decay channels by the ATLAS [10, 14, 23] and CMS [11, 15, 20] Collaborations at 13 TeV reach a relative uncertainty of 8–11%. The combination of the ATLAS measurements in the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ \rightarrow 4\ell$ decay channels under the SM assumptions results in an uncertainty of 7% in the total cross section for Higgs boson production [24].

In 2022, the Run 3 of the LHC started at the increased centre-of-mass energy of 13.6 TeV, providing the opportunity to extend Higgs boson cross section measurements to the increased energy and further test the SM predictions. Inclusive fiducial cross sections for Higgs boson production were measured at $\sqrt{s} = 13.6$ TeV in the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ \rightarrow 4\ell$ decay channels by the ATLAS Collaboration with relative uncertainties of about 17 and 26%, respectively [25]. The CMS Collaboration measured the inclusive fiducial cross section in the $H \rightarrow ZZ \rightarrow 4\ell$ decay channel at 13.6 TeV with a relative uncertainty of $\approx 20\%$ [26]. In addition, differential fiducial cross sections were measured as a function of the transverse momentum (p_T) and absolute value of the rapidity of the Higgs boson, denoted by p_T^H and $|y^H|$, respectively. This paper presents inclusive and differential measurements of the Higgs boson production in the $H \rightarrow \gamma\gamma$ channel from the CMS experiment at $\sqrt{s} = 13.6$ TeV. The measurements use data recorded with the CMS detector [27, 28] in 2022 and are performed in a fiducial phase space defined at the particle level with improved perturbative convergence of the theoretical calculations [29].

Although the $H \rightarrow \gamma\gamma$ decay has a branching fraction ($\mathcal{B}$) of $\approx 0.227\%$ [7], the excellent energy resolution of the CMS electromagnetic calorimeter (ECAL) yields a narrow peak in the invariant-mass distribution of the two photons, $m_{\gamma\gamma}$. The analysis follows the strategy used in previous measurements in this channel [9, 11, 30]. Events are selected with two reconstructed photons with $m_{\gamma\gamma} \in [100, 180]$ GeV, i.e. in a range around the Higgs boson mass of ≈ 125 GeV. The background and signal contributions are then estimated from a combined fit of a smoothly falling and a peaking function to the $m_{\gamma\gamma}$ distribution, reducing the reliance on Monte Carlo (MC) simulations for background modelling. The background function and its parameters are determined from the fit to data. The signal function is estimated from MC simulations corrected using $Z \rightarrow ee$ decays, taking advantage of the similarity of electron and photon electromagnetic showers. To improve the measurement sensitivity, events are categorized based on a per-event $m_{\gamma\gamma}$ resolution estimate. A combined fit to all categories is performed to extract the number of signal events. Compared to the previous measurement of fiducial inclusive and differential cross sections by the CMS Collaboration, which used 137 fb^{-1} of pp collision data at $\sqrt{s} = 13$ TeV [11], this analysis contains several improvements. In particular, a novel approach based on a neural network is used to correct the modelling of photon identification variables and the estimate of the per-photon energy resolution in MC simulations, so that their

distributions agree better with those observed in data.

The paper is structured as follows: the CMS detector is briefly introduced in Section 2. The data and simulation samples are described in Section 3. The event reconstruction, and the event selection and categorization are presented in Sections 4 and 5, respectively. The fiducial phase space and the observables for the differential measurements are introduced in Section 6. The statistical analysis and the systematic uncertainties are described in Sections 7 and 8, respectively. The results are presented in Section 9, followed by a summary in Section 10. Tabulated results are provided in the HEPData record for this analysis [31].

2 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal ECAL, and a brass and scintillator hadron calorimeter, each composed of a barrel and two endcap sections. The ECAL consists of 75 848 lead tungstate crystals, which provide coverage in pseudorapidity $|\eta| < 1.48$ in the barrel region (EB) and $1.48 < |\eta| < 3.0$ in the two endcap regions (EE). Preshower detectors consisting of two planes of silicon sensors interleaved with a total of three radiation lengths of lead are located in front of each EE detector. Forward calorimeters extend the pseudorapidity coverage provided by the barrel and endcap detectors. Muons are reconstructed using gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid.

Events of interest are selected using a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about $4\ \mu\text{s}$ [32]. The second level, known as the high-level trigger (HLT), consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 5 kHz before data storage [33, 34].

A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Refs. [27, 28].

3 Data samples and simulated events

This analysis uses pp collision data, collected in 2022 at $\sqrt{s} = 13.6\ \text{TeV}$, corresponding to an integrated luminosity of $34.7\ \text{fb}^{-1}$ [35]. The data were selected using a diphoton HLT [33] with p_T thresholds on the highest p_T (leading, γ_1) and second-highest p_T (subleading, γ_2) photon of 30 and 22 GeV, respectively. Additionally, it is required that $m_{\gamma\gamma} > 90\ \text{GeV}$. Both photons were required to pass loose identification criteria on the amount of surrounding energy deposits in the calorimeters and on variables that characterize the photon shower in the ECAL (shower shape variables) [36].

The four main Higgs boson production modes are gluon-gluon fusion (ggH), vector boson fusion (VBF), associated production with a W or Z boson (VH), and with a top quark pair (ttH). Samples for these processes are generated with MADGRAPH5_aMC@NLO (version 2.9.9) [37] at next-to-leading order (NLO) in perturbative quantum chromodynamics using the NNPDF3.1 NNLO [38] set of parton distribution functions (PDFs). The FxFx merging scheme [39] is used to match jets from matrix element calculations to those from parton shower for the ggH, VH, and ttH production processes. Events in the ggH production mode are weighted to match the predictions from the NNLOPS generator [40–42] as a function of the Higgs boson p_T and

the number of jets in the event. The samples are normalized to the cross sections provided by the LHC Higgs Working Group for $\sqrt{s} = 13.6$ TeV [43], based on an interpolation procedure using cross sections computed for the centre-of-mass energies of 13 and 14 TeV [7, 43], using the results from Refs. [44–65].

The main backgrounds are the irreducible background from non-resonant diphoton ($\gamma\gamma$) production and the reducible background from $\gamma + \text{jet}$ production, where a jet is misreconstructed as a photon. Multijet production accounts for an additional and smaller contribution in the reducible background. For the measurement of the inclusive and differential cross sections, the normalization and the shape of the $m_{\gamma\gamma}$ background distribution are estimated from the data. Simulated events for $\gamma\gamma$ and $\gamma + \text{jet}$ production are only used to optimize the event selection and categorization. Diphoton production is simulated with SHERPA (version 2.2.12) [66] at leading order (LO). The gluon-induced box process is simulated without additional final-state partons in the matrix element, whereas the quark-induced and quark-gluon-induced Born processes include up to three additional partons in the final state. The production of $\gamma + \text{jet}$ events is simulated with PYTHIA8 [67] (version 8.306) as a $2 \rightarrow 2$ LO process at the matrix-element level.

Several corrections are applied to the simulation samples or to the data to either assess and address mismodelling in the simulation or to improve the data calibration. These corrections are inferred from a comparison of data and simulation in events with $Z \rightarrow ee$ and $Z \rightarrow \mu\mu\gamma$ decays and are validated with these samples. In data, $Z \rightarrow ee$ events are collected using a single-electron trigger with a p_T threshold of 30 GeV and a double-electron trigger with p_T thresholds of 23 and 12 GeV for the leading and subleading electron, respectively [33, 34, 36]. Events with a radiative $Z \rightarrow \mu\mu\gamma$ decay are recorded with dimuon triggers with p_T thresholds of 17 and 8 GeV for the leading and subleading muon, respectively [33, 34, 68]. The simulation samples used for these corrections are generated with MADGRAPH5_aMC@NLO.

All simulated events are interfaced with PYTHIA for the simulation of the parton shower, fragmentation, and hadronization, using the CP5 underlying event tune [69]. The PYTHIA generator is also used to simulate additional pp interactions occurring in the same or neighbouring bunch crossings (pileup). Simulated events are reweighted to reproduce the distribution of the number of interaction vertices observed in the data. The average number of interactions per bunch crossing in 2022 data is 46, assuming a total inelastic pp cross section of 80 mb at $\sqrt{s} = 13.6$ TeV. For all processes, the response of the CMS detector is simulated using a detailed description of the CMS apparatus based on the GEANT4 package [70].

4 Event reconstruction

The primary vertex is taken as the vertex corresponding to the hardest scattering in the event, which is defined as the vertex that maximizes the p_T^2 sum of reconstructed particles, evaluated using tracking information alone [71]. A particle-flow (PF) algorithm [72] aims to reconstruct and identify each individual particle in an event (PF candidate), with an optimized combination of information from the various elements of the CMS detector.

Following a power cooling issue in September 2022, about 7% of the ECAL channels in one of the EEs were disabled [73]. Events with at least one jet with $p_T > 30$ GeV in this region are removed to prevent biases that could affect the cross section measurements reported in this paper. Simulation samples are split into two periods to reflect the detector conditions before and after the issue and are weighted according to the integrated luminosities of the corresponding data sets.

4.1 Photon reconstruction and identification

Energy deposits (clusters) in the ECAL form the basis of the photon reconstruction [36]. The ECAL clusters with energy well above the electronics noise level are combined if they are compatible with originating from the same photon, resulting in a so-called supercluster. Superclusters are selected as photons not matched to charged-particle trajectories associated with a reconstructed hard-scattering vertex. Photons in the transition regions between the barrel and the endcaps ($1.4442 < |\eta| < 1.5660$) are not considered.

An energy resolution of $\approx 1\%$ is achieved in the EB for unconverted or late-converting photons in the tens of GeV energy range. The energy resolution for other photons is $\approx 1.3\%$ up to $|\eta| = 1$ and $\approx 2.5\%$ up to $|\eta| = 1.44$ in the EB. In the EE, the energy resolution is $\approx 2.5\%$ for unconverted or late-converting photons, and between 3 and 4% for other photons.

Not all the photon energy is deposited in the operational ECAL crystals or accounted for in the clustering. This is mainly due to lateral and longitudinal shower leakage, intermodule gaps, unresponsive channels, or energy thresholds used to mitigate the noise in the cluster reconstruction. To improve the energy measurement, a dedicated simulation-based parametric regression is used. This method also provides a per-photon estimate for the energy resolution, σ_E . The input variables to the regression include the location of the supercluster and its seed (the crystal with the highest energy), and a range of shower shape variables. The relative energy resolution of photons with generated transverse momenta in the range of 1 to 100 GeV and with $|\eta| < 1.0$ improves from about 1.3 to 0.9% when the regression is applied. In the outer EE ($2.0 < |\eta| < 2.5$), the relative resolution for photons in the same momentum range improves from approximately 3.1 to 2.4%.

Differences between data and simulation in the energy scale and resolution are addressed by residual corrections. Electrons from $Z \rightarrow ee$ decays that are reconstructed and calibrated with the photon algorithms described above, except for the track-based veto, are used. The corrections are derived from a comparison of the dielectron invariant mass distribution in data and simulation [74]. The scale calibrations are applied to data and are derived from the positions of the Z boson peak in data and simulation as functions of time, the gain from the photodetector readout of the supercluster's seed crystal, as obtained from its multigain preamplifier [75], the photon energy, the supercluster η , and the R_9 variable. The R_9 variable is defined as the energy deposited in the three-by-three crystal matrix around the seed divided by the total uncorrected energy of the supercluster [36] and is useful to discriminate converted from unconverted photons. The resolution corrections broaden the width of the distribution in simulation, such that it corresponds to that observed in the data, and depend on the photon energy, the supercluster η , and R_9 .

A boosted decision tree (BDT) is trained to define photon identification (ID) criteria [73] in a similar way as described in Ref. [36]. The BDT score is used to separate prompt photons that are produced at the primary vertex and non-prompt photons, which mostly stem from collimated diphoton decays of neutral mesons, such as π^0 and η , inside hadronic jets. As input, the model uses shower shape variables, isolation variables built from the p_T of tracks and PF objects near the photon candidate, and the ratio of the energy deposited in the hadronic calorimeter behind the photon supercluster to the supercluster energy (H/E). The BDT is implemented in XG-BOOST [76] and is trained on prompt and non-prompt photons from simulated $\gamma + \text{jet}$ events. Besides the discriminant features, the energy and η of the photon supercluster as well as ρ , defined as the median of the transverse energy density in the event, are also used as input features. The BDT scores of prompt and non-prompt photons from simulated $\gamma + \text{jet}$ events are shown in Fig. 1 for reconstructed photons with $p_T^\gamma > 25 \text{ GeV}$. The leading and the subleading




Figure 1: Normalized distributions of the photon identification BDT scores for prompt (blue) and non-prompt (orange) photons from $\gamma + \text{jet}$ simulated events. The shaded region indicates the photons that are rejected by the photon preselection requirement of > -0.9 .

photon are included in the distributions. Prompt photons accumulate at high values of the BDT score, whereas the distribution of non-prompt photons is steeply falling towards larger values. The photon preselection used in this analysis rejects photons with a BDT score smaller than -0.9 , as indicated by the shaded region. This requirement is highly efficient for selecting prompt photons and rejects a large fraction of non-prompt photons, reducing the number of possible diphoton pairs.

Electron background contributions are reduced with a conversion-safe electron veto algorithm [77]. This veto rejects the photon candidate if its supercluster is close to a track compatible with an electron, unless the track is matched to a photon conversion vertex.

4.2 Corrections to simulated photons

Photon mismodelling in the simulation is a non-negligible source of systematic uncertainty in $H \rightarrow \gamma\gamma$ fiducial cross section measurements. This holds in particular for shower shape and isolation variables [9], but also for the per-photon energy resolution estimate, σ_E . Sources of the mismodelling include the imperfect description of the material budget in the simulation, the modelling of noise, and the time evolution of the detector response, in particular due to radiation damage [11, 36]. Shower shape and isolation variables, as well as σ_E , are hence corrected using probe electrons reconstructed as photons ($\gamma_{\text{probe},e}$) with the tag-and-probe method [78] in $Z \rightarrow ee$ events. A first correction method, called “chained quantile-regression”, was developed in Ref. [11] for the analysis of the Run 2 data. It involves the training of a large number of BDTs and was used to successfully correct shower shape and isolation variables.

In this analysis, a new method based on normalizing flows [79] is used for a more efficient derivation of the corrections. It takes as input the variables that are used for the photon identification BDT, as well as the per-photon estimate of the energy resolution, and provides as output per-photon corrections for all of these variables. Hence, the method does not only cor-

rect these values but also their correlations. The basis of the method is that normalizing flows are able to learn a high-dimensional mapping from a distribution of interest, in our case the BDT input variables and σ_E , to a simpler distribution of same dimension, often a multivariate Gaussian. We use a simple but efficient solution with one normalizing flow [80] trained on both data and simulation. It is parametrized as a function of an MC/data binary variable, as well as the p_T , η , and the azimuthal angle ϕ of the $\gamma_{\text{probe},e}$, and ρ . To perform the correction, the mapping to the multivariate Gaussian is performed for simulated photons, the MC/data boolean is flipped, and the inverse transformation provides the corrected values. The model is an autoregressive normalizing flow [81] and consists of five neural spline transformations [82] with ten spline bins, implemented using the PYTORCH [83] and ZUKO [84] libraries.

Before deriving these corrections, the distributions of the conditional variables ρ , p_T , η , and ϕ are reweighted in simulation to match the data distributions. In addition, the isolation variables, which show a discontinuous behaviour with a peak at zero followed by a continuous tail due to p_T thresholds on PF candidates and energy depositions in the calorimeters, are transformed to a continuous distribution [80]: isolation values in the peak are resampled to populate the gap between the peak and the start of the continuous tail.

The level of agreement between selected distributions in data and simulation for $\gamma_{\text{probe},e}$ from $Z \rightarrow ee$ decays before and after the corrections is shown in Fig. 2. The distributions for σ_E and for H/E are shown, where H/E serves as an example for the input variables that are used in the identification BDT with the largest observed data-to-simulation shape disagreement between data and MC. In addition, the BDT score is shown separately for photons in the EB and in the EE. After the corrections, the agreement of the BDT score between data and simulation is significantly improved. The average disagreement is as small as 1.7 (2.0)% in the EB (EE), where the average is calculated from the absolute difference between data and MC in all bins shown in Fig. 2 (lower row). The corrections are validated using photons from $Z \rightarrow \mu\mu\gamma$ events. Data and simulations are found to agree within the uncertainties for the BDT score and σ_E variables, which enter the event selection and categorization.

4.3 Lepton reconstruction

Electrons and muons are used for the derivation and validation of corrections in $Z \rightarrow ee$ and $Z \rightarrow \mu\mu\gamma$ events and for overlap removal procedures to resolve ambiguities between reconstructed objects. The reconstruction of electrons is based on charged-particle tracks matched to ECAL clusters [36]. Superclusters are built in the same way as for photons. The electron momentum is estimated from a combination of the ECAL energy and the momentum measurement in the tracker. Electrons are required to have $p_T > 15 \text{ GeV}$ and have to satisfy loose cut-based identification requirements with a signal efficiency of $\approx 90\%$. The variables used to define these identification criteria are described in Ref. [36]. Muons are reconstructed using a combination of a track in the central tracking system and either a single track or multiple hits in the muon detectors [85]. Muons are required to have $p_T > 10 \text{ GeV}$ and they need to pass tight identification criteria, which are based on the relative isolation with respect to hadrons and photons in a cone of radius $\Delta R = 0.4$, where $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ defines the angular distance, around the muon and the quality of the fit of the muon track. Electrons and muons have to be separated by an angular distance greater than 0.2 from both of the photons of the diphoton pair, which is reconstructed as described in Section 5.1, when they are used for overlap removal. No requirement on the angular distance to photons is placed when electrons and muons are used for the derivation of corrections in $Z \rightarrow ee$ and $Z \rightarrow \mu\mu\gamma$ events.




Figure 2: Data-to-simulation comparison for σ_E (upper left), H/E (upper right), the photon identification BDT score in EB (lower left) and EE (lower right) for electrons from $Z \rightarrow ee$ decays reconstructed as photons. The uncorrected distributions are shown in blue and the corrected distributions from the normalizing flow are shown in green. The error bars in the ratio panels include the statistical uncertainty from the data and the uncertainty from the limited number of simulated events. For the distributions of the photon identification BDT score, the shaded region corresponds to photons with a BDT score < 0.25 , which are excluded by the selection applied in the cross section measurements. For the σ_E distribution, the last bin contains the overflow.

4.4 Jet reconstruction

In this analysis, jets are only used for the differential measurement with respect to the number of jets. Jets are reconstructed from PF candidates using the anti- k_T algorithm [86, 87] with a distance parameter of 0.4. The pileup-per-particle identification algorithm [88] is used to mitigate pileup effects. The algorithm assigns a weight to each particle candidate before the jet clustering according to the likelihood that the candidate originated from pileup. Jets originating from noise and reconstruction failures are rejected using criteria on the energy composition and number of PF constituents of the jets [88].

Jet energy corrections (JECs) are derived from simulation to calibrate the measured jet momentum to that of particle-level jets [89]. In situ measurements in data of the momentum balance in dijet, $\gamma + \text{jet}$, $Z + \text{jet}$, and multijet events are used to account for residual differences in the jet energy scale between data and simulation. The jet energy resolution (JER) is found to be worse in data than in simulation. The resolution in the simulation is hence broadened to agree with that observed in data.

Jets are required to have $p_T > 30 \text{ GeV}$ and $|\eta| < 2.5$, as jets in the central part of the detector are subject to lower systematic uncertainties than in the forward region. Whereas the requirement on $|\eta|$ reduces the efficiency for the VBF process, inclusive Higgs boson production is dominated by the ggH process in the SM. Jets with $\Delta R < 0.4$ from a photon of the diphoton pair or a charged lepton are removed.

5 Event selection and categorization

5.1 Event selection

The event selection retains $H \rightarrow \gamma\gamma$ signal candidates while rejecting as much background as possible from both prompt and non-prompt photons. Each photon must have a supercluster with $|\eta| < 2.5$, excluding the ECAL barrel-endcap transition regions of $1.4442 < |\eta| < 1.5660$. Each photon must satisfy preselection criteria based on its shower shape, isolation and kinematic properties. These criteria are described in detail in Ref. [90] and they are defined to be slightly more stringent than the corresponding trigger requirements. In particular, the requirements $p_T^{\gamma_1} > 35 \text{ GeV}$ and $p_T^{\gamma_2} > 25 \text{ GeV}$ are applied to the p_T -leading and -subleading photons, respectively. Only the p_T -leading diphoton system composed of photons satisfying the preselection requirements is considered. The selection criteria for $p_T/m_{\gamma\gamma}$ are changed with respect to previous measurements [11, 91]. Previously, the leading (subleading) photon had to fulfil $p_T/m_{\gamma\gamma} > 1/3$ ($1/4$). In this analysis, the requirement for the subleading photon is unchanged, but a requirement of $\sqrt{p_T^{\gamma_1} p_T^{\gamma_2}}/m_{\gamma\gamma} > 1/3$ is applied instead of the above criterion for the leading photon. This requirement on the scaled geometric mean of $p_T^{\gamma_1}$ and $p_T^{\gamma_2}$ improves the perturbative convergence of the theoretical calculations [29].

The efficiencies of the diphoton trigger are measured with $Z \rightarrow ee$ events using the tag-and-probe method, from which corrections (“scale factors”) are derived for simulated events such that the efficiencies in simulation match those measured in the data [33]. The measurement is performed in bins of p_T , η , and R_9 . Scale factors for the preselection efficiencies and for the photon identification efficiency after the application of the normalizing flow correction are measured in a similar way, whereas the scale factors for the efficiencies of the electron veto criterion are computed from $Z \rightarrow \mu\mu\gamma$ events.

5.2 Event categorization

In order to maximize the sensitivity to the Higgs boson signal and minimize the dependence on the underlying model for its production and decay, the selected events are categorized based on σ_m/m [9], i.e. the estimator of the per-event diphoton invariant-mass resolution divided by $m_{\gamma\gamma}$. The resolution estimator σ_m is calculated from the per-photon resolution estimator σ_E obtained from the energy regression BDT (cf. Section 4.1) for each photon:

$$\frac{\sigma_m}{m} = \frac{1}{2} \sqrt{\left(\frac{\sigma_{E_1}}{E_1}\right)^2 + \left(\frac{\sigma_{E_2}}{E_2}\right)^2}, \quad (1)$$

where the contribution from the photon angles is neglected. The contribution to the mass resolution from the angular resolution is negligible with respect to the one from the energy resolution if the chosen Higgs boson decay vertex has $|\Delta z| < 1$ cm, with Δz being the difference of the z -coordinate between the true and the reconstructed vertex. This is referred to as the correct identification of the vertex, which occurs for $\approx 70\%$ of ggH events. The efficiency of the correct vertex identification increases with higher Higgs boson p_T and exceeds 85 (95)% for VH ($t\bar{t}H$) associated production across the entire range.

Since the BDT is trained on simulation and the energy resolution is known to be worse in data than in simulation, the term that adjusts the energy resolution in simulation (cf. Section 4.1) is added in quadrature to the value of σ_E from the energy regression BDT before σ_m/m is calculated.

The diphoton mass resolution in $H \rightarrow \gamma\gamma$ decays is typically 1–2%, depending on the measurement of the photon energies in the ECAL and the topology of the photons in the event [74]. As the relative energy resolution σ_E/E improves with photon energy, σ_m/m is correlated with the diphoton invariant mass. Hence, a categorization in σ_m/m can distort the $m_{\gamma\gamma}$ distribution of the background by depleting the low-mass region in high-resolution categories. The background distribution, instead, is assumed to be monotonically falling over the selected diphoton mass range in the statistical analysis. To avoid distortions of the invariant mass distribution, the mass-resolution estimator is decorrelated from the diphoton invariant mass. This is achieved with the quantile morphing algorithm described in Ref. [11]. It divides the σ_m/m distribution into bins of $m_{\gamma\gamma}$ and morphs the cumulative distribution function of σ_m/m in each bin to resemble that in a reference mass bin, chosen as $[125.0, 125.5]$ GeV.

The decorrelated mass-resolution estimator is used to categorize the events. Additionally, a requirement on the photon ID BDT score is placed on both photons to reject background contributions with non-prompt photons. This requirement and the category boundaries are optimized simultaneously using simulated events. The signal component is modelled with a combination of the four main production modes with a Higgs boson mass of 125 GeV, whereas the non-resonant background is composed of the diphoton and $\gamma + \text{jet}$ MC samples. Both the signal and background components are weighted according to the expected number of events in the selected phase space, assuming the SM cross sections for the Higgs boson production processes at $\sqrt{s} = 13.6$ TeV [43] and the LO cross section predictions for the background processes. Additional scaling factors for the normalizations of the diphoton and $\gamma + \text{jet}$ processes are determined from a fit to the distribution of the lower of the two photon ID BDT scores, restricted to events where the minimum score exceeds zero, in order to improve the modelling of the invariant mass distribution. The $m_{\gamma\gamma}$ distribution of the signal and background samples is fitted using the sum of an exponential function for the background and the sum of two Gaussian distributions for the signal. The figure of merit for the optimization is the approximate expected signal significance $S/\sqrt{B}$, where S and B are the expected number of signal and back-




Figure 3: Data-to-simulation comparison of the per-event decorrelated mass-resolution estimator σ_m/m using $Z \rightarrow ee$ events. Both electrons are reconstructed as photons and categorized either both in the EB (left) or at least one in the EE (right). The uncertainty band in the lower panel represents the systematic uncertainty based on the residual mismodelling of σ_E/E (5%). The error bars on the markers in the lower panels include the statistical uncertainty from data and the uncertainty from a limited number of simulated events. The last bin contains the overflow.

ground events, respectively, in the smallest $m_{\gamma\gamma}$ interval containing 68% of simulated signal events.

The modelling of the σ_m/m distribution in simulation is improved by including the per-photon energy resolution estimator σ_E into the set of variables corrected with the normalizing flow (cf. Section 4.2). Figure 3 shows the agreement between the σ_m/m distributions in $Z \rightarrow ee$ data and simulation before and after propagating the corrections for σ_E to σ_m/m . Both electrons in the $Z \rightarrow ee$ events are reconstructed as photons and the simulated events are reweighted to data in p_T , η , and ϕ of $\gamma_{\text{probe},e}$, as well as in ρ . The corrected MC distribution agrees with the distribution observed in data within the assigned systematic uncertainty in σ_E/E (5%), as described in Section 8.

Three σ_m/m categories are used in this analysis, as the improvement in sensitivity quickly saturates and does not noticeably improve with additional categories. The resulting category boundaries are $[0, 0.0105)$, $[0.0105, 0.0130)$, and $[0.0130, \infty)$. In all three categories, the minimum value for the photon ID BDT score is 0.25.

The combined acceptance and efficiency of the event selection described in this section is estimated from simulation to be $\approx 32\%$ for the ggH and VBF processes, about 27% for the VH process, and $\approx 28\%$ for the $t\bar{t}H$ process with respect to the total phase space.

6 Fiducial phase space and observables

The cross section for the Higgs boson production is measured in a fiducial phase space defined at the particle level, with the goal of reducing model dependence and extrapolation uncertainties. The fiducial criteria at the particle level are close to the event selection requirements at the detector level (Section 5). The criteria are based on the two p_T -leading photons in the event.

They must be within the fiducial acceptance of $|\eta| < 2.5$ with $1.4442 < |\eta| < 1.5660$ excluded to match the rejection of the ECAL barrel-endcap transition regions at the detector level. The photons must fulfil $\mathcal{I} < 10 \text{ GeV}$, where the isolation variable $\mathcal{I}$ is the scalar p_T sum of stable, visible final-state particles in a cone of radius 0.3 centred on the photon momentum direction. The geometric requirement of $\sqrt{p_T^{\gamma_1} p_T^{\gamma_2}} / m_{\gamma\gamma} > 1/3$ is applied, and the subleading photon must fulfil $p_T^{\gamma_2} / m_{\gamma\gamma} > 1/4$. The efficiency of these criteria, as determined from simulation, is $\approx 50.6\%$. This is slightly lower compared to the efficiency of $\approx 51.8\%$ that is obtained with the requirement $p_T^{\gamma_1} / m_{\gamma\gamma} > 1/3$ used in previous measurements [11, 30] instead of the geometric requirement.

Fiducial cross sections are also measured in bins of four kinematic observables: the transverse momentum, the absolute value of the rapidity of the diphoton system, the number of jets (N_{jets}) in the event, and the transverse momentum of the leading jet ($p_T^{\text{j}_1}$), defined as the jet with the highest p_T . Jets at the fiducial level are built with the anti- k_T clustering algorithm out of stable particles with a distance parameter of 0.4, excluding neutrinos. Jets are retained if they satisfy $p_T^{\text{jet}} > 30 \text{ GeV}$, $|\eta^{\text{jet}}| < 2.5$ and if they do not overlap with an electron or muon within $\Delta R < 0.4$, where the leptons have to fulfil the following requirements: Electrons (muons) must have $p_T > 15$ (10) GeV, $|\eta| < 2.5$ (2.4), and $\mathcal{I} / p_T < 0.2$. These criteria match the detector-level jet selections. The bin boundaries are chosen to provide an expected relative uncertainty in each bin of about 40 (60%) for p_T^H and $|y^H|$ (N_{jets} and $p_T^{\text{j}_1}$). The boundary values are shown in Table 1.

Table 1: Bin boundaries for the differential cross section measurement. The first $p_T^{\text{j}_1}$ bin corresponds to events without jets. For the N_{jets} binning, the right boundary should be considered as not included in the bin, i.e. [lower, upper).

Observable	Bin boundaries								
p_T^H (GeV)	0	15	30	45	80	120	200	350	∞
$ y^H $	0	0.15	0.3	0.6	0.9	2.5			
N_{jets}	0	1	2	3	∞				
$p_T^{\text{j}_1}$ (GeV)	0-jet	30	75	120	200	∞			

7 Statistical analysis

To extract the inclusive fiducial cross section, σ_{fid} , a binned profile likelihood fit to the $m_{\gamma\gamma}$ distributions in the three mass-resolution categories is performed with the COMBINE tool [92]. A bin width of 0.25 GeV has been chosen, which is sufficiently small compared to the typical mass resolution of 1.5–2 GeV of the SM Higgs boson signal as indicated in Fig. 4. Systematic uncertainties are treated as nuisance parameters with Gaussian constraints. The likelihood can be factorized over the $N_m = 320$ bins of the $m_{\gamma\gamma}$ distributions in each category:

$$\mathcal{L}(\sigma_{\text{fid}}, \vec{n}_{\text{bkg}}, \vec{\theta}_S, \vec{\theta}_B) = \text{Pdf}(\vec{\theta}_S) \text{Pdf}(\vec{\theta}_B) \prod_{i=1}^{N_{\text{cat}}} \text{Pois} \left(n_{\text{ev}}^i | n_{\text{sig}}^i + n_{\text{bkg}}^i \right) \\ \times \prod_{l=1}^{N_m} \left(\frac{\sigma_{\text{fid}} K^l(\vec{\theta}_S) S^l(m_{\gamma\gamma}^l | \vec{\theta}_S) L + n_{\text{OOA}}^i S_{\text{OOA}}^l(m_{\gamma\gamma}^l | \vec{\theta}_S) + n_{\text{bkg}}^i B^l(m_{\gamma\gamma}^l | \vec{\theta}_B)}{n_{\text{sig}}^i + n_{\text{bkg}}^i} \right)^{n_{\text{ev}}^i}, \quad (2)$$

where

- $\text{Pdf}(\vec{\theta})$ is the probability density function for the vector of nuisance parameters $\vec{\theta}$;
- $\vec{\theta}_S$ and $\vec{\theta}_B$ are the vectors of nuisance parameters associated with the signal and the background models, respectively;
- $\text{Pois}(n|\lambda)$ is the probability mass function of the Poisson distribution for n occurrences with an expectation value of λ ;
- n_{ev}^i is the number of observed data events in category i and n_{ev}^{li} is the number of observed data events in category i and $m_{\gamma\gamma}$ bin l ;
- $n_{\text{sig}}^i(\sigma_{\text{fid}}, \vec{\theta}_S) = \sigma_{\text{fid}} K^i(\vec{\theta}_S) L + n_{\text{OOA}}^i$ and n_{bkg}^i are the number of signal and background events in category i , and $\vec{n}_{\text{bkg}}$ is the vector with entries n_{bkg}^i ;
- $N_{\text{cat}} = 3$ is the number of mass-resolution categories;
- $K^i(\vec{\theta}_S)$ is the efficiency for reconstructing an event in category i ;
- $S^i(m_{\gamma\gamma}|\vec{\theta}_S)$ and $B^i(m_{\gamma\gamma}|\vec{\theta}_B)$ are the signal and background probability density functions in category i (cf. Sections 7.1 and 7.2);
- $m_{\gamma\gamma}^l$ is the centre of the l -th $m_{\gamma\gamma}$ bin;
- L is the integrated luminosity;
- $n_{\text{OOA}}^i S_{\text{OOA}}^i(m_{\gamma\gamma}|\vec{\theta}_S)$ is the out-of-acceptance (OOA) signal contribution in category i , where n_{OOA}^i is the number of OOA events and S_{OOA}^i the corresponding $m_{\gamma\gamma}$ signal model. n_{OOA}^i and S_{OOA}^i are estimated from simulation and are affected by the same set of nuisance parameters as in-fiducial signal events.

A similar approach is used to extract the differential cross sections. In those fits, σ_{fid} is promoted to a vector of fiducial cross sections measured in the particle-level bins of a specific observable. At the detector level, the same binning is used. Consequently, all quantities carrying the index i in Eq. (2) are extended with an additional index j to enumerate the same number of reconstruction-level bins. Similarly, several quantities are extended with an index k for the respective particle-level bin. For example, the per-category efficiency $K^i(\vec{\theta}_S)$ becomes the detector response matrix $K_k^{ij}(\vec{\theta}_S)$ that relates the events from a particle-level bin k with a detector-level bin j and category i . This permits to encode the unfolding directly in the likelihood. The condition numbers for the response matrices are less than ten, so no regularization is performed. The condition number is defined as the absolute value of the ratio between the largest and smallest matrix eigenvalues.

7.1 Signal model

The signal model is based on simulated events, after including all the corrections described before. A sum of up to five Gaussian functions is used as a parametric model to describe the $m_{\gamma\gamma}$ resonant shape for each combination of reconstruction-level bins and σ_m/m categories, separately for events passing the fiducial selection and the OOA contribution for each of the four main SM Higgs boson production modes. The fraction of OOA events is the largest for the VH and ttH production processes, reaching up to 3.4% in the worst-resolution category for the measurement of the inclusive cross section. In the best-resolution category, the fraction of OOA events is below 1% for the ggH, VBF, and VH processes. Additionally, different signal shapes are constructed for the cases of correct and incorrect identification of the primary vertex, as the invariant diphoton mass depends on the estimated photon direction and therefore the $m_{\gamma\gamma}$ shape differs significantly for the cases of a correctly or incorrectly identified primary vertex. The models for the correct and incorrect vertex identification are combined with their

relative weights obtained from simulation. This enables the inclusion of systematic uncertainties related to the vertex assignment in the fit (cf. Section 8). To improve the modelling of the luminous region, its spread along the beam axis is reweighted in simulation to correspond to the spread in data. The final nominal signal model is given by the weighted sum of the individual parametric models according to the respective SM cross section predictions, OOA fractions, and selection efficiencies. The combined parametric signal models for the three mass-resolution categories and the weighted sum across all categories are shown in Fig. 4. The values of the effective mass resolution σ_{eff} , defined as half of the smallest interval centred around the mean that contains 68.3% of the total area under the signal model histogram, are indicated. Signal shapes are constructed for Higgs boson masses of 120, 125 and 130 GeV. The parameters of the models for masses between these points are obtained using piecewise linear interpolation. A piecewise cubic interpolation is used to determine the $\mathcal{B}(\text{H} \rightarrow \gamma\gamma)$, the acceptance and efficiency, and the right-vertex fraction as a function of the Higgs boson mass based on the same mass hypotheses. The nominal Higgs boson mass is set to 125.38 GeV [74]. The interference between the $\text{H} \rightarrow \gamma\gamma$ signal and the continuous diphoton background [93] is not taken into account as it is at the percent level for an SM Higgs boson, which is well below the theoretical uncertainties.

7.2 Background model

The background model is defined in a data-driven way using the discrete profiling method [94]. Parametrized functions are used to model the smoothly falling background as a function of $m_{\gamma\gamma}$ in the range 100–180 GeV. A different model is constructed for each category. During the extraction of the fiducial cross sections from the fit to data (cf. Eq. (2)), the choice of the background function is treated as a discrete nuisance parameter and accounts for the imperfect a-priori knowledge of the $m_{\gamma\gamma}$ background shape. The resulting uncertainty is absorbed into the statistical uncertainty.

Initial fits are performed over the whole $m_{\gamma\gamma}$ range to determine the set of plausible function choices. The considered functions are grouped into families: sums of Bernstein polynomials, sums of exponential functions, Laurent series, and sums of power-law functions. In each family, multiple functions with different numbers of parameters are considered. An F -test [95] is performed to determine the maximum number of parameters to be used, whereas the minimum number is determined by a requirement on the goodness-of-fit to the data. A higher number of degrees of freedom for the fitting function is penalized by constructing the likelihood using $-2 \ln \mathcal{L}_B + N_B$, where $\mathcal{L}_B$ refers to the likelihood as a function of the background function parameters and N_B is the number of free parameters of a given background function. In the profile likelihood fit, at least four background functions are considered for each category with at least one function per family. In the measurement of the inclusive cross section, the sum of three exponential functions, the Bernstein polynomial sum of fourth degree, and the sum of three exponential functions are the best-fit functions in the best-resolution, medium-resolution and worst-resolution categories, respectively. For the measurement of the differential cross sections, a best-fit background function is determined for the three mass-resolution categories in each detector-level bin of the differential distribution.

8 Systematic uncertainties

The following systematic uncertainties affect the signal model by allowing changes in the location and the width of the Higgs boson peak (detailed in Section 7.1):

- *Photon energy scale and resolution*: This is the uncertainty related to the scale of the




Figure 4: Combined parametrized signal shapes per category and for the sum of all categories for the measurement of the inclusive cross section. The open squares denote the expectation from the simulation and the blue lines show the parametric models that describe the simulations. The uncertainty bars for the expectation from the simulation due to the limited number of simulated events are smaller than the marker size. The normalization of the histograms corresponds to the expected number of events, taking into account the cross sections of the considered production modes, the efficiency of the selection, and the integrated luminosity of 34.7 fb^{-1} . The effective mass resolution σ_{eff} (defined as half of the width of the smallest interval containing 68.3% of the area of the distribution) for each combined signal model is indicated in the grey area.

photon energy in data and the resolution corrections in simulation. The total uncertainty consists of three components: First, a uniform 0.1% uncertainty accounts for biases from the method related to the energy dependence of the resolution corrections to the simulation, efficiency scale factors, and kinematic differences between data and simulation. Additionally, the variations of the corrections from changing the fit window of the invariant dielectron mass from $[80, 100]$ to $[70, 110]$ GeV are considered. Furthermore, the derivation of the corrections is repeated with a photon ID BDT score requirement of > -0.9 instead of > 0.25 to estimate the impact of background contributions as well as the correlations between the energy and the photon ID BDT score. The energy scale uncertainties are the lowest for photons in the inner EB ($|\eta| < 1$) and amount to $\approx 0.1\%$ in this region. The relative uncertainty in the resolution term is around 5% for high- R_9 photons in the inner EB. It reaches 20 (40)% for high- (low-) R_9 photons in the EE. Six uncorrelated nuisance parameters parametrize this source of uncertainty in the statistical model.

- *GEANT4 electromagnetic shower modelling*: An uncertainty is introduced to account for the imperfect modelling of electromagnetic showers in GEANT4. A simulation with an alternative shower description modifies the energy scale for both electrons and photons. This results in an uncertainty of 0.05% in the photon energy scale.
- *Non-uniformity of light collection*: This uncertainty is related to the modelling of the light collection depending on the emission depth in the ECAL crystals, which is different for electrons and photons [74]. It is only considered for photons with $R_9 > 0.975$ as most photons with smaller R_9 values convert before reaching the calorimeter and therefore deposit most of their energy as electrons. This uncertainty affects the photon energy scale and amounts to about 0.14 (0.31)% for photons in the EB (EE).
- *Modelling of the material upstream of the ECAL*: The fraction of photons that convert before reaching the ECAL depends on the material before the ECAL whose description in the simulation is not perfect. The impact on the photon energy scale is estimated by varying the amount of upstream material in the simulation. It amounts to 0.02–0.05% for photons in the EB and reaches 0.25% for photons in the EE.
- *Vertex assignment*: The fraction of events with a reconstructed vertex within $|\Delta z| < 1$ cm of the true vertex, i.e. with the mass resolution driven by the energy resolution of the photons, is varied by $\pm 2\%$. The estimation of this uncertainty is detailed in Ref. [91].

Other sources of experimental systematic uncertainty affect the event yields. Their effect is parametrized with log-normal distributions. The following sources of experimental uncertainty are considered:

- *Integrated luminosity*: The uncertainty in the integrated luminosity collected in 2022 is 1.4% [35].
- *Cross section for pileup reweighting*: Simulated events are reweighted so that the distribution of the number of reconstructed primary vertices in MC matches that observed in data. The data pileup distribution is inferred from the recorded instantaneous luminosity profile together with an assumed inelastic pp cross section of 69.2 mb and compared to that in simulation to determine the nominal reweighting factors. Its uncertainty of 3.2 mb is propagated by recomputing the data pileup distribution, yielding alternative event weights that are propagated to the statistical analysis and parameterized with a single nuisance parameter. This uncertainty primarily affects the extracted cross section through the correlation between pileup and the diphoton

mass resolution estimator.

- *Trigger efficiency*: The systematic uncertainty in the trigger efficiency is determined by using alternative background templates for the efficiency measurement from $Z \rightarrow ee$ events. The uncertainty reaches 2.2% for low- R_9 photons, but it is generally below 0.3% for photons in the EB and with high values of R_9 , which is the case for most photons from Higgs boson decays that enter the best-resolution category.
- *Photon preselection efficiency*: The uncertainty in the photon preselection scale factor is evaluated by varying the signal and background shapes used to determine the preselection efficiency in data and simulation from $Z \rightarrow ee$ events and propagating these variations to the scale factors. The resulting uncertainty is generally below 1% for photons in the EB with $p_T < 60$ GeV, where the Z boson decays provide a large number of events. For photons in the EE, the uncertainty can be as large as 2.8%.
- *Electron veto efficiency*: $Z \rightarrow \mu\mu\gamma$ events are used to determine the efficiency of the electron veto in both data and simulation. The limited number of events in data is the dominant uncertainty in the resulting scale factor. The uncertainty reaches 2.5% for low- R_9 photons in the EE, but is below 0.5% otherwise.
- *Photon identification efficiency*: After applying the normalizing-flow-based corrections, the photon ID BDT score distribution is in good agreement between data and simulation. Scale factors are calculated using the tag-and-probe method with $Z \rightarrow ee$ events to correct the simulation for any remaining disagreement. The scale factors are compatible with unity within the uncertainties. Several sources of systematic uncertainty are taken into account, including alternative signal and background modelling templates. These uncertainties are combined and result in an uncertainty that is generally below 0.5% for photons in the EB. For EE photons with low R_9 , it can reach up to 2%.
- *Per-photon energy resolution*: After applying the normalizing-flow-based correction, a small residual disagreement in the distribution of σ_E/E between data and simulation remains for both $Z \rightarrow ee$ events as well as diphoton events with $m_{\gamma\gamma}/\text{GeV} \in [100, 120] \cup [130, 180]$. A conservative uncertainty of 5% is applied to σ_E/E and propagated to the invariant mass resolution of the diphoton system. This uncertainty mainly results in migrations between the mass-resolution categories.
- *Jet energy correction and jet energy resolution*: The uncertainty in the calibration of the JEC and JER [89] directly affects the selection efficiency of the jets for the differential measurement of the number of jets. The uncertainty in the JEC and JER is 2–5%, depending on the p_T and η of the jets.

In addition, several sources of theoretical uncertainty are taken into account. The normalization of each particle-level bin is fixed while evaluating the effect of all of these uncertainties except for those related to the limited number of events in the simulation samples. Thus, their effect only enters in category migrations between detector-level bins and mass-resolution categories. The following sources are considered:

- *Parton distribution function uncertainties*: The uncertainty in the imperfect knowledge of the PDFs is estimated by reweighting events according to the NNPDF3.1 [38] prescription using the compressed Hessian set of PDF eigenvectors. This source of uncertainty is parameterized with 100 independent nuisance parameters in the statistical model.
- *Renormalization and factorization scales uncertainty*: This uncertainty is related to the missing higher-order terms in the perturbation series for the cross section calcula-

tion. Simulated events are reweighted with alternative event weights where the scales are varied by a factor of two, excluding the $(2, 1/2)$ and $(1/2, 2)$ variations.

- α_S *uncertainty*: As the PDF uncertainty, the uncertainty in the value of the strong coupling constant is taken from the NNPDF3.1 prescription. It is evaluated by varying α_S by ± 0.002 from the nominal value of $\alpha_S(m_Z) = 0.118$.
- *Parton shower uncertainty*: The uncertainty in the modelling of the parton shower is estimated with reweighting factors that correspond to per-event cross section variations with the scales for initial- and final-state radiation varied up and down by a factor of two.

Finally, the uncertainty due to the limited number of events in the simulation samples is accounted for and parametrized using one nuisance parameter per category and for each of the two periods before and after the ECAL cooling issue [96, 97].

The dominant sources of systematic uncertainties for the measurement of the fiducial cross section are summarized in Table 2.

9 Results

The measured cross section in the fiducial phase space, defined in Section 6, is

$$\sigma_{\text{fid}} = 74 \pm 11 \text{ (stat)}_{-4}^{+5} \text{ (syst) fb} = 74 \pm 12 \text{ fb.} \quad (3)$$

Figure 5 shows the likelihood scans for the inclusive fiducial cross section measurement. It also shows the theoretical prediction of $67.8 \pm 3.8 \text{ fb} = 67.8 \pm 2.6 \text{ (scales)} \pm 2.3 \text{ (PDF} + \alpha_S) \pm 1.4 \text{ (}\mathcal{B}\text{) fb}$, calculated with MADGRAPH5_aMC@NLO reweighted to match the NNLOPS prediction for ggH, and interfaced with PYTHIA8 (version 8.240) [98] using the CP5 tune. The measured value agrees with this prediction within the uncertainties. The uncertainty in the prediction combines contributions from the Higgs boson production cross sections, the $\mathcal{B}(\text{H} \rightarrow \gamma\gamma)$, and the fiducial acceptance. The first two uncertainties are taken from Refs. [7] and [43], respectively, whereas the third is computed with the MC samples presented in Section 3. The variations of the fiducial acceptance are evaluated with the compressed Hessian eigenvector set of NNPDF3.1 [38], the α_S value varied by 0.002 around its nominal value of 0.118, and the renormalization and factorization scales varied by a factor of 2, while excluding the $(2, 1/2)$ and $(1/2, 2)$ variations.

The dominant sources of systematic uncertainty for the measurement of the inclusive fiducial cross section are shown in Table 2. The main contribution is from the per-photon energy resolution, related to migrations between the mass-resolution categories, followed by the photon energy scale and resolution, which affect the location and the width of the Higgs boson peak. Other experimental uncertainties not listed in Table 2 are below 0.5% and the theoretical uncertainty is below 1%.

The observed diphoton invariant mass distribution is shown in Fig. 6 together with the combined signal and background fit and the background component alone. The distributions from the three mass-resolution categories in the inclusive fiducial measurement are included and weighted by $S/(S+B)$, where S and B are the number of signal and background events in the central interval of width $2\sigma_{\text{eff}}$ around the fitted peak position per mass-resolution category.

The differential fiducial cross sections for the observables and the binning introduced in Section 6 are extracted from the maximum likelihood fit together with their uncertainties and the




Figure 5: Likelihood scans for the inclusive fiducial cross section measurement. The black line corresponds to considering both the statistical and systematic uncertainties. The blue dash-dotted line corresponds to considering only the statistical uncertainty, including the discrete profiling method for the background modelling uncertainty. The theoretical prediction from MADGRAPH5_aMC@NLO, including the NNLOPS reweighting for the ggH component, is shown in red. The shaded theory uncertainty band includes the uncertainties in the renormalization and factorization scales, in the parton distribution functions, in α_s , in the $\mathcal{B}(H \rightarrow \gamma\gamma)$, and in the fiducial acceptance.

correlation matrices. The spectra and correlation matrices for p_T^H , $|y^H|$, N_{jets} , and p_T^h are presented in Figs. 7, 8, 9, and 10, respectively. The cross sections measured in bins of jet-related observables exhibit stronger correlations due to the larger bin-by-bin migrations induced by the relatively poor energy resolution for jets compared to photons, whereas there is almost no correlation for the well measured variables p_T^H and $|y^H|$. The measured differential cross sections are compared to various theoretical predictions. For these theoretical predictions, the acceptances in the differential bins are calculated using the ggH predictions from three different generators, whereas the inclusive prediction is normalized to the next-to-next-to-next-to-leading order computation [7, 43]. The three predictions are taken from the MADGRAPH5_aMC@NLO simulation, with and without NNLOPS reweighting, and from the POWHEG 2.0 event generator [99–102]. The acceptance of the $xH = VBF + VH + t\bar{t}H$ component of the signal is taken from the MADGRAPH5_aMC@NLO simulation, whereas the inclusive prediction is normalized to the integrated cross sections reported in Ref. [43]. The uncertainties in the theoretical predictions are computed following the same strategy as for the inclusive cross section, described above.

To assess the compatibility with the SM predictions, p -values are computed for every observable from

$$p = \int_{\Delta}^{\infty} f(x; N_{\text{Bins}}) dx, \quad (4)$$

where $f(x; N_{\text{Bins}})$ is the probability density function for a chi-squared variable (χ^2) with N_{Bins} degrees of freedom and $\Delta = 2(\text{NLL}(\vec{\sigma}_{\text{SM}}) - \text{NLL}(\vec{\sigma})) = \chi_{\text{SM}}^2$ is twice the difference between




Figure 6: Diphoton invariant mass distribution in the inclusive fiducial measurement, weighted by $S/(S+B)$ for the different mass-resolution categories. The distribution is shown together with the signal+background fit (red line) and the background-only component (dashed line). In the lower panel, the signal component is shown, estimated by subtracting the background component from the signal+background fit. The green (yellow) bands indicate the $\pm 1\sigma$ ($\pm 2\sigma$) uncertainties in the background component. They are derived from pseudoexperiments using the best-fit background function from the signal+background fit.

Table 2: Magnitude of the systematic uncertainties (Impact) in the inclusive fiducial cross section measurement. The magnitude of the uncertainty from the photon energy scale and resolution is extracted by performing a fit with the corresponding group of nuisance parameters frozen to their best-fit values. The obtained confidence interval is then subtracted in quadrature from the total confidence interval from the fit where all nuisance parameters are profiled. The magnitudes of the other sources of systematic uncertainty are obtained by varying the corresponding nuisance parameter by one standard deviation, keeping the other nuisance parameters at their best-fit values.

Systematic uncertainty	Impact in %
Category migration from energy resolution	+3.5/−4.2
Photon energy scale and resolution group	+3.4/−2.8
Integrated luminosity	±1.4
Photon preselection efficiency	±1.4
Material budget	+1.3/−1.2
Photon identification efficiency	±1.0
Pileup reweighting	±0.8

the negative log likelihood (NLL) evaluated for the SM hypothesis and for the best-fit values. The p -values for the nominal prediction, MADGRAPH5_aMC@NLO with NNLOPS reweighting, are 0.14, 0.19, 0.85, and 0.65 for the measurements of the p_T^H , $|y^H|$, N_{jets} , and p_T^h differential cross sections, respectively. These values are above 5% and show good compatibility with the SM prediction.

The residual model dependence of the differential measurements is also tested. Differential fiducial cross sections are extracted from fits to an Asimov data set [103] that comprises the background component generated from the best-fit background modelling function and a signal component from the SM signal model. In these fits, the signal component is assumed to be entirely composed of VBF, VH, or $t\bar{t}H$. This introduces a model dependence, as the migration matrices do not correspond to the SM scenario and the normalization of the OOA component changes. The differences with the extracted differential cross sections using the SM signal component hypothesis are below 12% in every particle-level p_T^H bin. The average deviation is 8.3%, which is much smaller than the expected statistical uncertainty per bin. Thus, the presented differential cross section measurements are model independent within the statistical uncertainties given that the tested scenarios of 100% VBF, VH, and $t\bar{t}H$ contributions in the signal model are extreme considering the current experimental knowledge of Higgs boson production.

10 Summary

The fiducial inclusive cross section for Higgs boson production in proton-proton collisions has been measured at a centre-of-mass energy of 13.6 TeV using the $H \rightarrow \gamma\gamma$ decay channel. The data were collected with the CMS detector at the LHC and correspond to an integrated luminosity of 34.7 fb^{-1} . A new normalizing-flow-based method is applied to correct the imperfect modelling of reconstructed photon variables in the simulation and to reduce the associated systematic uncertainties. The fiducial phase space is defined at the particle level and requires two isolated photons within the pseudorapidity $|\eta| < 2.5$ and not within $1.4442 < |\eta| < 1.5660$. These photons must fulfil a requirement on the geometric mean of their transverse momenta scaled by their invariant mass, $\sqrt{p_T^{\gamma_1} p_T^{\gamma_2}} / m_{\gamma\gamma} > 1/3$, which improves the perturbative convergence of the theoretical predictions, as well as the requirement $p_T^{\gamma_2} / m_{\gamma\gamma} > 1/4$. The measured




Figure 7: Differential fiducial cross sections for p_T^H (left) and the corresponding correlation matrix (right). The measured cross section in each bin is divided by the corresponding bin width. The coloured lines denote the predictions from different event generation setups, explained in the legend and in the text. The dashed boxes show the uncertainties in theoretical predictions on both the ggH and xH components. The p -value is calculated for the nominal SM prediction, which is MADGRAPH5.aMC@NLO with NNLOPS (MG5.aMC@NLO + NNLOPS) reweighting. The lower panel in the left plot shows the ratio to the nominal SM prediction. The last bin extends to infinity and the normalization of the bin is indicated in the plot.




Figure 8: Differential fiducial cross sections for $|y^H|$ (left) and the corresponding correlation matrix (right). Other details as for the caption of Fig. 7. In this case, the last bin does not extend to infinity, but it is limited to 2.5.




Figure 9: Differential fiducial cross sections for N_{Jets} (left) and the corresponding correlation matrix (right). Other details as for the caption of Fig. 7.




Figure 10: Differential fiducial cross sections for p_T^1 (left) and the corresponding correlation matrix (right). Other details as for the caption of Fig. 7.

inclusive fiducial cross section is $\sigma_{\text{fid}} = 74 \pm 11 \text{ (stat)}^{+5}_{-4} \text{ (syst) fb}$ and is in agreement with the standard-model (SM) expectation of $67.8 \pm 3.8 \text{ fb}$. Differential cross sections are measured as functions of the Higgs boson transverse momentum, rapidity, the number of associated jets, and the transverse momentum of the leading jet in the event. Within the uncertainties, the differential cross sections agree with the SM predictions.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centres and personnel of the Worldwide LHC Computing Grid and other centres for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2021-4.1.2-NEMZ_KI-2024-00036 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Centre for High Performance Computing, Big Data and Quantum Computing and FAIR – Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant

CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MICIU/AEI/10.13039/501100011033, ERDF/EU, "European Union NextGenerationEU/PRTR", and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G670016 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

References

- [1] ATLAS Collaboration, "Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC", *Phys. Lett. B* **716** (2012) 1, doi:10.1016/j.physletb.2012.08.020, arXiv:1207.7214.
- [2] CMS Collaboration, "Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC", *Phys. Lett. B* **716** (2012) 30, doi:10.1016/j.physletb.2012.08.021, arXiv:1207.7235.
- [3] CMS Collaboration, "Observation of a new boson with mass near 125 GeV in pp collisions at $\sqrt{s} = 7$ and 8 TeV", *JHEP* **06** (2013) 081, doi:10.1007/JHEP06(2013)081, arXiv:1303.4571.
- [4] L. Evans and P. Bryant (editors), "LHC machine", *JINST* **3** (2008) S08001, doi:10.1088/1748-0221/3/08/S08001.
- [5] ATLAS Collaboration, "A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery", *Nature* **607** (2022) 52, doi:10.1038/s41586-022-04893-w, arXiv:2207.00092. [Erratum: doi:10.1038/s41586-022-05581-5].
- [6] CMS Collaboration, "A portrait of the Higgs boson by the CMS experiment ten years after the discovery.", *Nature* **607** (2022) 60, doi:10.1038/s41586-022-04892-x, arXiv:2207.00043. [Erratum:doi:10.1038/s41586-023-06164-8].
- [7] LHC Higgs Cross Section Working Group, "Handbook of LHC Higgs cross sections: 4. deciphering the nature of the Higgs sector", 2016. arXiv:1610.07922.
- [8] ATLAS Collaboration, "Measurements of fiducial and differential cross sections for Higgs boson production in the diphoton decay channel at $\sqrt{s} = 8$ TeV with ATLAS", *JHEP* **09** (2014) 112, doi:10.1007/JHEP09(2014)112, arXiv:1407.4222.
- [9] CMS Collaboration, "Measurement of differential cross sections for Higgs boson production in the diphoton decay channel in pp collisions at $\sqrt{s} = 8$ TeV", *Eur. Phys. J. C* **76** (2016) 13, doi:10.1140/epjc/s10052-015-3853-3, arXiv:1508.07819.
- [10] ATLAS Collaboration, "Measurements of the Higgs boson inclusive and differential fiducial cross-sections in the diphoton decay channel with pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector", *JHEP* **08** (2022) 027, doi:10.1007/JHEP08(2022)027, arXiv:2202.00487.

- [11] CMS Collaboration, “Measurement of the Higgs boson inclusive and differential fiducial production cross sections in the diphoton decay channel with pp collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **07** (2023) 091, doi:10.1007/JHEP07(2023)091, arXiv:2208.12279.
- [12] ATLAS Collaboration, “Fiducial and differential cross sections of Higgs boson production measured in the four-lepton decay channel in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector”, *Phys. Lett. B* **738** (2014) 234, doi:10.1016/j.physletb.2014.09.054, arXiv:1408.3226.
- [13] CMS Collaboration, “Measurement of differential and integrated fiducial cross sections for Higgs boson production in the four-lepton decay channel in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, *JHEP* **04** (2016) 005, doi:10.1007/JHEP04(2016)005, arXiv:1512.08377.
- [14] ATLAS Collaboration, “Measurements of the Higgs boson inclusive and differential fiducial cross sections in the 4ℓ decay channel at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J. C* **80** (2020) 942, doi:10.1140/epjc/s10052-020-8223-0, arXiv:2004.03969.
- [15] CMS Collaboration, “Measurements of inclusive and differential cross sections for the Higgs boson production and decay to four-leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **08** (2023) 040, doi:10.1007/JHEP08(2023)040, arXiv:2305.07532.
- [16] ATLAS Collaboration, “Measurement of fiducial differential cross sections of gluon-fusion production of Higgs bosons decaying to $WW^* \rightarrow e\nu\mu\nu$ with the ATLAS detector at $\sqrt{s} = 8$ TeV”, *JHEP* **08** (2016) 104, doi:10.1007/JHEP08(2016)104, arXiv:1604.02997.
- [17] CMS Collaboration, “Measurement of the transverse momentum spectrum of the Higgs boson produced in pp collisions at $\sqrt{s} = 8$ TeV using $H \rightarrow WW$ decays”, *JHEP* **03** (2017) 032, doi:10.1007/JHEP03(2017)032, arXiv:1606.01522.
- [18] ATLAS Collaboration, “Measurements of differential cross sections of Higgs boson production through gluon fusion in the $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ final state at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **83** (2023) 774, doi:10.1140/epjc/s10052-023-11873-5, arXiv:2301.06822.
- [19] ATLAS Collaboration, “Fiducial and differential cross-section measurements for the vector-boson-fusion production of the Higgs boson in the $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ decay channel at 13 TeV with the ATLAS detector”, *Phys. Rev. D* **108** (2023) 072003, doi:10.1103/PhysRevD.108.072003, arXiv:2304.03053.
- [20] CMS Collaboration, “Measurement of the inclusive and differential Higgs boson production cross sections in the leptonic WW decay mode at $\sqrt{s} = 13$ TeV”, *JHEP* **03** (2021) 003, doi:10.1007/JHEP03(2021)003, arXiv:2007.01984.
- [21] CMS Collaboration, “Measurement of the inclusive and differential Higgs boson production cross sections in the decay mode to a pair of τ leptons in pp collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. Lett.* **128** (2022) 081805, doi:10.1103/PhysRevLett.128.081805, arXiv:2107.11486.
- [22] ATLAS Collaboration, “Differential cross-section measurements of Higgs boson production in the $H \rightarrow \tau^+\tau^+$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **03** (2025) 010, doi:10.1007/JHEP03(2025)010, arXiv:2407.16320.

-
- [23] ATLAS Collaboration, “Measurements of Higgs boson production by gluon-gluon fusion and vector-boson fusion using $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ decays in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. D* **108** (2023) 032005, doi:10.1103/PhysRevD.108.032005, arXiv:2207.00338.
- [24] ATLAS Collaboration, “Measurement of the total and differential Higgs boson production cross-sections at $\sqrt{s} = 13$ TeV with the ATLAS detector by combining the $H \rightarrow ZZ^* \rightarrow 4\ell$ and $H \rightarrow \gamma\gamma$ decay channels”, *JHEP* **05** (2023) 028, doi:10.1007/JHEP05(2023)028, arXiv:2207.08615.
- [25] ATLAS Collaboration, “Measurement of the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ cross-sections in pp collisions at $\sqrt{s} = 13.6$ TeV with the ATLAS detector”, *Eur. Phys. J. C* **84** (2024) 78, doi:10.1140/epjc/s10052-023-12130-5, arXiv:2306.11379.
- [26] CMS Collaboration, “Measurements of Higgs boson production cross section in the four-lepton final state in proton-proton collisions at $\sqrt{s} = 13.6$ TeV”, 2025. arXiv:2501.14849. Accepted by JHEP.
- [27] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [28] CMS Collaboration, “Development of the CMS detector for the CERN LHC Run 3”, *JINST* **19** (2024) P05064, doi:10.1088/1748-0221/19/05/P05064, arXiv:2309.05466.
- [29] G. P. Salam and E. Slade, “Cuts for two-body decays at colliders”, *JHEP* **11** (2021) 220, doi:10.1007/JHEP11(2021)220, arXiv:2106.08329.
- [30] CMS Collaboration, “Measurement of inclusive and differential Higgs boson production cross sections in the diphoton decay channel in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **01** (2019) 183, doi:10.1007/JHEP01(2019)183, arXiv:1807.03825.
- [31] HEPDATA record for this analysis, 2025. doi:10.17182/hepdata.157577.
- [32] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [33] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [34] CMS Collaboration, “Performance of the CMS high-level trigger during LHC Run 2”, *JINST* **19** (2024) P11021, doi:10.1088/1748-0221/19/11/P11021, arXiv:2410.17038.
- [35] CMS Collaboration, “Luminosity measurement in proton-proton collisions at 13.6 TeV in 2022 at CMS”, CMS Physics Analysis Summary CMS-PAS-LUM-22-001, 2024.
- [36] CMS Collaboration, “Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC”, *JINST* **16** (2021) P05014, doi:10.1088/1748-0221/16/05/P05014, arXiv:2012.06888.

- [37] J. Alwall et al., “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, *JHEP* **07** (2014) 079, doi:10.1007/JHEP07(2014)079, arXiv:1405.0301.
- [38] NNPDF Collaboration, “Parton distributions from high-precision collider data”, *Eur. Phys. J. C* **77** (2017) 663, doi:10.1140/epjc/s10052-017-5199-5, arXiv:1706.00428.
- [39] R. Frederix and S. Frixione, “Merging meets matching in MC@NLO”, *JHEP* **12** (2012) 061, doi:10.1007/JHEP12(2012)061, arXiv:1209.6215.
- [40] K. Hamilton, P. Nason, E. Re, and G. Zanderighi, “NNLOPS simulation of Higgs boson production”, *JHEP* **10** (2013) 222, doi:10.1007/JHEP10(2013)222, arXiv:1309.0017.
- [41] K. Hamilton, P. Nason, and G. Zanderighi, “MINLO: Multi-scale improved NLO”, *JHEP* **10** (2012) 155, doi:10.1007/JHEP10(2012)155, arXiv:1206.3572.
- [42] A. Kardos, P. Nason, and C. Oleari, “Three-jet production in POWHEG”, *JHEP* **04** (2014) 043, doi:10.1007/JHEP04(2014)043, arXiv:1402.4001.
- [43] A. Karlberg et al., “Ad interim recommendations for the Higgs boson production cross sections at $\sqrt{s} = 13.6$ TeV”, 2024. arXiv:2402.09955.
- [44] C. Anastasiou et al., “Higgs boson gluon-fusion production in QCD at three loops”, *Phys. Rev. Lett.* **114** (2015) 212001, doi:10.1103/PhysRevLett.114.212001, arXiv:1503.06056.
- [45] C. Anastasiou et al., “High precision determination of the gluon fusion Higgs boson cross-section at the LHC”, *JHEP* **05** (2016) 58, doi:10.1007/JHEP05(2016)058, arXiv:1602.00695.
- [46] M. Ciccolini, A. Denner, and S. Dittmaier, “Strong and electroweak corrections to the production of a Higgs boson+2 jets via weak interactions at the Large Hadron Collider”, *Phys. Rev. Lett.* **99** (2007) 161803, doi:10.1103/PhysRevLett.99.161803, arXiv:0707.0381.
- [47] M. Ciccolini, A. Denner, and S. Dittmaier, “Electroweak and QCD corrections to Higgs production via vector-boson fusion at the LHC”, *Phys. Rev. D* **77** (2008) 013002, doi:10.1103/PhysRevD.77.013002, arXiv:0710.4749.
- [48] P. Bolzoni, F. Maltoni, S.-O. Moch, and M. Zaro, “Higgs production via vector-boson fusion at NNLO in QCD”, *Phys. Rev. Lett.* **105** (2010) 011801, doi:10.1103/PhysRevLett.105.011801, arXiv:1003.4451.
- [49] P. Bolzoni, F. Maltoni, S.-O. Moch, and M. Zaro, “Vector boson fusion at next-to-next-to-leading order in QCD: Standard model Higgs boson and beyond”, *Phys. Rev. D* **85** (2012) 035002, doi:10.1103/PhysRevD.85.035002, arXiv:1109.3717.
- [50] O. Brein, A. Djouadi, and R. Harlander, “NNLO QCD corrections to the Higgs-strahlung processes at hadron colliders”, *Phys. Lett. B* **579** (2004) 149, doi:10.1016/j.physletb.2003.10.112, arXiv:hep-ph/0307206.

-
- [51] M. L. Ciccolini, S. Dittmaier, and M. Krämer, “Electroweak radiative corrections to associated WH and ZH production at hadron colliders”, *Phys. Rev. D* **68** (2003) 073003, doi:10.1103/PhysRevD.68.073003, arXiv:hep-ph/0306234.
 - [52] W. Beenakker et al., “Higgs radiation off top quarks at the Tevatron and the LHC”, *Phys. Rev. Lett.* **87** (2001) 201805, doi:10.1103/PhysRevLett.87.201805, arXiv:hep-ph/0107081.
 - [53] W. Beenakker et al., “NLO QCD corrections to $t\bar{t}H$ production in hadron collisions.”, *Nucl. Phys. B* **653** (2003) 151, doi:10.1016/S0550-3213(03)00044-0, arXiv:hep-ph/0211352.
 - [54] S. Dawson, L. H. Orr, L. Reina, and D. Wackeroth, “Associated top quark Higgs boson production at the LHC”, *Phys. Rev. D* **67** (2003) 071503, doi:10.1103/PhysRevD.67.071503, arXiv:hep-ph/0211438.
 - [55] S. Dawson et al., “Associated Higgs production with top quarks at the Large Hadron Collider: NLO QCD corrections”, *Phys. Rev. D* **68** (2003) 034022, doi:10.1103/PhysRevD.68.034022, arXiv:hep-ph/0305087.
 - [56] Z. Yu et al., “QCD NLO and EW NLO corrections to $t\bar{t}H$ production with top quark decays at hadron collider”, *Phys. Lett. B* **738** (2014) 1, doi:10.1016/j.physletb.2014.09.022, arXiv:1407.1110.
 - [57] S. S. Frixione et al., “Weak corrections to Higgs hadroproduction in association with a top-quark pair”, *JHEP* **09** (2014) 65, doi:10.1007/JHEP09(2014)065, arXiv:1407.0823.
 - [58] F. Demartin, F. Maltoni, K. Mawatari, and M. Zaro, “Higgs production in association with a single top quark at the LHC”, *Eur. Phys. J. C* **75** (2015) 267, doi:10.1140/epjc/s10052-015-3475-9, arXiv:1504.0611.
 - [59] F. Demartin et al., “ tWH associated production at the LHC”, *Eur. Phys. J. C* **77** (2017) 34, doi:10.1140/epjc/s10052-017-4601-7, arXiv:1607.05862.
 - [60] A. Denner et al., “Standard model Higgs-boson branching ratios with uncertainties”, *Eur. Phys. J. C* **71** (2011) 1753, doi:10.1140/epjc/s10052-011-1753-8, arXiv:1107.5909.
 - [61] A. Djouadi, J. Kalinowski, M. Muehlleitner, and M. Spira, “HDECAY: Twenty₊₊ years after”, *Comput. Phys. Commun.* **238** (2019) 214, doi:10.1016/j.cpc.2018.12.010, arXiv:1801.09506.
 - [62] A. Bredenstein, A. Denner, S. Dittmaier, and M. M. Weber, “Precise predictions for the Higgs-boson decay $H \rightarrow WW/ZZ \rightarrow 4$ leptons”, *Phys. Rev. D* **74** (2006) 013004, doi:10.1103/PhysRevD.74.013004, arXiv:hep-ph/0604011.
 - [63] A. Bredenstein, A. Denner, S. Dittmaier, and M. M. Weber, “Radiative corrections to the semileptonic and hadronic Higgs-boson decays $H \rightarrow WW/ZZ \rightarrow 4$ fermions”, *JHEP* **02** (2007) 80, doi:10.1088/1126-6708/2007/02/080, arXiv:hep-ph/0611234.
 - [64] S. Boselli et al., “Higgs boson decay into four leptons at NLOPS electroweak accuracy”, *JHEP* **06** (2015) 23, doi:10.1007/JHEP06(2015)023, arXiv:1503.07394.

- [65] S. Actis, G. Passarino, C. Sturm, and S. Uccirati, “NNLO computational techniques: the cases $H \rightarrow \gamma\gamma$ and $H \rightarrow gg$ ”, *Nucl. Phys. B* **811** (2009) 182, doi:10.1016/j.nuclphysb.2008.11.024, arXiv:0809.3667.
- [66] Sherpa Collaboration, “Event generation with Sherpa 2.2”, *SciPost Phys.* **7** (2019) 034, doi:10.21468/SciPostPhys.7.3.034, arXiv:1905.09127.
- [67] C. Bierlich et al., “A comprehensive guide to the physics and usage of PYTHIA 8.3”, *SciPost Phys. Codeb.* **2022** (2022) 8, doi:10.21468/SciPostPhysCodeb.8, arXiv:2203.11601.
- [68] CMS Collaboration, “Performance of the CMS muon trigger system in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **16** (2021) P07001, doi:10.1088/1748-0221/16/07/P07001, arXiv:2102.04790.
- [69] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) 4, doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.
- [70] GEANT4 Collaboration, “GEANT4 – a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [71] CMS Collaboration, “Technical proposal for the Phase-II upgrade of the Compact Muon Solenoid”, CMS Technical Proposal CERN-LHCC-2015-010, CMS-TDR-15-02, 2015.
- [72] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965.
- [73] CMS Collaboration, “Electron and photon reconstruction and identification performance at CMS in 2022 and 2023”, CMS Detector Performance Note CMS-DP-2024-052, 2024.
- [74] CMS Collaboration, “A measurement of the Higgs boson mass in the diphoton decay channel”, *Phys. Lett. B* **805** (2020) 135425, doi:10.1016/j.physletb.2020.135425, arXiv:2002.06398.
- [75] CMS Collaboration, “The CMS electromagnetic calorimeter project: Technical design report”, CMS Technical Design Report CERN-LHCC-97-033, CMS-TDR-4, 1997.
- [76] T. Chen and C. Guestrin, “XGBoost: A scalable tree boosting system”, in *Proc. 22nd ACM SIGKDD Int. Conf. Know. Discov. Data Min.*, p. 785. 2016. doi:10.1145/2939672.2939785.
- [77] CMS Collaboration, “Performance of photon reconstruction and identification with the CMS detector in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JINST* **10** (2015) P08010, doi:10.1088/1748-0221/10/08/P08010, arXiv:1502.02702.
- [78] CMS Collaboration, “Measurements of inclusive W and Z cross sections in pp collisions at $\sqrt{s} = 7$ TeV”, *JHEP* **01** (2011) 080, doi:10.1007/JHEP01(2011)080, arXiv:1012.2466.
- [79] G. Papamakarios et al., “Normalizing flows for probabilistic modeling and inference”, *J. Mach. Learn. Res.* **22** (2021) 1, doi:10.48550/arXiv.1912.02762, arXiv:1912.02762.

-
- [80] C. Daumann et al., “One flow to correct them all: Improving simulations in high-energy physics with a single normalising flow and a switch”, *Comput. Softw. Big. Sci.* **8** (2024) 15, doi:10.1007/s41781-024-00125-0, arXiv:2403.18582.
- [81] G. Papamakarios, T. Pavlakou, and I. Murray, “Masked autoregressive flow for density estimation”, 2018. arXiv:1705.07057.
- [82] C. Durkan, A. Bekasov, I. Murray, and G. Papamakarios, “Neural spline flows”, 2019. arXiv:1906.04032.
- [83] A. Paszke et al., “PYTORCH: An imperative style, high-performance deep learning library”, in *Proc. 33rd Conference on Neural Information Processing Systems (NeurIPS 2019): Vancouver, Canada, December 08–14, 2019*. 2019. arXiv:1912.01703.
- [84] F. Rozet, F. Divo, and S. Schnake, “Zuko: Normalizing flows in PyTorch”, 2022. doi:10.5281/zenodo.10070008.
- [85] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [86] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- k_T jet clustering algorithm”, *JHEP* **04** (2008) 063, doi:10.1088/1126-6708/2008/04/063, arXiv:0802.1189.
- [87] M. Cacciari, G. P. Salam, and G. Soyez, “Fastjet user manual”, *Eur. Phys. J. C* **72** (2012) 1896, doi:10.1140/epjc/s10052-012-1896-2, arXiv:1111.6097.
- [88] CMS Collaboration, “Pileup mitigation at CMS in 13 TeV data”, *JINST* **15** (2020) P09018, doi:10.1088/1748-0221/15/09/P09018, arXiv:2003.00503.
- [89] CMS Collaboration, “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV”, *JINST* **12** (2017) P02014, doi:10.1088/1748-0221/12/02/P02014, arXiv:1607.03663.
- [90] CMS Collaboration, “Measurements of Higgs boson properties in the diphoton decay channel in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **11** (2018) 185, doi:10.1007/JHEP11(2018)185, arXiv:1804.02716.
- [91] CMS Collaboration, “Measurements of Higgs boson production cross sections and couplings in the diphoton decay channel at $\sqrt{s} = 13$ TeV”, *JHEP* **07** (2021) 027, doi:10.1007/JHEP07(2021)027, arXiv:2103.06956.
- [92] CMS Collaboration, “The CMS statistical analysis and combination tool: COMBINE”, *Comput. Softw. Big Sci.* **8** (2024) 19, doi:10.1007/s41781-024-00121-4, arXiv:2404.06614.
- [93] J. Campbell, M. Carena, R. Harnik, and Z. Liu, “Interference in the $gg \rightarrow h \rightarrow \gamma\gamma$ on-shell rate and the Higgs boson total width”, *Phys. Rev. Lett.* **119** (2017) 181801, doi:10.1103/PhysRevLett.119.181801, arXiv:1704.08259. [Addendum: doi:10.1103/PhysRevLett.119.199901].
- [94] P. D. Dauncey, M. Kenzie, N. Wardle, and G. J. Davies, “Handling uncertainties in background shapes: the discrete profiling method”, *JINST* **10** (2015) P04015, doi:10.1088/1748-0221/10/04/P04015, arXiv:1408.6865.

- [95] R. A. Fisher, “On the interpretation of χ^2 from contingency tables, and the calculation of p”, *J. R. Stat. Soc.* **85** (1922) 87, doi:10.1111/j.2397-2335.1922.tb00768.x.
- [96] R. J. Barlow and C. Beeston, “Fitting using finite Monte Carlo samples”, *Comput. Phys. Commun.* **77** (1993) 219, doi:10.1016/0010-4655(93)90005-W.
- [97] J. S. Conway, “Incorporating nuisance parameters in likelihoods for multisource spectra”, in *Proc. PHYSTAT 2011*, p. 115. 2011. arXiv:1103.0354. doi:10.5170/CERN-2011-006.115.
- [98] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [99] P. Nason, “A new method for combining NLO QCD with shower Monte Carlo algorithms”, *JHEP* **11** (2004) 040, doi:10.1088/1126-6708/2004/11/040, arXiv:hep-ph/0409146.
- [100] S. Frixione, P. Nason, and C. Oleari, “Matching NLO QCD computations with parton shower simulations: the POWHEG method”, *JHEP* **11** (2007) 070, doi:10.1088/1126-6708/2007/11/070, arXiv:0709.2092.
- [101] S. Alioli, P. Nason, C. Oleari, and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX”, *JHEP* **06** (2010) 043, doi:10.1007/JHEP06(2010)043, arXiv:1002.2581.
- [102] E. Bagnaschi, G. Degrandi, P. Slavich, and A. Vicini, “Higgs production via gluon fusion in the POWHEG approach in the SM and in the MSSM”, *JHEP* **02** (2012) 088, doi:10.1007/JHEP02(2012)088, arXiv:1111.2854.
- [103] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, “Asymptotic formulae for likelihood-based tests of new physics”, *Eur. Phys. J. C* **71** (2011) 1554, doi:10.1140/epjc/s10052-011-1554-0, arXiv:1007.1727. [Erratum: doi:10.1140/epjc/s10052-013-2501-z].

A The CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, V. Makarenko , A. Tumasyan¹ 

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, L. Benato , T. Bergauer , K. Damanakis , M. Dragicevic , C. Giordano, P.S. Hussain , M. Jeitler² , N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck² , R. Schöfbeck² , D. Schwarz , M. Shooshtari, M. Sonawane , W. Waltenberger , C.-E. Wulz² 

Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , H. Kwon , T. Van Laer, P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

J. Bierkens , N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, Y. Hong , S. Lowette , I. Makarenko , D. Müller , J. Song , S. Tavernier , M. Tytgat³ , G.P. Van Onsem , S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios , A. Khalilzadeh, F.A. Khan , A. Malara , M.A. Shahzad, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer , F. Zhang 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , J. Knolle , L. Lambrecht , D. Marckx , K. Skovpen , N. Van Den Bossche , J. van der Linden , J. Vandenbroeck , L. Wezenbeek 

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

S. Bein , A. Benecke , A. Bethani , G. Bruno , A. Cappati , J. De Favereau De Jeneret , C. Delaere , A. Giammanco , A.O. Guzel , V. Lemaitre, J. Lidrych , P. Mastrapasqua , S. Turkcapar 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , C. Hensel , T. Menezes De Oliveira , C. Mora Herrera⁴ , P. Rebello Teles , M. Soeiro, E.J. Tonelli Manganote⁵ , A. Vilela Pereira⁴ 

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁶, M. Costa Reis , E.M. Da Costa , G.G. Da Silveira⁷ , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza, S. S. Jesus , T. Laux Kuhn⁷ , M. Macedo , K. Mota Amarilo , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel , F. Torres Da Silva De Araujo⁸ 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁷ , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , B. Lopes Da Costa, I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula , V. Scheurer

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , P. Danev, R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 

University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , D. Laroze , S. Thakur 

Universidad Técnica Federico Santa María, Valparaiso, Chile

W. Brooks 

Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu, X. Wang 

Institute of High Energy Physics, Beijing, China

G.M. Chen⁹ , H.S. Chen⁹ , M. Chen⁹ , Y. Chen , Q. Hou , X. Hou⁹, F. Iemmi¹⁰ , A. Kapoor¹⁰ , H. Liao , Z.-A. Liu¹¹ , S. Song⁹, J. Tao , C. Wang⁹, J. Wang , A. Zada⁹ , H. Zhang , Z. Zhang⁹, J. Zhao 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, Q. Guo, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao, C. Pan, S. Qian, S.J. Qian , X. Qin, X. Sun , D. Wang , J. Wang, H. Yang, Y. Zhao, C. Zhou 

Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 

Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer¹², B. Li¹³, H. Wang , K. Yi¹⁴ , J. Zhang 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

Y. Li

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao¹⁵ 

Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo , A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

C. Rendón , M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

N. Godinovic , D. Lelas , A. Sculac 

University of Split, Faculty of Science, Split, Croatia

M. Kovac , A. Petkovic , T. Sculac 

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov , T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , C. Nicolaou, L. Paizanos, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

Charles University, Prague, Czech Republic

M. Finger , M. Finger Jr. , A. Kveton 

Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

A.A. Abdelalim^{16,17} , S. Elgammal¹⁸, A. Ellithi Kamel¹⁹

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

M. Abdullah Al-Mashad , A. Hussein, H. Mohammed 

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht , M. Kadastik, T. Lange , C. Nielsen , J. Pata , M. Raidal , N. Seeba , L. Tani , C. Veelken 

Department of Physics, University of Helsinki, Helsinki, Finland

A. Milieva, K. Osterberg , M. Voutilainen 

Helsinki Institute of Physics, Helsinki, Finland

N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew , K.T.S. Kallonen , R. Kumar Verma , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , N.R. Mancilla Xinto, M. Myllymäki , M.m. Rantanen , S. Saariokari , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

H. Kirschenmann , P. Luukka , H. Petrow 

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, P. Devouge, J.L. Faure, F. Ferri , P. Gagne, S. Ganjour , P. Gras , F. Guilloux , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi , L. Portales , S. Ronchi, M.Ö. Sahin , A. Savoy-Navarro²⁰ , P. Simkina , M. Titov , M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

F. Beaudette , G. Boldrini , P. Busson , C. Charlot , M. Chiusi , T.D. Cuisset , F. Damas , O. Davignon , A. De Wit , T. Debnath , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , L. Kalipoliti , G. Liu , M. Manoni , M. Nguyen , S. Obraztsov , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , G. Sokmen, L. Urda Gómez , A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram²¹ , J. Andrea , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet 

G. Saha , M.A. Sessini , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo , P. Depasse , C. Dozen²² , H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg , G. Grenier , B. Ille , E. Jourd'huy, I.B. Laktineh, M. Lethuillier , B. Massoteau, L. Mirabito, S. Perries, A. Purohit , M. Vander Donckt , J. Xiao 

Georgian Technical University, Tbilisi, Georgia

G. Adamov, I. Lomidze , Z. Tsamalaidze²³ 

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser , P. Natland, V. Oppenländer, A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

C. Daumann, S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann , M. Erdmann , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , N. Kumar , M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , F. Nowotny, A. Pozdnyakov , W. Redjeb , H. Reithler , U. Sarkar , V. Sarkisovi , A. Schmidt , C. Seth, A. Sharma , J.L. Spah , S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , N. Hoeflich , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman²⁴ , K. Borras²⁵ , A. Campbell , S. Chatterjee , L.X. Coll Saravia , G. Eckerlin, D. Eckstein , E. Gallo²⁴ , A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka²⁶ , W. Lohmann²⁷ , F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , J. Park , E. Ranken , A. Raspereza , D. Rastorguev , L. Rygaard, M. Scham^{28,25} , S. Schnake²⁵ , P. Schütze , C. Schwanenberger²⁴ , D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford , F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , K. Wichmann, L. Wiens²⁵ , C. Wissing , Y. Yang , S. Zakharov, A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , M. Antonello , S. Bollweg, M. Bonanomi , K. El Morabit , Y. Fischer , M. Frahm, E. Garutti , A. Grohsjean , J. Haller , D. Hundhausen, H.R. Jabusch , G. Kasieczka , P. Keicher , R. Klanner , W. Korcar , T. Kramer , C.c. Kuo, V. Kutzner , F. Labe , J. Lange , A. Lobanov , L. Moureaux , M. Mrowietz, A. Nigamova , K. Nikolopoulos, Y. Nissan, A. Paasch , K.J. Pena Rodriguez , N. Prouvost, T. Quadfasel , B. Raciti , M. Rieger , D. Savoiu , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, R. Ward,

B. Wiederspan, M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , E. Butz , Y.M. Chen , T. Chwalek , A. Dierlamm , G.G. Dincer , U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann²⁹ , R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , O. Lavoryk , J.M. Lawhorn , M. Link, A. Lintuluoto , S. Maier , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , R. Schmieder, N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa, L. Stockmeier, K. Tauqeer, M. Toms , B. Topko , N. Trevisani , C. Verstege , T. Voigtländer , R.F. Von Cube , J. Von Den Driesch, M. Wassmer , F. Wittig, R. Wolf , W.D. Zeuner, X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis , A. Kyriakis , A. Papadopoulos²⁹, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

G. Melachroinos, Z. Painesis , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 

National Technical University of Athens, Athens, Greece

T. Chatzistavrou, G. Karapostoli , K. Kousouris , E. Siamarkou, G. Tsipolitis 

University of Ioánnina, Ioánnina, Greece

I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

D. Druzhkin , C. Hajdu , D. Horvath^{30,31} , K. Márton, A.J. Rádl³² , F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkuti³³ , M.M.A. Gadallah³⁴ , Á. Kadlecik , G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 

HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo³³ , F. Nemes³³ , T. Novak , I. Szanyi³⁵ 

Panjab University, Chandigarh, India

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra³⁶ , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand, J.B. Singh , A. Singla 

University of Delhi, Delhi, India

A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya 

Indian Institute of Technology Kanpur, Kanpur, India

S. Mukherjee 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

Indian Institute of Technology Madras, Madras, India

M.M. Ameen , P.K. Behera , S. Chatterjee , G. Dash , A. Dattamunsi, P. Jana , P. Kalbhor , S. Kamble , J.R. Komaragiri³⁷ , T. Mishra , P.R. Pujahari , N.R. Saha , A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay 

IISER Mohali, India, Mohali, India

B.K. Sirasva

Tata Institute of Fundamental Research-A, Mumbai, India

L. Bhatt, S. Dugad, G.B. Mohanty , M. Shelake, P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , S. Barman³⁸ , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, B.M. Joshi , S. Kumar , M. Maity³⁸, G. Majumder , K. Mazumdar , S. Parolia , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati³⁹ , D. Maity⁴⁰ , P. Mal , K. Naskar⁴⁰ , A. Nayak⁴⁰ , S. Nayak, K. Pal , R. Raturi, P. Sadangi, S.K. Swain , S. Varghese⁴⁰ , D. Vats⁴⁰ 

Indian Institute of Science Education and Research (IISER), Pune, India

S. Acharya⁴¹ , A. Alpana , S. Dube , B. Gomber⁴¹ , P. Hazarika , B. Kansal , A. Laha , B. Sahu⁴¹ , R. Sharma , S. Sharma , K.Y. Vaish 

Indian Institute of Technology Hyderabad, Telangana, India

S. Ghosh 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi⁴² , A. Jafari⁴³ , M. Zeinali⁴⁴ 

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Bashiri, S. Chenarani⁴⁵ , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie , M. Mohammadi Najafabadi , S. Tizchang⁴⁶ 

University College Dublin, Dublin, Ireland

M. Felcini , M. Grunewald 

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia^{a,b} , M. Barbieri^{a,b}, M. Buonsante^{a,b} , A. Colaleo^{a,b} , D. Creanza^{a,c} , B. D'Anzi^{a,b} , N. De Filippis^{a,c} , M. De Palma^{a,b} , W. Elmetenawee^{a,b,16} , N. Ferrara^{a,b} , L. Fiore^a , L. Longo^a , M. Louka^{a,b}, G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^a , V. Mastrapasqua^{a,b} , S. My^{a,b} , S. Nuzzo^{a,b} , A. Pellecchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} , R. Radogna^{a,b} , D. Ramos^a , A. Ranieri^a , L. Silvestris^a , F.M. Simone^{a,c} , Ü. Sözbilir^a , A. Stamerra^{a,b} , D. Troiano^{a,b} , R. Venditti^{a,b} , P. Verwilligen^a , A. Zaza^{a,b} 

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , P. Capiluppi^{a,b} , A. Castro^{+a,b} , F.R. Cavallo^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} , D. Fasanella^a , P. Giacomelli^a , L. Guiducci^{a,b} , S. Lo Meo^{a,47} , M. Lorusso^{a,b} , L. Lunerti^a , S. Marcellini^a , G. Masetti^a , F.L. Navarria^{a,b} 

G. Paggi^{a,b} , A. Perrotta^a , F. Primavera^{a,b} , A.M. Rossi^{a,b} , S. Rossi Tisbeni^{a,b} ,
T. Rovelli^{a,b} , G.P. Siroli^{a,b} 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,48} , A. Di Mattia^a , A. Lapertosa^a , R. Potenza^{a,b}, A. Tricomi^{a,b,48} 

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

J. Altork^{a,b}, P. Assiouras^a , G. Barbagli^a , G. Bardelli^a , M. Bartolini^{a,b}, A. Calandri^{a,b} ,
B. Camaiani^{a,b} , A. Cassese^a , R. Ceccarelli^a , V. Ciulli^{a,b} , C. Civinini^a ,
R. D'Alessandro^{a,b} , L. Damenti^{a,b}, E. Focardi^{a,b} , T. Kello^a , G. Latino^{a,b} , P. Lenzi^{a,b} ,
M. Lizzo^a , M. Meschini^a , S. Paoletti^a , A. Papanastassiou^{a,b}, G. Sguazzoni^a ,
L. Viliani^a 

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁴⁹ , D. Piccolo 

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

M. Alves Gallo Pereira^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} 

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a , F. Brivio^a , F. Cetorelli^{a,b} , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a ,
S. Gennai^a , R. Gerosa^{a,b} , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^a , G. Lavizzari^{a,b},
M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a ,
L. Moroni^a , M. Paganoni^{a,b} , S. Palluotto^{a,b} , D. Pedrini^a , A. Perego^{a,b} ,
B.S. Pinolini^a, G. Pizzati^{a,b} , S. Ragazzi^{a,b} , T. Tabarelli de Fatis^{a,b} 

INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo^a , A. Cagnotta^{a,b} , F. Carnevali^{a,b}, C. Di Fraia^{a,b} , F. Fabozzi^{a,c} ,
L. Favilla^{a,d}, A.O.M. Iorio^{a,b} , L. Lista^{a,b,50} , P. Paolucci^{a,29} , B. Rossi^a 

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università degli Studi di Cagliari^c, Cagliari, Italy

R. Ardino^a , P. Azzi^a , N. Bacchetta^{a,51} , D. Bisello^{a,b} , P. Bortignon^a , G. Bortolato^{a,b},
A.C.M. Bulla^a , R. Carlin^{a,b} , P. Checchia^a , T. Dorigo^{a,52} , F. Gasparini^{a,b} ,
U. Gasparini^{a,b} , S. Giorgetti^a, M. Gulmini^{a,53} , E. Lusiani^a , M. Margoni^{a,b} ,
J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , M. Tosi^{a,b} ,
A. Triossi^{a,b} , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} , G. Zumerle^{a,b} 

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

A. Braghieri^a , S. Calzaferri^a , D. Fiorina^a , P. Montagna^{a,b} , M. Pelliccioni^a , V. Re^a ,
C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b} 

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal^{a,b} , M.E. Ascioti^{a,b}, G.M. Bilei^a , C. Carrivale^{a,b}, D. Ciangottini^{a,b} ,
L. Della Penna^a, L. Fanò^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,54} ,
A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} 

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

C. Aimè^{a,b} , C.A. Alexe^{a,c} , P. Asenov^{a,b} , P. Azzurri^a , G. Bagliesi^a ,
R. Bhattacharya^a , L. Bianchini^{a,b} , T. Boccali^a , E. Bossini^a , D. Bruschini^{a,c} ,
R. Castaldi^a , F. Cattafesta^{a,c} , M.A. Ciocci^{a,b} , M. Cipriani^{a,b} , V. D'Amante^{a,d} ,
R. Dell'Orso^a , S. Donato^{a,b} , R. Forti^{a,b} , A. Giassi^a , F. Ligabue^{a,c} 

A.C. Marini^{a,b} , D. Matos Figueiredo^a , A. Messineo^{a,b} , S. Mishra^a , V.K. Muraleedharan Nair Bindhu^{a,b} , M. Musich^{a,b} , S. Nandan^a , F. Palla^a , M. Riggirello^{a,c} , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^{a,55} , T. Sarkar^a , A. Scribano^a , P. Spagnolo^a , F. Tenchini^{a,b} , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , F. Vaselli^{a,c} , A. Venturi^a , P.G. Verdini^a 

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Akrap^{a,b} , C. Basile^{a,b} , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , F. De Ruggi^{a,b} , D. Del Re^{a,b} , E. Di Marco^{a,b} , M. Diemoz^a , F. Errico^{a,b} , L. Frosina^{a,b} , R. Gargiulo^{a,b} , B. Harikrishnan^{a,b} , F. Lombardi^{a,b} , E. Longo^{a,b} , L. Martikainen^{a,b} , J. Mijuskovic^{a,b} , G. Organtini^{a,b} , N. Palmeri^{a,b} , R. Paramatti^{a,b} , C. Quaranta^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a , V. Vladimirov^{a,b} 

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^{a,c} , R. Bellan^{a,b} , C. Biino^a , C. Borca^{a,b} , N. Cartiglia^a , S. Coli^a , M. Costa^{a,b} , R. Covarelli^{a,b} , N. Demaria^a , L. Finco^a , M. Grippo^{a,b} , B. Kiani^{a,b} , L. Lanteri^{a,b} , F. Legger^a , F. Luongo^{a,b} , C. Mariotti^a , L. Markovic^{a,b} , S. Maselli^a , A. Mecca^{a,b} , L. Menzio^{a,b} , P. Meridiani^a , E. Migliore^{a,b} , M. Monteno^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , F. Rotondo^a , M. Ruspa^{a,c} , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} , C. Tarricone^{a,b} , D. Trocino^a , G. Umoret^{a,b} , R. White^{a,b} 

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar^{a,b} , S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , K. De Leo^a , G. Della Ricca^{a,b} , R. Delli Gatti^{a,b} 

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , J. Kim, T. Kim, D. Lee, H. Lee, J. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , S. Sekmen , B. Tae, Y.C. Yang 

Department of Mathematics and Physics - GWNU, Gangneung, Korea

M.S. Kim 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon 

Hanyang University, Seoul, Korea

E. Asilar , J. Choi⁵⁶ , D. Kim , T.J. Kim , J.A. Merlin, Y. Ryou

Korea University, Seoul, Korea

S. Han, B. Hong , K. Lee, K.S. Lee , S. Lee , J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh , J. Shin , S. Yang 

Sejong University, Seoul, Korea

Y. Kang , H. S. Kim , Y. Kim, S. Lee

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , Y. Kim, Y.W. Kim , S. Ko , H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , J. Shin, U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea

G. Cho, S. Ha , K. Hwang , B. Kim , S. Kim, K. Lee , H.D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , M.R. Kim , Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy , Y. Gharbia 

Kuwait University - College of Science - Department of Physics, Safat, Kuwait

F. Alazemi 

Riga Technical University, Riga, Latvia

K. Dreimanis , O.M. Eberlins , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs, R. Plese , A. Potrebko , M. Seidel , D. Sidiropoulos Kontos 

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Yusuff⁵⁷ , Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , A. Cota Rodriguez , L.E. Cuevas Picos, H.A. Encinas Acosta, L.G. Gallegos Maríñez, M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, R. Lopez-Fernandez , J. Mejia Guisao , R. Reyes-Almanza , A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , F.E. Neri Huerta , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

University of Montenegro, Podgorica, Montenegro

I. Bujanja , N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, W.A. Khan 

AGH University of Krakow, Krakow, Poland

V. Avati, A. Bellora⁵⁸ , L. Forthomme , L. Grzanka , M. Malawski , K. Piotrkowski

National Centre for Nuclear Research, Swierk, Poland

M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad 

Warsaw University of Technology, Warsaw, Poland

P. Fokow , K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , T. Camporesi , G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff 

Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , P. Milenovic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

D. Devetak , M. Dordevic , J. Milosevic , L. Nadderd , V. Rekovic , M. Stojanovic 

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. Llorente Merino , C. Martin Perez , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , J. Vazquez Escobar 

Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , A. Cardini , J. Cuevas , J. Del Riego Badas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , P. Leguina , M. Obeso Menendez , E. Palencia Cortezon , J. Prado Pico , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasosa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , C. Quintana San Emeterio , L. Scodellaro , I. Vila , R. Vilar Cortabitarte , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

B. Kailasapathy⁵⁹ , D.D.C. Wickramarathna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna⁶⁰ , K. Liyanage , N. Perera 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , J. Baechler , D. Barney , M. Bianco , A. Bocci , L. Borgonovi , C. Botta , A. Bragagnolo , C.E. Brown , C. Caillol , G. Cerminara , P. Connor , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck 

M.M. Defranchis , M. Deile , M. Dobson , W. Funk , A. Gaddi, S. Giani, D. Gigi, K. Gill , F. Glege , M. Glowacki, A. Gruber, J. Hegeman , J.K. Heikkilä , B. Huber , V. Innocente , T. James , P. Janot , J. Jaroslavceva, O. Kaluzinska , O. Karacheban²⁷ , G. Karathanasis , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , M. Magherini , L. Malgeri , M. Mannelli , M. Matthewman, A. Mehta , F. Meijers , S. Mersi , E. Meschi , M. Migliorini , V. Milosevic , F. Monti , F. Moortgat , M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , M. Pari, G. Petrucciani , A. Pfeiffer , M. Pierini , M. Pitt , H. Qu , D. Rabadý , B. Ribeiro Lopes , F. Riti , P. Rosado , M. Rovere , H. Sakulin , R. Salvatico , S. Sanchez Cruz , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas⁶¹ , A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea , E. Vernazza , J. Wanczyk⁶² , J. Wang, S. Wuchterl , M. Zarucki , P. Zehetner , P. Zejdl , G. Zevi Della Porta 

PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T. Bevilacqua⁶³ , L. Caminada⁶³ , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli⁶³ , L. Noehte⁶³ , T. Rohe , A. Samalan

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , M. Backhaus , G. Bonomelli , C. Cazzaniga , K. Datta , P. De Bryas Dexmiers D'archiac⁶² , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer , T.G. Harte, W. Lustermann , A.-M. Lyon , M. Malucchi , R.A. Manzoni , M. Marchegiani , L. Marchese , A. Mascellani⁶² , F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , B. Ristic , R. Seidita , A. Tarabini , D. Valsecchi , R. Wallny 

Universität Zürich, Zurich, Switzerland

C. Amsler⁶⁴ , P. Bäertschi , M.F. Canelli , G. Celotto, K. Cormier , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechti , A. Macchiolo , F. Meng , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr, F. Stäger , R. Tramontano 

National Central University, Chung-Li, Taiwan

D. Bhowmik, C.M. Kuo, P.K. Rout , S. Taj, P.C. Tiwari³⁷ 

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao, S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker , L.s. Tsai, D. Tsionou, H.y. Wu, E. Yazgan 

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas 

Tunis El Manar University, Tunis, Tunisia

Y. Maghrbi 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu⁶⁵ , Y. Guler⁶⁶ , E. Gurpinar Guler⁶⁶ , C. Isik , O. Kara, A. Kayis Topaksu , Y. Komurcu , G. Onengut , K. Ozdemir⁶⁷ , B. Tali⁶⁸ , U.G. Tok , E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

M. Yalvac⁶⁹ 

Bogazici University, Istanbul, Turkey

B. Akgun^{id}, I.O. Atakisi^{id}, E. Gülmez^{id}, M. Kaya^{70 id}, O. Kaya^{71 id}, M.A. Sarkisla⁷², S. Tekten^{73 id}

Istanbul Technical University, Istanbul, Turkey

A. Cakir^{id}, K. Cankocak^{65,74 id}, S. Sen^{75 id}

Istanbul University, Istanbul, Turkey

O. Aydılek^{76 id}, B. Hacısahinoglu^{id}, I. Hos^{77 id}, B. Kaynak^{id}, S. Ozkorucuklu^{id}, O. Potok^{id}, H. Sert^{id}, C. Simsek^{id}, C. Zorbilmez^{id}

Yildiz Technical University, Istanbul, Turkey

S. Cerci^{id}, B. Isildak^{78 id}, D. Sunar Cerci^{id}, T. Yetkin^{22 id}

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev^{id}, O. Dadazhanova, B. Grynyov^{id}

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk^{id}

University of Bristol, Bristol, United Kingdom

J.J. Brooke^{id}, A. Bundock^{id}, F. Bury^{id}, E. Clement^{id}, D. Cussans^{id}, H. Flacher^{id}, J. Goldstein^{id}, H.F. Heath^{id}, M.-L. Holmberg^{id}, L. Kreczko^{id}, S. Paramesvaran^{id}, L. Robertshaw, J. Segal, V.J. Smith^{id}

Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell^{id}, A. Belyaev^{79 id}, C. Brew^{id}, R.M. Brown^{id}, D.J.A. Cockerill^{id}, C. Cooke^{id}, A. Elliot^{id}, K.V. Ellis, J. Gajownik, K. Harder^{id}, S. Harper^{id}, J. Linacre^{id}, K. Manolopoulos, M. Moallemi^{id}, D.M. Newbold^{id}, E. Olaiya, D. Petyt^{id}, T. Reis^{id}, A.R. Sahasransu^{id}, G. Salvi^{id}, T. Schuh, C.H. Shepherd-Themistocleous^{id}, I.R. Tomalin^{id}, K.C. Whalen^{id}, T. Williams^{id}

Imperial College, London, United Kingdom

I. Andreou^{id}, R. Bainbridge^{id}, P. Bloch^{id}, O. Buchmuller, C.A. Carrillo Montoya^{id}, D. Colling^{id}, J.S. Dancu, I. Das^{id}, P. Dauncey^{id}, G. Davies^{id}, M. Della Negra^{id}, S. Fayer, G. Fedi^{id}, G. Hall^{id}, H.R. Hoorani^{id}, A. Howard, G. Iles^{id}, C.R. Knight^{id}, P. Krueper, J. Langford^{id}, K.H. Law^{id}, J. León Holgado^{id}, L. Lyons^{id}, A.-M. Magnan^{id}, B. Maier^{id}, S. Mallios, A. Mastronikolis, M. Mieskolainen^{id}, J. Nash^{80 id}, M. Pesaresi^{id}, P.B. Pradeep, B.C. Radburn-Smith^{id}, A. Richards, A. Rose^{id}, L. Russell^{id}, K. Savva^{id}, C. Seez^{id}, R. Shukla^{id}, A. Tapper^{id}, K. Uchida^{id}, G.P. Uttley^{id}, T. Virdee^{29 id}, M. Vojinovic^{id}, N. Wardle^{id}, D. Winterbottom^{id}

Brunel University, Uxbridge, United Kingdom

J.E. Cole^{id}, A. Khan, P. Kyberd^{id}, I.D. Reid^{id}

Baylor University, Waco, Texas, USA

S. Abdullin^{id}, A. Brinkerhoff^{id}, E. Collins^{id}, M.R. Darwish^{id}, J. Dittmann^{id}, K. Hatakeyama^{id}, V. Hegde^{id}, J. Hiltbrand^{id}, B. McMaster^{id}, J. Samudio^{id}, S. Sawant^{id}, C. Sutantawibul^{id}, J. Wilson^{id}

Catholic University of America, Washington, DC, USA

R. Bartek^{id}, A. Dominguez^{id}, S. Raj^{id}, A.E. Simsek^{id}, S.S. Yu^{id}

The University of Alabama, Tuscaloosa, Alabama, USA

B. Bam , S.C. Behera , A. Buchot Perraguin , R. Chudasama , S.I. Cooper , C. Crovella , G. Fidalgo , S.V. Gleyzer , A. Khukhunaishvili , K. Matchev , E. Pearson, C.U. Perez , P. Rumerio⁸¹ , E. Usai , R. Yi 

Boston University, Boston, Massachusetts, USA

G. De Castro, Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , J. Fulcher , F. Golf , S. Jeon , J. O'cain, I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , A.G. Zecchinelli 

Brown University, Providence, Rhode Island, USA

G. Barone , G. Benelli , D. Cutts , S. Ellis, L. Gouskos , M. Hadley , U. Heintz , K.W. Ho , J.M. Hogan⁸² , T. Kwon , G. Landsberg , K.T. Lau , J. Luo , S. Mondal , J. Roloff, T. Russell, S. Sagir⁸³ , X. Shen , M. Stamenkovic , N. Venkatasubramanian

University of California, Davis, Davis, California, USA

S. Abbott , B. Barton , C. Brainerd , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , O. Kukral , G. Mocellin , S. Ostrom , W. Wei , S. Yoo 

University of California, Los Angeles, California, USA

K. Adamidis, M. Bachtis , D. Campos, R. Cousins , A. Datta , G. Flores Avila , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , Y.f. Lo, E. Manca , A. Nunez Del Prado, D. Saltzberg , V. Valuev 

University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , G. Hanson 

University of California, San Diego, La Jolla, California, USA

A. Aportela, A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , L. Giannini , Y. Gu, J. Guiang , V. Krutelyov , R. Lee , J. Letts , H. Li, M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , D. Primosch, M. Quinnan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil , Z. Zhao

University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan , C. Campagnari , S. Carron Montero⁸⁴, K. Downham , C. Grieco , M.M. Hussain, J. Incandela , J. Kim , A.J. Li , P. Masterson , J. Richman , S.N. Santpur , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart , T.Á. Vámi , X. Yan , D. Zhang

California Institute of Technology, Pasadena, California, USA

A. Albert, S. Bhattacharya , A. Bornheim , O. Cerri, R. Kansal , J. Mao , H.B. Newman , G. Reales Gutiérrez, T. Sievert, M. Spiropulu , J.R. Vlimant , R.A. Wynne, S. Xie 

Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , P. Bryant , M. Cremonesi, V. Dutta , E.Y. Ertorer , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil, M. Kanemura, C. Liu , P. Meiring , T. Mudholkar , S. Murthy , P. Palit , K. Park, M. Paulini , A. Roberts , A. Sanchez , W. Terrill 

University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo 

Cornell University, Ithaca, New York, USA

J. Alexander , X. Chen , D.J. Cranshaw , J. Dickinson , J. Fan , X. Fan , J. Grassi , S. Hogan , P. Kotamnives, J. Monroy , G. Niendorf, M. Oshiro , J.R. Patterson , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou 

Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , C. Cosby , G. Cummings , I. Dutta , V.D. Elvira , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , T.C. Herwig , J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima , K.H.M. Kwok , S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , K. Maeshima , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg, V. Papadimitriou , N. Pastika , K. Pedro , C. Pena⁸⁵ , C.E. Perez Lara , F. Ravera , A. Reinsvold Hall⁸⁶ , L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , C. Wang , I. Zoi

University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , V. Sulimov , Y. Takahashi , J. Wang

Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhim , A. Askew , S. Bower , R. Hashmi , R.S. Kim , S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, M. Wulansatiti , R. Yohay

Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani , S. Butalla , S. Das , M. Hohlmann , M. Lavinsky, E. Yanes

University of Illinois Chicago, Chicago, Illinois, USA

M.R. Adams , N. Barnett, A. Baty , C. Bennett, R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , H. Gupta , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , D. S. Lemos , C. Mills , S. Nanda , G. Nigmatkulov , B. Ozek , T. Phan, D. Pilipovic , R. Pradhan , E. Prifti, P. Roy, T. Roy , N. Singh, M.B. Tonjes , N. Varelas , M.A. Wadud , J. Yoo

The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini , D. Blend, K. Dilsiz⁸⁷ , G. Karaman , O.K. Köseyan , A. Mestvirishvili⁸⁸ , O. Neogi, H. Ogul⁸⁹ , Y. Onel , A. Penzo , C. Snyder, E. Tiras⁹⁰ 

Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M.V. Srivastav , M. Swartz

The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , S. Popescu⁹¹ , C. Rogan , C. Royon , S. Rudrabhatla , S. Sanders , C. Smith , G. Wilson

Kansas State University, Manhattan, Kansas, USA

B. Allmond , R. Gujju Gurunadha , N. Islam, A. Ivanov , K. Kaadze , Y. Maravin 

J. Natoli , D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman, S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim, S. Lascio , P. Major , A.C. Mignerey , C. Palmer , C. Papageorgakis , M.M. Paranjpe, E. Popova⁹² , A. Shevelev , L. Wang , L. Zhang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

C. Baldenegro Barrera , J. Bendavid , S. Bright-Thorney , I.A. Cali , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, G. Grosso, P. Harris, D. Hoang, G.M. Innocenti, D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn , A. Novak , M.I. Park , C. Paus , C. Reissel , C. Roland , G. Roland , S. Rothman , G.S.F. Stephans , D. Walter , Z. Wang , B. Wyslouch , T. J. Yang 

University of Minnesota, Minneapolis, Minnesota, USA

B. Crossman , W.J. Jackson, C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , M. Revering , R. Rusack , O. Sancar, R. Saradhy , N. Strobbe 

University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , A. Rohilla , J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 

State University of New York at Buffalo, Buffalo, New York, USA

H. Bandyopadhyay , L. Hay , H.w. Hsia , I. Iashvili , A. Kalogeropoulos , A. Kharchilava , A. Mandal , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams , P. Young 

Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , J. Bonilla , B. Bylsma, M. Campana , J. Dervan , Y. Haddad , Y. Han , I. Israr , A. Krishna , J. Li , M. Lu , N. Manganelli , R. McCarthy , D.M. Morse , T. Orimoto , A. Parker , L. Skinnari , C.S. Thoreson, E. Tsai , D. Wood 

Northwestern University, Evanston, Illinois, USA

S. Dittmer , K.A. Hahn , Y. Liu , M. McGinnis , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliencio , M. Velasco, J. Wang 

University of Notre Dame, Notre Dame, Indiana, USA

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth , K. Hurtado Anampa , T. Ivanov , C. Jessop , A. Karneyeu , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , C. Moore , Y. Musienko²³ , H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan, M. Wayne , H. Yockey, L. Zygala 

The Ohio State University, Columbus, Ohio, USA

A. Basnet , M. Carrigan , R. De Los Santos , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas , K. Wei, D.A. Wenzl, B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, USA

H. Bouchamaoui , K. Coldham, P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , Y. Lai , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , F. Simpson , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , R. Sharma

Purdue University, West Lafayette, Indiana, USA

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung , D. Kondratyev , M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , J.F. Schulte , F. Wang , A. Wildridge , W. Xie , Y. Yao , Y. Zhong 

Purdue University Northwest, Hammond, Indiana, USA

J. Dolen , N. Parashar , A. Pathak 

Rice University, Houston, Texas, USA

D. Acosta , A. Agrawal , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner, F.J.M. Geurts , T. Huang , I. Krommydas , N. Lewis, W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 

University of Rochester, Rochester, New York, USA

O. Bessidskaia Bylund, A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , A. Garcia-Bellido , H.S. Hare, O. Hindrichs , N. Parmar , P. Parygin⁹² , R. Taus 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis , M. Heindl , C. Houghton , D. Jaroslawski , S. Konstantinou , I. Laflotte , A. Lath , J. Martins , B. Rand, J. Reichert , P. Saha , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora 

University of Tennessee, Knoxville, Tennessee, USA

D. Ally , A.G. Delannoy , S. Fiorendi , J. Harris, S. Higginbotham , T. Holmes , A.R. Kanuganti , N. Karunarathna , J. Lawless, L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , T. Akhter , K. Androsov , A. Bolshov, O. Bouhali⁹³ , R. Eusebi , J. Gilmore , T. Kamon , H. Kim , S. Luo , R. Mueller , A. Safonov 

Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , Y. Feng , N. Gogate , Y. Kazhykarim, K. Lamichhane , S.W. Lee , C. Madrid , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , D. Rathjens , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen , J. Zhang

University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , H. Chung, B. Cox , J. Hakala , R. Hirosky , M. Jose, A. Ledovskoy , C. Mantilla , C. Neu , C. Ramón Álvarez 

Wayne State University, Detroit, Michigan, USA

S. Bhattacharya , P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind , S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, S. Lomte, R. Loveless , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , L. Pétré , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi 

W. Vetens , A. Warden 

Authors affiliated with an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , Yu. Andreev , T. Aushev , D. Budkouski , R. Chistov⁹⁴ , M. Danilov⁹⁴ , T. Dimova⁹⁴ , A. Ershov⁹⁴ , S. Gninenko , I. Golutvin[†] , I. Gorbunov , A. Gribushin⁹⁴ , A. Kamenev , V. Karjavine , M. Kirsanov , V. Klyukhin⁹⁴ , O. Kodolova^{95,92} , V. Korenkov , A. Kozyrev⁹⁴ , N. Krasnikov , A. Lanev , A. Malakhov , V. Matveev⁹⁴ , A. Nikitenko^{96,95} , V. Palichik , V. Perelygin , S. Petrushanko⁹⁴ , S. Polikarpov⁹⁴ , O. Radchenko⁹⁴ , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen⁹⁴ , V. Smirnov , O. Teryaev , I. Tlisova⁹⁴ , A. Toropin , N. Voytishin , B.S. Yuldashev^{†97}, A. Zarubin , I. Zhizhin

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

L. Dudko , K. Ivanov , V. Kim²³ , V. Murzin , V. Oreshkin , D. Sosnov 

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Ghent University, Ghent, Belgium

⁴Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁵Also at FACAMP - Faculdades de Campinas, Sao Paulo, Brazil

⁶Also at Universidade Estadual de Campinas, Campinas, Brazil

⁷Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁸Also at The University of the State of Amazonas, Manaus, Brazil

⁹Also at University of Chinese Academy of Sciences, Beijing, China

¹⁰Also at China Center of Advanced Science and Technology, Beijing, China

¹¹Also at University of Chinese Academy of Sciences, Beijing, China

¹²Now at Henan Normal University, Xinxiang, China

¹³Also at University of Shanghai for Science and Technology, Shanghai, China

¹⁴Now at The University of Iowa, Iowa City, Iowa, USA

¹⁵Also at Center for High Energy Physics, Peking University, Beijing, China

¹⁶Also at Helwan University, Cairo, Egypt

¹⁷Now at Zewail City of Science and Technology, Zewail, Egypt

¹⁸Now at British University in Egypt, Cairo, Egypt

¹⁹Now at Cairo University, Cairo, Egypt

²⁰Also at Purdue University, West Lafayette, Indiana, USA

²¹Also at Université de Haute Alsace, Mulhouse, France

²²Also at Istinye University, Istanbul, Turkey

²³Also at an institute formerly covered by a cooperation agreement with CERN

²⁴Also at University of Hamburg, Hamburg, Germany

²⁵Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²⁶Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

²⁷Also at Brandenburg University of Technology, Cottbus, Germany

²⁸Also at Forschungszentrum Jülich, Jülich, Germany

²⁹Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

³⁰Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

³¹Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania

³²Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

³³Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

- ³⁴Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- ³⁵Also at The University of Kansas, Lawrence, Kansas, USA
- ³⁶Also at Punjab Agricultural University, Ludhiana, India
- ³⁷Also at Indian Institute of Science (IISc), Bangalore, India
- ³⁸Also at University of Visva-Bharati, Santiniketan, India
- ³⁹Also at IIT Bhubaneswar, Bhubaneswar, India
- ⁴⁰Also at Institute of Physics, Bhubaneswar, India
- ⁴¹Also at University of Hyderabad, Hyderabad, India
- ⁴²Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ⁴³Also at Isfahan University of Technology, Isfahan, Iran
- ⁴⁴Also at Sharif University of Technology, Tehran, Iran
- ⁴⁵Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴⁶Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁴⁷Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁴⁸Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁴⁹Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁵⁰Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- ⁵¹Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- ⁵²Also at Lulea University of Technology, Lulea, Sweden
- ⁵³Also at Laboratori Nazionali di Legnaro dell'INFN, Legnaro, Italy
- ⁵⁴Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵⁵Also at UPES - University of Petroleum and Energy Studies, Dehradun, India
- ⁵⁶Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
- ⁵⁷Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁸Also at INFN Sezione di Torino, Università di Torino, Torino, Italy; Università del Piemonte Orientale, Novara, Italy
- ⁵⁹Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁶⁰Also at Saegis Campus, Nugegoda, Sri Lanka
- ⁶¹Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁶²Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶³Also at Universität Zürich, Zurich, Switzerland
- ⁶⁴Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶⁵Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶⁶Also at Konya Technical University, Konya, Turkey
- ⁶⁷Also at Izmir Bakircay University, Izmir, Turkey
- ⁶⁸Also at Adiyaman University, Adiyaman, Turkey
- ⁶⁹Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey
- ⁷⁰Also at Marmara University, Istanbul, Turkey
- ⁷¹Also at Milli Savunma University, Istanbul, Turkey
- ⁷²Also at Tubitak, Kavaklıdere, Ankara, Turkey
- ⁷³Also at Kafkas University, Kars, Turkey
- ⁷⁴Now at Istanbul Okan University, Istanbul, Turkey
- ⁷⁵Also at Hacettepe University, Ankara, Turkey
- ⁷⁶Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- ⁷⁷Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey

⁷⁸Also at Yildiz Technical University, Istanbul, Turkey

⁷⁹Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom

⁸⁰Also at Monash University, Faculty of Science, Clayton, Australia

⁸¹Also at Università di Torino, Torino, Italy

⁸²Also at Bethel University, St. Paul, Minnesota, USA

⁸³Also at Karamanoğlu Mehmetbey University, Karaman, Turkey

⁸⁴Also at California Lutheran University, Thousand Oaks, California, USA

⁸⁵Also at California Institute of Technology, Pasadena, California, USA

⁸⁶Also at United States Naval Academy, Annapolis, Maryland, USA

⁸⁷Also at Bingöl University, Bingöl, Turkey

⁸⁸Also at Georgian Technical University, Tbilisi, Georgia

⁸⁹Also at Sinop University, Sinop, Turkey

⁹⁰Also at Erciyes University, Kayseri, Turkey

⁹¹Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania

⁹²Now at another institute formerly covered by a cooperation agreement with CERN

⁹³Also at Texas A&M University at Qatar, Doha, Qatar

⁹⁴Also at another institute formerly covered by a cooperation agreement with CERN

⁹⁵Also at Yerevan Physics Institute, Yerevan, Armenia

⁹⁶Also at Imperial College, London, United Kingdom

⁹⁷Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan