

CERN-EP-2025-041  
2025/12/16

CMS-SMP-23-008

# Measurement of event shapes in minimum-bias events from proton-proton collisions at $\sqrt{s} = 13$ TeV

The CMS Collaboration\*

## Abstract

A measurement of event-shape variables is presented, using a data sample produced in a special run with approximately one inelastic proton-proton collision per bunch crossing. The data were collected with the CMS detector at a center-of-mass energy of 13 TeV, corresponding to an integrated luminosity of  $64 \mu\text{b}^{-1}$ . A number of observables related to the overall distribution of charged particles in the collisions are corrected for detector effects and compared with simulations. Inclusive event-shape distributions, as well as differential distributions of event shapes as functions of charged-particle multiplicity, are studied. None of the models investigated are able to satisfactorily describe the data. Moreover, there are significant features common amongst all generator setups studied, particularly showing data being more isotropic than any of the simulations. Multidimensional unfolded distributions are provided, along with their correlations.

*Published in Physical Review D as doi:10.1103/rqvb-gfb7.*



# 1 Introduction

Recent measurements of observables in proton-proton (pp) collisions related to the overall structure of the event, such as the correlations of outgoing particles [1–4] and the multiplicity of strange hadrons [5, 6], highlight the difficulties in modeling these collisions. Models that attempt to describe these effects, for instance those developed in the context of heavy ion collisions, include nonperturbative effects, such as collective flow [7–12] or rope hadronization [13–15]. This motivates the continued measurement program of minimum-bias pp collisions and the study of observables sensitive to nonperturbative effects. Furthermore, the dominance of soft quantum chromodynamics (QCD) effects, even in high-precision electroweak observables, such as the anomalous magnetic moment of the muon [16, 17], highlights the need for additional experimental input to nonperturbative QCD.

One nonperturbative phenomenon of particular interest is the QCD instanton [18–26]. Recent calculations [27] suggest that instanton processes at the CERN LHC may have a large cross section of  $\mathcal{O}(\text{mb})$ . Several strategies for detecting instanton events have been suggested [28–32], and one of the common characteristics used in many of those searches is the relatively isotropic distribution of particles produced by the instanton.

While we do not provide direct limits or instanton interpretation, these effects and other similar modifications of event shapes motivate our measurement as a means to test the limits of the existing pp collision modeling. In particular, we study minimum-bias collisions that define a set of nearly inclusive collision events, with minimal selection criteria applied to ensure the collisions and outgoing particles are well measured by the detector.

This paper presents measurements of observables related to the overall kinematics of events, often referred to as event shapes. The measurements use minimum-bias pp collisions at a center-of-mass energy of 13 TeV collected by the CMS detector at the CERN LHC, corresponding to an integrated luminosity of  $64 \mu\text{b}^{-1}$  [33, 34]. We study eight observables related to the global event shapes using tracks reconstructed in the detector (detector level). Then we unfold the same observables to the level of stable charged particles (particle level). Similar observables have been measured in  $e^+e^-$  collisions at the CERN LEP collider [35] and were compared with simulations in previous CMS event generator tuning studies [36]. The measurements in this paper include effects of the hadronic initial state not present in those measurements. The charged-particle multiplicity has previously been studied at 13 TeV center-of-mass energy [37–39]. Some of the other observables have also been studied at the CERN LHC in the underlying event in top quark pair ( $t\bar{t}$ ) production [40], or using jets as inputs [41], and at lower center-of-mass energies [42–44]. The measurements presented here extend those analyses and elucidate correlations between the event shapes and the charged-particle multiplicity. The observables are always computed using tracks (detector level) or charged particles (particle level). Sphericity, thrust, broadening and isotropy are computed in the center-of-mass frames of the tracks or charged particles for each event. The full list of observables is as follows:

- Particle multiplicity,  $N$ .
- Total invariant mass,  $m_{\text{ch}} = \sqrt{(\sum_i p_i)^2}$ , where  $p_i$  is the four-momentum of particle  $i$ .
- Sphericity: a measure of how isotropically the momenta  $p$  are distributed in an event [45, 46]. The tensor  $S$  is first defined with components

$$S^{\alpha\beta} = \frac{\sum_i p_i^\alpha p_i^\beta}{\sum_i |\vec{p}_i|^2}, \quad (1)$$

in which  $\alpha, \beta \in \{x, y, z\}$  refer to Cartesian coordinates, and  $i$  is the index for the

final-state charged particles that passed the selections based on the detector acceptance, described in Section 4. The sphericity is constructed from the two smallest eigenvalues  $\lambda_2$  and  $\lambda_3$ :  $\mathcal{S} = \frac{3}{2}(\lambda_2 + \lambda_3)$ .

- Thrust: a measure of how highly collimated the momenta in an event are along one particular axis, the thrust axis [46]. It is defined as

$$\mathcal{T} = 1 - \max_{\vec{n}} \frac{\sum_i \vec{p}_i \cdot \vec{n}}{\sum_i |\vec{p}_i|}, \quad (2)$$

where  $\vec{n}$  is a unit vector, which is chosen to maximize the numerator [47].

- Transverse thrust: the thrust in the transverse plane, calculated by setting  $p_i^z$  to 0 in the thrust definition (Eq. 2).
- Broadening: a measure of the fraction of energy that is perpendicular to the thrust axis. The thrust axis defines the left  $\mathcal{L}$  and right  $\mathcal{R}$  hemispheres of the event. The left and right broadening are defined as

$$\mathcal{B}_{\mathcal{L}} = \sum_{i \in \mathcal{L}} \frac{|\vec{p}_i \times \vec{n}|}{2 \sum_j |\vec{p}_j|}, \quad \mathcal{B}_{\mathcal{R}} = \sum_{i \in \mathcal{R}} \frac{|\vec{p}_i \times \vec{n}|}{2 \sum_j |\vec{p}_j|}. \quad (3)$$

The total broadening is defined as  $\mathcal{B} = \mathcal{B}_{\mathcal{L}} + \mathcal{B}_{\mathcal{R}}$ .

- Transverse sphericity: the sphericity in the transverse plane, a measure of how spread out the momentum is across directions in the transverse plane,

$$\mathcal{S}_0 = \frac{\pi^2}{4} \min_{\vec{n}=(n_x, n_y, 0)} \left( \frac{\sum_i |\vec{p}_{Ti} \times \vec{n}|}{\sum_i p_{Ti}} \right)^2. \quad (4)$$

- Isotropy: a measure of how isotropically energy is distributed in an event. It is defined as the energy mover's distance [48] from the spatial distribution of the tracks to a uniform radiation pattern [49]. The detailed definition of this variable is given in Ref. [50], which also explores the efficacy of the isotropy to distinguish the isotropic events from the jetlike ones characterized by the majority of the transverse momentum being clustered along a few axes. For this measurement we use the spherical geometry, with the distance measure  $d_{ij} = 2(1 - \cos \theta_{ij})$ , where  $\theta_{ij}$  is the angle between the particles.

We investigate a number of minimum-bias simulations from generators including HERWIG (v7.2.2) [51, 52], EPOS-LHC [7], and PYTHIA (v8.230) [53], with multiple different tunes of PYTHIA, to evaluate the current quality of the modeling of event-shape distributions, and to provide a reference for the future search for currently unmodeled instanton contributions to these observables.

A machine learning (ML)-based unfolding algorithm [54, 55] is used to correct for detector effects in the observed distributions and to estimate the distributions for stable charged particles. This ML-based approach is a multidimensional unfolding, which follows a similar approach to traditional binned iterative unfolding, but extended to an unbinned analysis.

For each event, the value of every unfolded observable is provided and a simultaneous unfolding of all observables is performed. This allows the full joint information of all unfolded observables to be taken into account simultaneously in the unfolding and for further selections or binning choices to be made after the unfolding is performed.

The histograms of individual event-shape observables are provided in Section 7, as well as two-dimensional (2D) histograms of event-shape observables as functions of the number of charged particles and their invariant mass in Section 7 and Appendix B. Correlation matrices for the systematic uncertainties of the histogrammed event-shape observables are provided in Appendix C. Tabulated results are provided in the HEPData record [56], and the unbinned unfolding results are provided in the CERN Document Server [57].

## 2 The CMS detector and object reconstruction

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field strength of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two end-cap sections. Forward calorimeters extend the pseudorapidity coverage provided by the barrel and end-cap detectors. Muons are measured in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Refs. [58, 59].

The primary vertices (PVs) are estimates of the positions of the pp collisions reconstructed from the detector objects. We use the standard CMS vertex reconstruction algorithm [60], which clusters prompt tracks, associates them with PVs, and then fits the PV positions from the tracks. A particle-flow algorithm [61] aims to reconstruct and identify each individual particle in an event, with an optimized combination of all subdetector information. In this process, the identification of the particle type (photon, electron, muon, charged hadron, or neutral hadron) plays an important role in the determination of the particle direction and energy. Electrons (e.g., coming from photon conversions in the tracker material or from B meson semileptonic decays) are identified as a primary charged-particle track and potentially many ECAL energy clusters corresponding to this track followed to the ECAL and to possible bremsstrahlung photons emitted along the way through the tracker material. Muons (e.g., from B meson semileptonic decays) are identified as tracks in the central tracker consistent with either a track or several hits in the muon system, and associated with calorimeter deposits compatible with the muon hypothesis. Charged hadrons are identified as charged-particle tracks, neither identified as electrons nor as muons.

The energy of electrons is determined from a combination of the track momentum at the main interaction vertex, the corresponding ECAL cluster energy, and the energy sum of all bremsstrahlung photons attached to the track. The energy of muons is obtained from the corresponding track momentum. The energy of charged hadrons is determined from a combination of the track momentum and the corresponding ECAL and HCAL energies, corrected for the response function of the calorimeters to hadronic showers.

## 3 Data and simulated samples

Data were collected with the CMS detector in 2018 under low-pileup conditions, corresponding to an average number of inelastic pp collisions per bunch crossing of approximately 1. The usage of the low-pileup data mitigates pileup effects in this precision measurement [62]. Despite the low luminosity of this dataset, the measurement is not statistically limited. Only a fraction of the total data collected by the CMS detector in 2018 is used, corresponding to 7

million events and an integrated luminosity of  $64 \mu\text{b}^{-1}$  [33, 34]. An unbiased trigger is used to randomly select collision events that are recorded independently of the event content [63].

Several Monte Carlo (MC) simulations of minimum-bias pp collisions are used to perform the unfolding and compare with the observed data. One nominal choice of minimum-bias event generation setup is used as the central prediction, while the others are used for estimating uncertainties as well as for comparison and interpretation. Some details of the nominal simulation sample and of those used for estimating systematic uncertainties are listed in Table 1. In all cases, GEANT4 [64] is used for the detector simulation. For most samples, approximately 2 million unweighted events are used.

The PYTHIA CP1 tune [36] is chosen as the nominal model for its relatively good description of the observed distributions of track multiplicity and track pseudorapidity to minimize uncertainties and bias in the unfolding due to mismodeling. The PYTHIA A3 tune [65], EPOS-LHC [7], and the HERWIG CH3 tune [66] were chosen as systematic variations that provide a similar level of agreement with the observations, while also covering different tuning and physics model choices. PYTHIA and HERWIG are both general purpose MC event generators commonly used to perform pp collision analyses. EPOS-LHC uses a core-corona event model for hadronic collisions originally developed in the context of heavy ion collisions but adapted for pp collisions and includes some LHC data in the parameter tuning.

Table 1: Samples of simulated minimum-bias events used for the unfolding. The value of the strong coupling constant,  $\alpha_s$ , corresponds to that used in the parton distribution function (PDF) set. The samples are simulated without additional pileup interactions.

|                       | Generator           | PDF ( $\alpha_s(m_Z)$ )     |
|-----------------------|---------------------|-----------------------------|
| Nominal sample        | PYTHIA8 (CP1) [36]  | NNPDF3.1 QCD LO (0.130)     |
| Systematic variations | EPOS-LHC [7]        |                             |
|                       | PYTHIA8 (A3) [65]   | NNPDF2.3 QCD+QED LO (0.130) |
|                       | HERWIG 7 (CH3) [66] | NNPDF3.1 QCD NNLO (0.118)   |

Additional MC simulations are listed in Appendix A, which are used for cross-checks and validation but are not included in the main analysis. Some of these samples were used as pseudodata samples to validate the performance of the unfolding against known particle-level distributions, while others were investigated to ensure that they did not give substantially different results to the samples used in the analysis. The main samples used in the analysis are all simulated without additional pileup interactions. A simulated PYTHIA sample with the CP5 tune with pileup drawn from a Poisson distribution of mean 1 was used to check the impact of pileup on the observables—the effects were negligible.

## 4 Event and object selections

This analysis is performed using charged particles (primarily charged hadrons) without clustering into jets. All calculations are performed directly with tracks associated with particle-flow candidates and reconstructed vertices without distinguishing leptons and hadrons.

The analysis uses a minimal set of selections at detector level, applied to both data and simulation, to select events with one inelastic proton-proton collision. A selection is made on the PV, such that it is within a broad region around the nominal interaction point: within  $\pm 24$  cm along the beamline direction and within  $\pm 2$  cm in the transverse plane. Furthermore, the vertex position must be well reconstructed from the tracks associated with it, as determined by

the number and position of tracks associated with the vertex [60]. Every event is required to have only one such PV and at least three tracks fulfilling the track selection criteria associated with the PV. All tracks are required to have transverse momentum  $p_T > 0.5 \text{ GeV}$ , pseudorapidity  $|\eta| < 2.4$ , and loose selections are placed on the track impact parameters based on the vertex resolution: the track impact parameter in the transverse (longitudinal) direction must be at most 8 (13) times the vertex resolution in the transverse (longitudinal) direction.

A minimal particle-level event selection defines the fiducial phase space that is the target of the unfolding. These selections are intended to define a minimally biased set of inelastic collisions, while minimizing extrapolation between the detector-level selection and the particle-level phase space. Events are required to have at least two charged particles with  $p_T > 0.5 \text{ GeV}$  and  $|\eta| < 2.4$ . The particles selected at the particle level are electrically charged particles and required to be stable (lifetime  $c\tau > 10 \text{ mm}$ , corresponding to “status 1” particles of the HEPMC standard [67]). Simulated events that pass the detector-level selection but do not meet the fiducial phase space selection criteria are included in the analysis and treated as background.

## 5 Unfolding procedure

We use an ML-based unfolding procedure, MULTIFOLD, which performs an unbinned, multi-dimensional unfolding and uses iteration as its method of regularization [54, 55].

The MULTIFOLD algorithm is similar to the Richardson-Lucy unfolding [68, 69], also known as iterative Bayesian unfolding or the d’Agostini method [70], but uses ML to extend the methods to unbinned cases. The algorithm is based on the fact that, when performing a classification to identify an event as coming from one sample or another, an ideal classifier will learn the likelihood ratio, i.e. the ratio of the probability of observing such an event given that it is from one class or the other. Therefore, a sufficiently well-designed and trained neural network classifier should approximate the likelihood ratio [71]. The likelihood ratio estimated from the classifier is then used to reweight events from one sample to resemble the other sample in an unbinned way.

The unfolding uses multiple reweighting steps for each iteration, first correcting the simulation toward the data at the detector level, then correcting the original simulation toward this corrected simulation at the particle level. Additional steps are also employed to account for acceptance and efficiency effects, as described in Ref. [55].

These steps are iterated a number of times, with the iteration acting as a form of regularization. Fewer iterations correspond to more regularization and tend to be more biased toward the input simulated distribution. The number of iterations used in the unfolding was chosen by checking goodness-of-fit measures of the unfolded result on a simulated pseudodata sample from an alternative simulation. The goodness of fit of the histograms reported in Section 7 was calculated using a  $\chi^2$  for each observable, and the number of iterations was chosen such that larger numbers of iterations did not provide any clear improvement in the goodness of fit. Both the bias of the unfolded estimates and the frequency with which the stated uncertainty intervals contain the true value of the parameter being estimated (coverage) converge quickly, and no significant changes were found to these properties after two iterations.

For every event, the neural networks are provided with the values of all observables being unfolded, such that all observables are simultaneously unfolded, and their correlations are taken into account. The values passed for every event are the number of objects, their invariant mass, sphericity, thrust, transverse thrust, broadening, transverse sphericity, and isotropy, as

defined in Section 1. The objects used to calculate the observables are the tracks for detector-level reweighting, or charged particles for particle-level reweighting.

The classification networks consist of three fully connected layers of 100 nodes each. The cross-entropy, which measures the ability of the network to identify the sample from which an event originates, is used as the loss function. The training is performed for every step of the procedure and every iteration, and is done using 80% of the sample for training and 20% for validation. Early stopping is used, where the training is stopped after no improvement in the loss function evaluated on the validation sample is seen for 10 epochs (complete passes through the training sample). Then the network parameters are taken from the last epoch that showed improvement in the training. Different initial values of the network weights during training can lead to random fluctuations in the output. This effect was mitigated by an ensembling technique: four trainings and evaluations with independent initializations of the neural network weights are performed, and the final event weight is the average of the ensemble. The unfolding is performed using two iterations.

As an input to the unfolding, the nominal MC simulated events have uniform weights that normalize them to the total inelastic cross section of 13 TeV pp collisions reported by CMS [72]. All MC samples are weighted such that the sum of weights for the events passing the particle-level selection is equal to that for the nominal MC sample. The weighting ensures a fixed fiducial cross section for the nominal MC and its variations at the start of the unfolding.

In each reweighting step of the training, the program uses balanced weights between the set to be weighted and the target, so that the weighting function from the ML model is based on only the shape differences between the distributions of the two sets. The weights from the unfolding are multiplied by the previous ones in each iteration.

The final result of the unbinned unfolding is a set of reweighted MC events, whose weights provide an estimate of the probability density of the observed data at each point in the phase space. These results are binned after the unfolding to provide numerous histograms displaying the estimated differential cross sections as a function of the observables of interest.

A number of tests have been performed to validate the unfolding procedure. For these tests, one or more of the alternative simulated samples listed in Table A.1 is treated as pseudodata to be unfolded. As well as visual checks of the unfolding closure, binned goodness-of-fit tests are performed to validate that the unfolded results provide a better description of the pseudodata than the original simulation. A bottom-line test [73] is performed to ensure that the unfolding does not artificially introduce discriminating power between models that was not already present at the detector level. Frequentist bias and coverage tests were also performed using parametric sampling [74]. For most bins, the estimated bias is compatible with zero, and in all cases it is considerably smaller than the overall uncertainty. No cases of undercoverage were observed.

## 6 Uncertainties

The statistical uncertainty in the observed data is estimated by a bootstrapping method [75]. This method first assigns weights following a Poisson(1) distribution to the data events to generate alternative data samples with statistical fluctuations, and then performs pseudoexperiments of unfolding these alternative data samples. The standard deviations of the unfolded results become estimations of the statistical uncertainty.

The effects of systematic uncertainties are also estimated using pseudoexperiments, which start



from templates defining the shifts in the unfolding model from individual systematic uncertainty sources. The templates define these shifts as weights on the events of the nominal MC sample. The sources of the systematic uncertainty and the estimation methods are as follows:

- MC modeling: modeling of the underlying physics can affect the result of the unfolded measurements in several distinct ways. We consider two distinct effects:
  - The regularization, which is induced by the limited number of iterations in the unfolding algorithm and imperfections in the neural network classifier performance can lead to biased results. This bias depends on the choice of the nominal MC model. To estimate the effect of this systematic uncertainty, alternative templates are estimated by weighting the nominal MC to the alternative MC samples from other tunes or generators at the particle level. The resulting reweighted nominal sample has the same particle-level distributions as the alternative MC but maintains the same migration function as the nominal ones.
  - Differences in the underlying physics modeling of quantities that are not unfolded, for example the track pseudorapidity distribution, can also affect the detector response and thus lead to a poor modeling of the migration from the particle-level distributions to the detector-level ones. The templates of this systematic deviation are derived in two steps. In the first step, the alternative MC samples are weighted to the nominal sample at the particle level, which results in reweighted alternative MC samples with the same particle-level distributions as the nominal MC and the same migration function as the alternative MC. In the second step, the nominal sample is weighted at both the particle level and the detector level to the reweighted alternative MC samples from the first step. The second step results in reweighted nominal samples with the same particle-level and detector-level distributions as the output from the first step. The function of the second step is to transfer the weights on the alternative MC samples to those on the nominal samples, keeping the distributions the same. The resulting reweighted nominal samples have the same particle-level distribution as the nominal MC and the same migration function as the alternative MC.

For each of these two effects, we consider multiple alternative MC models as sources of uncertainty. Specifically, we consider one alternative PYTHIA tune as well as the EPOS-LHC and HERWIG setups, as described in Section 3. Therefore, we have a total of six independent modeling uncertainties. Alternative choices of systematic samples coming from generator tune eigenvariations or other models with reasonable descriptions of the data were also evaluated and they resulted in similar sizes of the uncertainties.

- Track reconstruction efficiency: this uncertainty accounts for potential deviations of the track reconstruction efficiency in simulations from the actual efficiency in observations. To estimate the effect of this uncertainty in the event-shape measurement, we randomly drop 2.1% of the tracks with  $p_T < 20$  GeV and 1% of those with  $p_T > 20$  GeV in the simulation. These numbers are based on the tracking efficiency studies in the  $D^*(2010)^\pm$  meson reconstruction measurement [76, 77]. The template of this systematic deviation is estimated by weighting the nominal MC samples to the samples with randomly dropped tracks at the particle and detector levels.
- Statistical uncertainty of the simulated model: the statistical uncertainty of the MC

sample leads to a systematic uncertainty in the unfolding results. This uncertainty is estimated by the same bootstrapping method that is used in estimating the uncertainty from the event counts of observed data.

The effects of pileup were negligible.

For the MC regularization bias, migration function, and track reconstruction uncertainties, we derived a set of weights on the nominal MC simulation, such that the effect of the uncertainty can be described as a function of a continuous nuisance parameter. The weights are derived from the alternative samples using a ML-reweighting procedure similar to the reweighting procedure used in the unfolding. A neural network classifier is trained to distinguish between the nominal and alternative samples and then the classifier score is used to define the reweighting value. The nuisance parameters for each uncertainty are assumed to be Gaussian-constrained, and the weight of any event,  $w_i$ , as a function of a given nuisance parameter,  $\theta$ , is given by

$$w_i(\theta) = w_{\text{nom}} (w_i(1)/w_{\text{nom}})^\theta, \quad (5)$$

in which  $w_{\text{nom}}$  is the weight of the nominal MC event, and  $w_i(1)$  is the weight of the event in the corresponding systematic uncertainty template, which is defined to be the +1 standard deviation from the nominal.

The total uncertainty and its covariance matrices are estimated by 100 pseudoexperiments of unfolding with weighted nominal MC simulation and weighted data separately, which emulate the fluctuations of the templates. For the data and MC statistical uncertainty, the weights in the pseudoexperiments are taken from the Poisson(1) distribution and applied to the data and MC samples, respectively. For the MC bias, migration function, and track reconstruction efficiency uncertainties, a nuisance parameter value  $\theta_j \sim N(0, 1)$  is generated for each template  $j$ . The total weight of any given MC event for each pseudoexperiment is the product of the weights from all the sources of systematic uncertainty.

The systematic uncertainties in the results are larger than the statistical uncertainties for the chosen bin widths in this paper. For one-dimensional (1D) distributions, the typical uncertainty is  $\mathcal{O}(5\%)$ , depending on the kinematic bin. The uncertainty from tracking efficiency is typically the largest, of the order of a few percent. The uncertainty from the migration functions for different physics models combined across all models is of a similar size. The uncertainty from the estimated regularization bias between the nominal model and other models is typically  $\mathcal{O}(1\%)$  or less for each model, and the combined uncertainty from regularization bias from all models is typically a little smaller than the uncertainties from tracking efficiency or migration functions of the models. For the bin widths presented here, the statistical uncertainties in the 1D distributions are significantly less than 1%.

## 7 Results

Projections of the full unfolded distribution into 1D and 2D histograms are provided and discussed below. Additional 2D histograms and the correlation matrices of the systematic uncertainties for the 1D projections are given in Appendices B and C. The uncertainty decomposition of the unfolding is given in Appendix D.

Figures 1–3 show the distributions of unfolded observables in 1D using the PYTHIA CP1 tune as the nominal model.

The number of charged particles and their invariant mass, shown in Fig. 1, are different in data and simulation. The trend of the simulation fluctuates around the data, and does not follow

a single monotonic trend. All generators tend to overestimate the relative number of events at low charged-particle multiplicity ( $\lesssim 10$ ), compensated by an underestimate at intermediate values. At high multiplicity and masses, the predictions diverge from each other, and the predictions span the range from underestimating to overestimating the data. The same trend is observed for both the number of charged particles and their invariant mass.




Figure 1: The unfolded results for (left) the particle multiplicity, and (right) invariant mass, compared with the nominal MC from the PYTHIA CP1 tune and predictions from the PYTHIA A14, CP5, A3 tunes, the EPOS-LHC generator, and the HERWIG CH3 tune. The statistical uncertainty on each of the predictions is shown by its shaded band.

The event-shape observables shown in Fig. 2 show a consistent trend in which the unfolded data are more isotropic than the simulation. This trend is present for all measures of the event shapes used here, and for all generator configurations considered. Although the level of agreement, particularly in the tails of the distributions, depends on the observables and generator setup being considered, the overall trend is consistent and none of the simulations correctly capture the data.

Event-shape observables using only the transverse component of particle momenta are shown in Fig. 3. They show very similar trends to the event-shape observables that include longitudinal momentum information, with the data distributions being more isotropic than the simulations.

Figure 4 shows the unfolded sphericity and transverse sphericity distributions in slices of the charged-particle multiplicity. Other observables are included in Appendix B.

Within most slices of charged-particle multiplicity, the same general trend of the data being more isotropic than the simulation is present. This occurs over different charged-particle multiplicity slices where the trend of data to simulation in each slice differs, suggesting that the poor modeling of event shapes is likely not primarily coming from the distribution of the number of charged particles. This observation has also been confirmed by looking at slices with a single charged-particle multiplicity, and again is consistent across generator setups.




Figure 2: The unfolded results for (upper left) sphericity, (upper right) thrust, (lower left) broadening, and (lower right) isotropy compared with the nominal MC from the PYTHIA CP1 tune and MC predictions from the PYTHIA A14, CP5, A3 tunes, the EPOS-LHC generator, and the HERWIG CH3 tune. The statistical uncertainty on each of the predictions is shown by its shaded band.

Checks with the generator setups listed in Table A.1, including different tunes, PDF variations, and variations of  $\alpha_s$  used in the parton shower, all show the same qualitative behavior.

Comparisons to event-shape measurements performed at LEP [35] were previously performed




Figure 3: The unfolded results for (left) transverse thrust, and (right) transverse sphericity compared with the nominal MC from the PYTHIA CP1 tune and predictions from the PYTHIA A14, CP5, A3 tunes, the EPOS-LHC generator, and the HERWIG CH3 tune. The statistical uncertainty on each of the predictions is shown by its shaded band.

for the CMS PYTHIA tunes [36], where very different trends were observed. In those comparisons, which used measurements done in  $e^+e^- \rightarrow q\bar{q}$  events, the CP2 tune, very similar to the CP1 tune used in this analysis, was found to predict too many isotropic events. The opposite trends of modeling deficiencies for the event shapes of hadrons from  $e^+e^-$  collisions at LEP and from pp collisions in this measurement suggest that the poor modeling observed in this analysis is not related to the hadronic final-state evolutions to a large extent, but instead related to the initial states of pp collisions.

Furthermore, the effect of varying  $\alpha_s$  in the final-state shower was checked by using the CP5  $\alpha_s$  variation tunes [36], which were derived by fitting the  $\alpha_s$  parameter while fixing other values to their CP5 settings, and then taking the variations corresponding to the bounds of the 68% confidence interval estimated for  $\alpha_s$  from the fit. The predictions for the measured distributions have only minor changes, less than what is observed for other variations of settings shown here. The CUEP8M1 [78] and CUEP8M2T4 [40, 79] tunes both gave results very similar to those for the CP1 tune. The PYTHIA 6 Z2Star tune [80] shows larger discrepancies with the measured distributions than the models considered in the figures, probably because it was based on measurements of pp collisions at energies lower than 13 TeV.

These results are consistent with what was observed in a study of underlying event modeling in  $t\bar{t}$  events at CMS [40], where the underlying event simulations were observed to be somewhat less isotropic than the recorded data.




Figure 4: The unfolded distributions of sphericity (upper) and transverse sphericity (lower) in slices of charged-particle multiplicity compared with the nominal MC from the PYTHIA CP1 tune and MC predictions from the PYTHIA A14, CP5, A3 tunes, the EPOS-LHC generator, and the HERWIG CH3 tune. The statistical uncertainty on each of the predictions is shown by its shaded band.

## 8 Summary

A measurement of event shapes in a minimum-bias selection of proton-proton collisions at a center-of-mass energy of 13 TeV has been presented. Low-pileup data collected with the CMS

detector in 2018 were used, and the kinematics of reconstructed tracks were used to unfold the distributions to the level of stable charged particles. An unbinned multidimensional unfolding algorithm was used to obtain these results. All one- and two-dimensional distributions as well as their covariances are provided.

The results show a consistent trend of mismodeling event shapes common across all simulation configurations considered. The simulations under study include several different underlying physics models, from the PYTHIA, EPOS-LHC, and HERWIG generators, as well as a number of tunes of the PYTHIA generator. For each of these simulations, the isotropy of the event shapes is underpredicted across a range of six different event-shape observables.

Some mismodeling of the charged-particle multiplicity and the invariant mass of the charged particles is also observed. The interplay between these two trends was investigated by considering the event-shape observables in slices of charged-particle multiplicity. These investigations show that the observed data continue to prefer more isotropic distributions even within slices of charged-particle multiplicity and when considering only the plane transverse to the beam direction. These observations suggest that the mismodeling of event shapes is likely not coming only from the distribution of the number of charged particles or poorly modeled longitudinal components.

These event-shape observables are important in probing soft and nonperturbative effects in quantum chromodynamics (QCD) at the LHC. The unfolded data [56, 57] should be used by the community to improve and develop existing proton-proton collision models. The continued development of these models will be critical for understanding phenomena such as quark-gluon plasma and topological effects in non-Abelian gauge theories, such as QCD instantons.

## Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centres and personnel of the Worldwide LHC Computing Grid and other centres for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710,

824093, 101115353, 101002207, 101001205, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Beijing Municipal Science & Technology Commission, No. Z191100007219010, the Fundamental Research Funds for the Central Universities, the Ministry of Science and Technology of China under Grant No. 2023YFA1605804, and the Natural Science Foundation of China under Grant No. 12061141002 (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2021-4.1.2-NEMZ.KI-2024-00036 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Centre for High Performance Computing, Big Data and Quantum Computing and FAIR – Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MICIU/AEI/10.13039/501100011033, ERDF/EU, “European Union NextGenerationEU/PRTR”, and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G670016 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

## References

- [1] ALICE Collaboration, “Long- and short-range correlations and their event-scale dependence in high-multiplicity pp collisions at  $\sqrt{s} = 13$  TeV”, *JHEP* **05** (2021) 290, doi:10.1007/jhep05(2021)290, arXiv:2101.03110.
- [2] CMS Collaboration, “Observation of long-range, near-side angular correlations in proton-proton collisions at the LHC”, *JHEP* **09** (2010) 091, doi:10.1007/jhep09(2010)091, arXiv:1009.4122.
- [3] CMS Collaboration, “Evidence for collectivity in pp collisions at the LHC”, *Phys. Lett. B* **765** (2017) 193, doi:10.1016/j.physletb.2016.12.009, arXiv:1606.06198.
- [4] CMS Collaboration, “Multiplicity and rapidity dependence of strange hadron production in pp, pPb, and PbPb collisions at the LHC”, *Phys. Lett. B* **768** (2017) 103, doi:10.1016/j.physletb.2017.01.075, arXiv:1605.06699.
- [5] ALICE Collaboration, “Enhanced production of multi-strange hadrons in high-multiplicity proton-proton collisions”, *Nat. Phys.* **13** (2017) 535, doi:10.1038/nphys4111, arXiv:1606.07424.



- [6] ALICE Collaboration, “Studying strangeness and baryon production mechanisms through angular correlations between charged  $\Xi$  baryons and identified hadrons in pp collisions at  $\sqrt{s} = 13$  TeV”, *JHEP* **09** (2024) 102, doi:10.1007/JHEP09(2024)102, arXiv:2308.16706.
- [7] T. Pierog et al., “EPOS LHC: Test of collective hadronization with data measured at the CERN Large Hadron Collider”, *Phys. Rev. C* **92** (2015) 034906, doi:10.1103/PhysRevC.92.034906, arXiv:1306.0121.
- [8] P. Bartalini and J. R. Gaunt, eds., “Multiple parton interactions at the LHC”, volume 29. WSP, 2019. Chapter 19, ‘Multiple Scattering in EPOS’. doi:10.1142/10646, ISBN 978-981-322-775-0, 978-981-322-777-4.
- [9] K. Werner, “Core-corona procedure and microcanonical hadronization to understand strangeness enhancement in proton-proton and heavy ion collisions in the EPOS4 framework”, *Phys. Rev. C* **109** (2024) 014910, doi:10.1103/PhysRevC.109.014910, arXiv:2306.10277.
- [10] K. Werner, “Parallel scattering, saturation, and generalized Abramovskii-Gribov-Kancheli (AGK) theorem in the EPOS4 framework, with applications for heavy-ion collisions at  $\sqrt{s_{NN}}$  of 5.02 TeV and 200 GeV”, *Phys. Rev. C* **109** (2024) 034918, doi:10.1103/PhysRevC.109.034918, arXiv:2310.09380.
- [11] K. Werner, “Revealing a deep connection between factorization and saturation: New insight into modeling high-energy proton-proton and nucleus-nucleus scattering in the EPOS4 framework”, *Phys. Rev. C* **108** (2023) 064903, doi:10.1103/PhysRevC.108.064903, arXiv:2301.12517.
- [12] K. Werner and B. Guiot, “Perturbative QCD concerning light and heavy flavor in the EPOS4 framework”, *Phys. Rev. C* **108** (2023) 034904, doi:10.1103/PhysRevC.108.034904, arXiv:2306.02396.
- [13] C. Bierlich, G. Gustafson, L. Lönnblad, and A. Tarasov, “Effects of overlapping strings in pp collisions”, *JHEP* **03** (2015) 148, doi:10.1007/JHEP03(2015)148, arXiv:1412.6259.
- [14] C. Bierlich, “Hadronisation models and colour reconnection”, *PoS DIS2016* (2016) 051, doi:10.22323/1.265.0051, arXiv:1606.09456.
- [15] C. Bierlich, G. Gustafson, and L. Lönnblad, “A shoving model for collectivity in hadronic collisions”, 2016. arXiv:1612.05132.
- [16] Muon g-2 Collaboration, “Measurement of the positive muon anomalous magnetic moment to 0.20 ppm”, *Phys. Rev. Lett.* **131** (2023) 161802, doi:10.1103/PhysRevLett.131.161802, arXiv:2308.06230.
- [17] S. Borsanyi et al., “Leading hadronic contribution to the muon magnetic moment from lattice QCD”, *Nature* **593** (2021) 51, doi:10.1038/s41586-021-03418-1, arXiv:2002.12347.
- [18] A. M. Polyakov, “Compact gauge fields and the infrared catastrophe”, *Phys. Lett. B* **59** (1975) 82, doi:10.1016/0370-2693(75)90162-8.

- 
- [19] A. A. Belavin, A. M. Polyakov, A. S. Schwartz, and Y. S. Tyupkin, "Pseudoparticle solutions of the Yang-Mills equations", *Phys. Lett. B* **59** (1975) 85, doi:10.1016/0370-2693(75)90163-X.
- [20] G. 't Hooft, "Computation of the quantum effects due to a four-dimensional pseudoparticle", *Phys. Rev. D* **14** (1976) 3432, doi:10.1103/PhysRevD.14.3432. See also [21] for erratum.
- [21] G. t. Hooft, "Erratum: Computation of the quantum effects due to a four-dimensional pseudoparticle", *Phys. Rev. D* **18** (Sep, 1978) 2199, doi:10.1103/PhysRevD.18.2199.3.
- [22] R. Jackiw and C. Rebbi, "Vacuum periodicity in a Yang-Mills quantum theory", *Phys. Rev. Lett.* **37** (1976) 172, doi:10.1103/PhysRevLett.37.172.
- [23] C. G. Callan, R. F. Dashen, and D. J. Gross, "The structure of the gauge theory vacuum", *Phys. Lett. B* **63** (1976) 334, doi:10.1016/0370-2693(76)90277-X.
- [24] E. Witten, "Current algebra theorems for the U(1) 'Goldstone boson'", *Nucl. Phys. B* **156** (1979) 269, doi:10.1016/0550-3213(79)90031-2.
- [25] S. Coleman, "The uses of instantons", p. 805. Springer US, Boston, MA, 1979. doi:10.1007/978-1-4684-0991-8\_16.
- [26] E. Shuryak, "Lectures on nonperturbative QCD (nonperturbative topological phenomena in QCD and related theories)", 2020. arXiv:1812.01509.
- [27] V. V. Khoze, F. Krauss, and M. Schott, "Large effects from small QCD instantons: making soft bombs at hadron colliders", *JHEP* **04** (2020) 201, doi:10.1007/jhep04(2020)201, arXiv:1911.09726.
- [28] V. A. Khoze, V. V. Khoze, D. Milne, and M. Ryskin, "Hunting for QCD instantons at the LHC in events with large rapidity gaps", *Phys. Rev. D* **104** (2021) 054013, doi:10.1103/physrevd.104.054013, arXiv:2104.01861.
- [29] V. A. Khoze, V. V. Khoze, D. Milne, and M. Ryskin, "Central instanton production", *Phys. Rev. D* **105** (2022) 036008, doi:10.1103/physrevd.105.036008, arXiv:2111.02159.
- [30] M. Tasevsky, V. A. Khoze, D. Milne, and M. Ryskin, "Searches for QCD instantons with forward proton tagging", *Eur. Phys. J. C* **83** (2023) 35, doi:10.1140/epjc/s10052-023-11178-7, arXiv:2208.14089.
- [31] S. Amoroso, D. Kar, and M. Schott, "How to discover QCD instantons at the LHC", *Eur. Phys. J. C* **81** (2021) 624, doi:10.1140/epjc/s10052-021-09412-1, arXiv:2012.09120.
- [32] E. Shuryak and I. Zahed, "How to observe the QCD instanton/sphaleron processes at hadron colliders?", 2021. arXiv:2102.00256.
- [33] CMS Collaboration, "Precision luminosity measurement in proton-proton collisions at  $\sqrt{s} = 13$  TeV in 2015 and 2016 at CMS", *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.

- [34] CMS Collaboration, “CMS luminosity measurement for the 2018 data-taking period at  $\sqrt{s} = 13$  TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-18-002, 2019.
- [35] ALEPH Collaboration, “Studies of QCD at e+ e- centre-of-mass energies between 91-GeV and 209-GeV”, *Eur. Phys. J. C* **35** (2004) 457, doi:10.1140/epjc/s2004-01891-4.
- [36] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA 8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) 4, doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.
- [37] CMS Collaboration, “Measurement of charged particle spectra in minimum-bias events from proton-proton collisions at  $\sqrt{s} = 13$  TeV”, *Eur. Phys. J. C* **78** (2018) 697, doi:10.1140/epjc/s10052-018-6144-y, arXiv:1806.11245.
- [38] ALICE Collaboration, “Multiplicity dependence of charged-particle production in pp, p-Pb, Xe-Xe and Pb-Pb collisions at the LHC”, *Phys. Lett. B* **845** (2023) 138110, doi:10.1016/j.physletb.2023.138110, arXiv:2211.15326.
- [39] ATLAS Collaboration, “Charged-particle distributions at low transverse momentum in  $\sqrt{s} = 13$  TeV  $pp$  interactions measured with the ATLAS detector at the LHC”, *Eur. Phys. J. C* **76** (2016) 502, doi:10.1140/epjc/s10052-016-4335-y, arXiv:1606.01133.
- [40] CMS Collaboration, “Study of the underlying event in top quark pair production in pp collisions at 13 TeV”, *Eur. Phys. J. C* **79** (2019) doi:10.1140/epjc/s10052-019-6620-z, arXiv:1807.02810.
- [41] ATLAS Collaboration, “Measurements of multijet event isotropies using optimal transport with the ATLAS detector”, *JHEP* **10** (2023) 060, doi:10.1007/JHEP10(2023)060, arXiv:2305.16930.
- [42] CMS Collaboration, “First measurement of hadronic event shapes in pp collisions at  $\sqrt{s} = 7$  TeV”, *Phys. Lett. B* **699** (2011) 48, doi:10.1016/j.physletb.2011.03.060, arXiv:1102.0068.
- [43] ALICE Collaboration, “Transverse sphericity of primary charged particles in minimum bias proton-proton collisions at  $\sqrt{s} = 0.9, 2.76$  and 7 TeV”, *Eur. Phys. J. C* **72** (2012) 2124, doi:10.1140/epjc/s10052-012-2124-9, arXiv:1205.3963.
- [44] ATLAS Collaboration, “Measurement of charged-particle event shape variables in  $\sqrt{s} = 7$  TeV proton-proton interactions with the ATLAS detector”, *Phys. Rev. D* **88** (2013) 032004, doi:10.1103/PhysRevD.88.032004, arXiv:1207.6915.
- [45] G. Parisi, “Superinclusive cross sections”, *Phys. Lett. B* **74** (1978) 65, doi:10.1016/0370-2693(78)90061-8.
- [46] J. F. Donoghue, F. E. Low, and S.-Y. Pi, “Tensor analysis of hadronic jets in quantum chromodynamics”, *Phys. Rev. D* **20** (1979) 2759, doi:10.1103/PhysRevD.20.2759.
- [47] S. Brandt and H. Dahmen, “Axes and scalar measures of two-jet and three-jet events”, *Z. Phys. C* **1** (1979) 61, doi:10.1007/BF01450381.
- [48] P. T. Komiske, E. M. Metodiev, and J. Thaler, “Energy flow networks: deep sets for particle jets”, *JHEP* **01** (2019) 121, doi:10.1007/jhep01(2019)121, arXiv:1810.05165.

- 
- [49] C. Cesarotti and J. Thaler, “A robust measure of event isotropy at colliders”, *JHEP* **08** (2020) 084, doi:10.1007/JHEP08(2020)084, arXiv:2004.06125.
- [50] C. Cesarotti, M. Reece, and M. J. Strassler, “The efficacy of event isotropy as an event shape observable”, *JHEP* **07** (2021) 215, doi:10.1007/jhep07(2021)215, arXiv:2011.06599.
- [51] M. Bähr et al., “HERWIG++ physics and manual”, *Eur. Phys. J. C* **58** (2008) 639, doi:10.1140/epjc/s10052-008-0798-9, arXiv:0803.0883.
- [52] J. Bellm et al., “HERWIG 7.0/HERWIG++ 3.0 release note”, *Eur. Phys. J. C* **76** (2016) 196, doi:10.1140/epjc/s10052-016-4018-8, arXiv:1512.01178.
- [53] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [54] A. Andreassen et al., “Omnifold: A method to simultaneously unfold all observables”, *Phys. Rev. Lett.* **124** (2020) 182001, doi:10.1103/PhysRevLett.124.182001, arXiv:1911.09107.
- [55] A. Andreassen et al., “Scaffolding simulations with deep learning for high-dimensional deconvolution”, 2021. arXiv:2105.04448.
- [56] “HEPData record for this analysis”, 2025. doi:10.17182/hepdata.157862.
- [57] “CERN Document Server repository for this analysis”, 2025. doi:10.17181/d5j30-5wr08.
- [58] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [59] CMS Collaboration, “Development of the CMS detector for the CERN LHC Run 3”, *JINST* **19** (2024) P05064, doi:10.1088/1748-0221/19/05/P05064, arXiv:2309.05466.
- [60] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014) P10009, doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569.
- [61] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965.
- [62] CMS Collaboration, “Pileup mitigation at CMS in 13 TeV data”, *JINST* **15** (2020), no. 09, doi:10.1088/1748-0221/15/09/P09018, arXiv:2003.00503.
- [63] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [64] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [65] ATLAS Collaboration, “The PYTHIA 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie-Landshoff diffractive model”, ATLAS Note ATL-PHYS-PUB-2016-017, 2016.

- [66] CMS Collaboration, “Development and validation of HERWIG 7 tunes from CMS underlying-event measurements”, *Eur. Phys. J. C* **81** (2021) 312, doi:10.1140/epjc/s10052-021-08949-5, arXiv:2011.03422.
- [67] A. Buckley et al., “The HepMC3 event record library for Monte Carlo event generators”, *Comput. Phys. Comm.* **260** (2021) 107310, doi:10.1016/j.cpc.2020.107310, arXiv:1912.08005.
- [68] W. H. Richardson, “Bayesian-based iterative method of image restoration”, *J. Opt. Soc. Am.* **62** (1972) 55, doi:10.1364/JOSA.62.000055.
- [69] L. B. Lucy, “An iterative technique for the rectification of observed distributions”, *Astronomical Journal* **79** (1974) 745, doi:10.1086/111605.
- [70] G. D’Agostini, “A multidimensional unfolding method based on Bayes’ theorem”, *Nucl. Instrum. Meth. A* **362** (1995) 487, doi:10.1016/0168-9002(95)00274-X.
- [71] K. Cranmer, J. Pavez, and G. Louppe, “Approximating likelihood ratios with calibrated discriminative classifiers”, 6, 2015. arXiv:1506.02169.
- [72] CMS Collaboration, “Measurement of the inelastic proton-proton cross section at  $\sqrt{s} = 13$  TeV”, *JHEP* **07** (2018) 161, doi:10.1007/JHEP07(2018)161, arXiv:1802.02613.
- [73] R. D. Cousins, S. J. May, and Y. Sun, “Should unfolded histograms be used to test hypotheses?”, 2016. arXiv:1607.07038.
- [74] S. M. S. Lee and G. A. Young, “Parametric bootstrapping with nuisance parameters”, *Stat. Probab. Lett.* **71** (2005) 143, doi:10.1016/j.spl.2004.10.026.
- [75] R. J. Tibshirani and B. Efron, “An introduction to the bootstrap”. Chapman and Hall New York, 1993. Monographs on statistics and applied probability 57. doi:10.1201/9780429246593.
- [76] CMS Collaboration, “Tracking performance for charged pions with Run2 Legacy data”, CMS Detector Performance Note CMS-DP-2022-012, 2022.
- [77] CMS Collaboration, “Muon tracking efficiency for 2018 dataset using tag and probe method”, CMS Detector Performance Note CMS-DP-2020-013, 2019.
- [78] CMS Collaboration, “Event generator tunes obtained from underlying event and multiparton scattering measurements”, *Eur. Phys. J. C* **76** (2016) 155, doi:10.1140/epjc/s10052-016-3988-x, arXiv:1512.00815.
- [79] CMS Collaboration, “Investigations of the impact of the parton shower tuning in PYTHIA 8 in the modelling of  $t\bar{t}$  at  $\sqrt{s} = 8$  and 13 TeV”, CMS Physics Analysis Summary CMS-PAS-TOP-16-021, 2016.
- [80] CMS Collaboration, “Measurement of the underlying event activity at the LHC with  $\sqrt{s} = 7$  TeV and comparison with  $\sqrt{s} = 0.9$  TeV”, *JHEP* **09** (2011) 109, doi:10.1007/JHEP09(2011)109, arXiv:1107.0330.
- [81] CMS Collaboration, “CMS PYTHIA 8 colour reconnection tunes based on underlying-event data”, *Eur. Phys. J. C* **83** (2023) 587, doi:10.1140/epjc/s10052-023-11630-8, arXiv:2205.02905.
- [82] ATLAS Collaboration, “ATLAS PYTHIA 8 tunes to 7 TeV data”, ATLAS Note ATL-PHYS-PUB-2014-021, 2014.

## A Additional simulated samples for validation

Table A.1 gives additional simulated samples that are used as pseudodata to validate the performance of the unfolding algorithm or to extensively investigate the impacts of various models on the observables to be measured.

Table A.1: Samples of simulated minimum-bias events used for validation. The value of  $\alpha_S$  corresponds to that used in the PDF set.

|                    | Generator                                             | PDF ( $\alpha_S(m_Z)$ )          |
|--------------------|-------------------------------------------------------|----------------------------------|
| Pseudodata samples | PYTHIA8 (CP5) [36]                                    | NNPDF3.1 QCD+LUXQED NNLO (0.118) |
|                    | PYTHIA8 (CUETP8M1) [78]                               | NNPDF2.3 QCD+QED LO (0.130)      |
| Other validation   | PYTHIA8 (CP5) with pileup                             | NNPDF3.1 QCD+LUXQED NNLO (0.118) |
|                    | PYTHIA8 (CP5) $\alpha_S^{\text{FSR}}$ variations [36] | NNPDF3.1 QCD+LUXQED NNLO (0.118) |
|                    | PYTHIA8 CUETP8M2T4 [40, 79]                           | NNPDF3.0 QCD LO (0.130)          |
|                    | PYTHIA8 (CP5) color reconnection tunes [81]           | NNPDF3.1 QCD+LUXQED NNLO (0.118) |
|                    | PYTHIA8 (CP2)                                         | NNPDF3.1 QCD LO (0.130)          |
|                    | PYTHIA8 (A14) [82]                                    | NNPDF3.1 QCD LO (0.130)          |
|                    | PYTHIA8 (A14) eigenvariations                         | NNPDF3.1 QCD LO (0.130)          |
|                    | PYTHIA8 (A14) CTEQL1                                  | CTEQL1 (0.1298)                  |
|                    | PYTHIA8 (A14) MSTW2008LO                              | MSTW2008LO (0.13939)             |
|                    | PYTHIA8 (A14) HERAPDF1.5LO                            | HERAPDF1.5LO (0.130)             |

## B Additional two-dimensional distributions

Figures B.1, B.2 and B.3 show the unfolded distributions of isotropy, broadening, thrust, transverse thrust and invariant mass of charged particles in slices of the charged-particle multiplicity, as well as the MC predictions for comparison.




Figure B.1: The unfolded distributions of event isotropy (upper) and broadening (lower) in slices of charged-particle multiplicity compared with the nominal MC from the PYTHIA CP1 tune and MC predictions from the PYTHIA A14, CP5, A3 tunes, the EPOS-LHC generator, and the HERWIG CH3 tune. The statistical uncertainty on each of the predictions is shown by its shaded band.




Figure B.2: The unfolded distributions of thrust (upper) and transverse thrust (lower) in slices of charged-particle multiplicity compared with the nominal MC from the PYTHIA CP1 tune and MC predictions from the PYTHIA A14, CP5, A3 tunes, the EPOS-LHC generator, and the HERWIG CH3 tune. The statistical uncertainty on each of the predictions is shown by its shaded band.






Figure B.3: The unfolded invariant mass of charged particles in slices of charged-particle multiplicity compared with the nominal MC from the PYTHIA CP1 tune and MC predictions from the PYTHIA A14, CP5, A3 tunes, the EPOS-LHC generator, and the HERWIG CH3 tune. The statistical uncertainty on each of the predictions is shown by its shaded band.

## C Correlation matrices

Figures C.1 and C.2 provide the correlation matrices for the systematic uncertainties in the histograms given in Figs. 1, 2, and 3.




Figure C.1: The correlations of the unfolding systematic uncertainty between bins of the event-shape observables shown for (upper left) sphericity, (upper right) thrust, (lower left) broadening, and (lower right) transverse sphericity.




Figure C.2: The correlations of the unfolding systematic uncertainty between bins of the event-shape observables shown for (upper left) transverse thrust, (upper right) isotropy, (lower left) particle multiplicity, and (lower right) invariant mass.

## D Uncertainty decomposition

Table D.1 shows the uncertainty decomposition of unfolding, quantified by ratios of the uncertainties from individual sources to the total uncertainties, averaged over bins of transverse sphericity.

Table D.1: Relative uncertainties of unfolding from different sources, averaged over bins of transverse sphericity. The uncertainty from regularization and that from the model deficiency of the migration are estimated from the square root of the quadrature sum of the differences between the unfolding result using the nominal MC simulation and those using reweighted MC samples to systematic variations. The MC statistical uncertainty is estimated from the statistical uncertainty of the histogram of weighted MC samples after the unfolding. The data statistical uncertainty is estimated from the standard deviations of the unfolding results to alternative data samples with weight following a Poisson(1) distribution. The average relative uncertainties for the distributions of other observables are similar in size.

| Uncertainty source                | Relative uncertainty |
|-----------------------------------|----------------------|
| Track reconstruction efficiency   | 1.7%                 |
| Model deficiency of the migration | 1.6%                 |
| Bias from regularization          | 1.2%                 |
| MC event count                    | 0.26%                |
| Data event count                  | 0.26%                |

## E The CMS Collaboration

### Yerevan Physics Institute, Yerevan, Armenia

V. Chekhovsky, A. Hayrapetyan, V. Makarenko , A. Tumasyan<sup>1</sup> 

### Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, L. Benato , T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic , P.S. Hussain , M. Jeitler<sup>2</sup> , N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck<sup>2</sup> , R. Schöfbeck<sup>2</sup> , D. Schwarz , M. Sonawane , W. Waltenberger , C.-E. Wulz<sup>2</sup> 

### Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , H. Kwon , T. Van Laer, P. Van Mechelen 

### Vrije Universiteit Brussel, Brussel, Belgium

N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, Y. Hong , S. Lowette , I. Makarenko , D. Müller , S. Tavernier , M. Tytgat<sup>3</sup> , G.P. Van Onsem , S. Van Putte , D. Vannerom 

### Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios , A. Khalilzadeh, F.A. Khan , A. Malara , M.A. Shahzad, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 

### Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , J. Knolle , L. Lambrecht , D. Marckx , K. Skovpen , N. Van Den Bossche , J. van der Linden , J. Vandenbroeck , L. Wezenbeek 

### Université Catholique de Louvain, Louvain-la-Neuve, Belgium

S. Bein , A. Benecke , A. Bethani , G. Bruno , C. Caputo , J. De Favereau De Jeneret , C. Delaere , I.S. Donertas , A. Giammanco , A.O. Guzel , Sa. Jain , V. Lemaitre, J. Lidrych , P. Mastrapasqua , T.T. Tran , S. Turkcapar 

### Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , G. Correia Silva , C. Hensel , T. Menezes De Oliveira , C. Mora Herrera<sup>4</sup> , P. Rebello Teles , M. Soeiro, E.J. Tonelli Manganote<sup>5</sup> , A. Vilela Pereira<sup>4</sup> 

### Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato<sup>6</sup>, E.M. Da Costa , G.G. Da Silveira<sup>7</sup> , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza, T. Laux Kuhn<sup>7</sup> , M. Macedo , J. Martins , K. Mota Amarilo , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel 

### Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes<sup>7</sup> , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula , V. Scheurer

### Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 

**University of Sofia, Sofia, Bulgaria**

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 

**Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile**

S. Keshri , D. Laroze , S. Thakur 

**Beihang University, Beijing, China**

T. Cheng , T. Javaid , L. Yuan 

**Department of Physics, Tsinghua University, Beijing, China**

Z. Hu , Z. Liang, J. Liu

**Institute of High Energy Physics, Beijing, China**

G.M. Chen<sup>8</sup> , H.S. Chen<sup>8</sup> , M. Chen<sup>8</sup> , F. Iemmi , C.H. Jiang, A. Kapoor<sup>9</sup> , H. Liao , Z.-A. Liu<sup>10</sup> , R. Sharma<sup>11</sup> , J.N. Song<sup>10</sup>, J. Tao , C. Wang<sup>8</sup>, J. Wang , Z. Wang<sup>8</sup>, H. Zhang , J. Zhao 

**State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China**

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, X. Sun , D. Wang , H. Yang, Y. Zhao, C. Zhou 

**Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China**

S. Yang 

**Sun Yat-Sen University, Guangzhou, China**

Z. You 

**University of Science and Technology of China, Hefei, China**

K. Jaffel , N. Lu 

**Nanjing Normal University, Nanjing, China**

G. Bauer<sup>12</sup>, B. Li<sup>13</sup>, H. Wang , K. Yi<sup>14</sup> , J. Zhang 

**Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China**

Y. Li

**Zhejiang University, Hangzhou, Zhejiang, China**

Z. Lin , C. Lu , M. Xiao 

**Universidad de Los Andes, Bogota, Colombia**

C. Avila , D.A. Barbosa Trujillo , A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

**Universidad de Antioquia, Medellin, Colombia**

J. Jaramillo , C. Rendón , M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

**University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia**

D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac 

**University of Split, Faculty of Science, Split, Croatia**

M. Kovac , A. Petkovic , T. Sculac 

**Institute Rudjer Boskovic, Zagreb, Croatia**

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov<sup>15</sup> ,

T. Susa 

**University of Cyprus, Nicosia, Cyprus**

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , J. Mousa , C. Nicolaou, L. Paizanos, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

**Charles University, Prague, Czech Republic**

M. Finger , M. Finger Jr. , A. Kveton 

**Escuela Politecnica Nacional, Quito, Ecuador**

E. Ayala 

**Universidad San Francisco de Quito, Quito, Ecuador**

E. Carrera Jarrin 

**Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**

B. El-mahdy , S. Khalil<sup>16</sup> , E. Salama<sup>17,18</sup> 

**Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt**

M. Abdullah Al-Mashad , M.A. Mahmoud 

**National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**

K. Ehataht , M. Kadastik, T. Lange , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken 

**Department of Physics, University of Helsinki, Helsinki, Finland**

K. Osterberg , M. Voutilainen 

**Helsinki Institute of Physics, Helsinki, Finland**

N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew , K.T.S. Kallonen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Myllymäki , M.m. Rantanen , J. Tuominiemi 

**Lappeenranta-Lahti University of Technology, Lappeenranta, Finland**

H. Kirschenmann , P. Luukka , H. Petrow 

**IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France**

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi , L. Portales , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro<sup>19</sup> , P. Simkina , M. Titov , M. Tornago 

**Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France**

F. Beaudette , G. Boldrini , P. Busson , A. Cappati , C. Charlot , M. Chiusi , T.D. Cuisset , F. Damas , O. Davignon , A. De Wit , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , B. Harikrishnan , L. Kalipoliti , G. Liu , M. Manoni , M. Nguyen , S. Obraztsov , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , G. Sokmen, L. Urda Gómez , E. Vernazza , A. Zabi , A. Zghiche 

**Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France**

J.-L. Agram<sup>20</sup> , J. Andrea , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet , G. Saha , M.A. Sessini , P. Van Hove , P. Vaucelle 

**Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France**

A. Di Florio 

**Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France**

D. Amram , S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo , P. Depasse , C. Dozen<sup>21</sup> , H. El Mamouni , J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg , G. Grenier , B. Ille , E. Jourdhuy , I.B. Laktineh , M. Lethuillier , L. Mirabito , S. Perries , A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao 

**Georgian Technical University, Tbilisi, Georgia**

I. Lomidze , T. Toriashvili<sup>22</sup> , Z. Tsamalaidze<sup>15</sup> 

**RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser , A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

**RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany**

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann , M. Erdmann , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , N. Kumar , M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , F. Nowotny , A. Pozdnyakov , Y. Rath , W. Redjeb , F. Rehm , H. Reithler , V. Sarkisovi , A. Schmidt , C. Seth , A. Sharma , J.L. Spah , F. Torres Da Silva De Araujo<sup>23</sup> , S. Wiedenbeck , S. Zaleski 

**RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany**

C. Dziwok , G. Flügge , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 

**Deutsches Elektronen-Synchrotron, Hamburg, Germany**

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso , Y. An , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman<sup>24</sup> , K. Borras<sup>25</sup> , A. Campbell , A. Cardini , F. Colombina , M. De Silva , G. Eckerlin , D. Eckstein , L.I. Estevez Banos , E. Gallo<sup>24</sup> , A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , B. Kaech , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange , D. Leyva Pernia , K. Lipka<sup>26</sup> , W. Lohmann<sup>27</sup> , F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , J. Park , E. Ranken , A. Raspereza , D. Rastorguev , J. Rübenach , L. Rygaard , M. Scham<sup>28,25</sup> , S. Schnake<sup>25</sup> , P. Schütze , C. Schwanenberger<sup>24</sup> , D. Selivanova , K. Sharke , M. Shchedrolosiev , D. Stafford , F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , K. Wichmann , L. Wiens<sup>25</sup> , C. Wissing , Y. Yang , S. Zakharov , A. Zimmermann Castro Santos 

**University of Hamburg, Hamburg, Germany**

A. Albrecht , S. Albrecht , M. Antonello , S. Bollweg , M. Bonanomi , P. Connor , K. El Morabit , Y. Fischer , E. Garutti , A. Grohsjean , J. Haller , D. Hundhausen , H.R. Jabusch , G. Kasieczka , P. Keicher , R. Klanner , W. Korcari , T. Kramer , C.c. Kuo , V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , L. Moureaux , M. Mrowietz , A. Nigamova , Y. Nissan , A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoiu , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews , B. Wiederspan , M. Wolf 



**Karlsruher Institut fuer Technologie, Karlsruhe, Germany**

S. Brommer , E. Butz , Y.M. Chen , T. Chwalek , A. Dierlamm , G.G. Dincer , U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann<sup>29</sup> , R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , O. Lavoryk , J.M. Lawhorn , M. Link, A. Lintuluoto , S. Maier , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , R. Schmieder, N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa, L. Stockmeier, K. Tauqeer, M. Toms , B. Topko , N. Trevisani , T. Voigtländer , R.F. Von Cube , J. Von Den Driesch, M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo 

**Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece**

G. Anagnostou, G. Daskalakis , A. Kyriakis , A. Papadopoulos<sup>29</sup>, A. Stakia 

**National and Kapodistrian University of Athens, Athens, Greece**

G. Melachroinos, Z. Painesis , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 

**National Technical University of Athens, Athens, Greece**

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsipolitis 

**University of Ioánnina, Ioánnina, Greece**

I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

**HUN-REN Wigner Research Centre for Physics, Budapest, Hungary**

C. Hajdu , D. Horvath<sup>30,31</sup> , K. Márton, A.J. Rádl<sup>32</sup> , F. Sikler , V. Veszpremi 

**MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary**

M. Csanád , K. Farkas , A. Fehérkuti<sup>33</sup> , M.M.A. Gadallah<sup>34</sup> , Á. Kadlecik , P. Major , G. Pásztor , G.I. Veres 

**Faculty of Informatics, University of Debrecen, Debrecen, Hungary**

B. Ujvari , G. Zilizi 

**HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary**

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

**Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary**

T. Csorgo<sup>33</sup> , F. Nemes<sup>33</sup> , T. Novak 

**Panjab University, Chandigarh, India**

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra<sup>35</sup> , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand, J.B. Singh , A. Singla 

**University of Delhi, Delhi, India**

A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya 

**Indian Institute of Technology Kanpur, Kanpur, India**

S. Mukherjee 

**Saha Institute of Nuclear Physics, HBNI, Kolkata, India**

S. Baradia , S. Barman<sup>36</sup> , S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

### **Indian Institute of Technology Madras, Madras, India**

M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , G. Dash , P. Jana , P. Kalbhor , S. Kamble , J.R. Komaragiri<sup>37</sup> , D. Kumar<sup>37</sup> , T. Mishra , B. Parida<sup>38</sup> , P.R. Pujahari , N.R. Saha , A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay 

### **Tata Institute of Fundamental Research-A, Mumbai, India**

S. Dugad, G.B. Mohanty , M. Shelake, P. Suryadevara

### **Tata Institute of Fundamental Research-B, Mumbai, India**

A. Bala , S. Banerjee , S. Bhowmik<sup>39</sup> , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, B.M. Joshi , S. Kumar , G. Majumder , K. Mazumdar , S. Parolia , A. Thachayath 

### **National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India**

S. Bahinipati<sup>40</sup> , C. Kar , D. Maity<sup>41</sup> , P. Mal , K. Naskar<sup>41</sup> , A. Nayak<sup>41</sup> , S. Nayak, K. Pal , R. Raturi, P. Sadangi, S.K. Swain , S. Varghese<sup>41</sup> , D. Vats<sup>41</sup> 

### **Indian Institute of Science Education and Research (IISER), Pune, India**

S. Acharya<sup>42</sup> , A. Alpana , S. Dube , B. Gomber<sup>42</sup> , P. Hazarika , B. Kansal , A. Laha , B. Sahu<sup>42</sup> , S. Sharma , K.Y. Vaish 

### **Isfahan University of Technology, Isfahan, Iran**

H. Bakhshiansohi<sup>43</sup> , A. Jafari<sup>44</sup> , M. Zeinali<sup>45</sup> 

### **Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**

S. Bashiri, S. Chenarani<sup>46</sup> , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie , M. Mohammadi Najafabadi , S. Tizchang<sup>47</sup> 

### **University College Dublin, Dublin, Ireland**

M. Felcini , M. Grunewald 

### **INFN Sezione di Bari<sup>a</sup>, Università di Bari<sup>b</sup>, Politecnico di Bari<sup>c</sup>, Bari, Italy**

M. Abbrescia<sup>a,b</sup> , A. Colaleo<sup>a,b</sup> , D. Creanza<sup>a,c</sup> , B. D'Anzi<sup>a,b</sup> , N. De Filippis<sup>a,c</sup> , M. De Palma<sup>a,b</sup> , W. Elmetenawee<sup>a,b,48</sup> , N. Ferrara<sup>a,b</sup> , L. Fiore<sup>a</sup> , G. Iaselli<sup>a,c</sup> , L. Longo<sup>a</sup> , M. Louka<sup>a,b</sup> , G. Maggi<sup>a,c</sup> , M. Maggi<sup>a</sup> , I. Margjeka<sup>a</sup> , V. Mastrapasqua<sup>a,b</sup> , S. My<sup>a,b</sup> , S. Nuzzo<sup>a,b</sup> , A. Pellecchia<sup>a,b</sup> , A. Pompili<sup>a,b</sup> , G. Pugliese<sup>a,c</sup> , R. Radogna<sup>a,b</sup> , D. Ramos<sup>a</sup> , A. Ranieri<sup>a</sup> , L. Silvestris<sup>a</sup> , F.M. Simone<sup>a,c</sup> , Ü. Sözbilir<sup>a</sup> , A. Stamerra<sup>a,b</sup> , D. Troiano<sup>a,b</sup> , R. Venditti<sup>a,b</sup> , P. Verwilligen<sup>a</sup> , A. Zaza<sup>a,b</sup> 

### **INFN Sezione di Bologna<sup>a</sup>, Università di Bologna<sup>b</sup>, Bologna, Italy**

G. Abbiendi<sup>a</sup> , C. Battilana<sup>a,b</sup> , D. Bonacorsi<sup>a,b</sup> , P. Capiluppi<sup>a,b</sup> , A. Castro<sup>†a,b</sup> , F.R. Cavallo<sup>a</sup> , M. Cuffiani<sup>a,b</sup> , G.M. Dallavalle<sup>a</sup> , T. Diotallevi<sup>a,b</sup> , F. Fabbri<sup>a</sup> , A. Fanfani<sup>a,b</sup> , D. Fasanella<sup>a</sup> , P. Giacomelli<sup>a</sup> , L. Giommi<sup>a,b</sup> , C. Grandi<sup>a</sup> , L. Guiducci<sup>a,b</sup> , S. Lo Meo<sup>a,49</sup> , M. Lorusso<sup>a,b</sup> , L. Lunerti<sup>a</sup> , S. Marcellini<sup>a</sup> , G. Masetti<sup>a</sup> , F.L. Navarria<sup>a,b</sup> , G. Paggi<sup>a,b</sup> , A. Perrotta<sup>a</sup> , F. Primavera<sup>a,b</sup> , A.M. Rossi<sup>a,b</sup> , S. Rossi Tisbeni<sup>a,b</sup> , T. Rovelli<sup>a,b</sup> , G.P. Siroli<sup>a,b</sup> 

### **INFN Sezione di Catania<sup>a</sup>, Università di Catania<sup>b</sup>, Catania, Italy**

S. Costa<sup>a,b,50</sup> , A. Di Mattia<sup>a</sup> , A. Lapertosa<sup>a</sup> , R. Potenza<sup>a,b</sup> , A. Tricomi<sup>a,b,50</sup> 

### **INFN Sezione di Firenze<sup>a</sup>, Università di Firenze<sup>b</sup>, Firenze, Italy**

P. Assiouras<sup>a</sup> , G. Barbagli<sup>a</sup> , G. Bardelli<sup>a,b</sup> , M. Bartolini<sup>a,b</sup> , B. Camaiani<sup>a,b</sup> , A. Cassese<sup>a</sup> , R. Ceccarelli<sup>a</sup> , V. Ciulli<sup>a,b</sup> , C. Civinini<sup>a</sup> , R. D'Alessandro<sup>a,b</sup> , L. Damenti<sup>a,b</sup> , E. Focardi<sup>a,b</sup> , T. Kello<sup>a</sup> , G. Latino<sup>a,b</sup> , P. Lenzi<sup>a,b</sup> , M. Lizzo<sup>a</sup> , M. Meschini<sup>a</sup> , S. Paoletti<sup>a</sup> , A. Papanastassiou<sup>a,b</sup> , G. Sguazzoni<sup>a</sup> , L. Viliani<sup>a</sup> 

#### **INFN Laboratori Nazionali di Frascati, Frascati, Italy**

L. Benussi , S. Bianco , S. Meola<sup>51</sup> , D. Piccolo 

#### **INFN Sezione di Genova<sup>a</sup>, Università di Genova<sup>b</sup>, Genova, Italy**

M. Alves Gallo Pereira<sup>a</sup> , F. Ferro<sup>a</sup> , E. Robutti<sup>a</sup> , S. Tosi<sup>a,b</sup> 

#### **INFN Sezione di Milano-Bicocca<sup>a</sup>, Università di Milano-Bicocca<sup>b</sup>, Milano, Italy**

A. Benaglia<sup>a</sup> , F. Brivio<sup>a</sup> , F. Cetorelli<sup>a,b</sup> , F. De Guio<sup>a,b</sup> , M.E. Dinardo<sup>a,b</sup> , P. Dini<sup>a</sup> , S. Gennai<sup>a</sup> , R. Gerosa<sup>a,b</sup> , A. Ghezzi<sup>a,b</sup> , P. Govoni<sup>a,b</sup> , L. Guzzi<sup>a</sup> , G. Lavizzari<sup>a,b</sup> , M.T. Lucchini<sup>a,b</sup> , M. Malberti<sup>a</sup> , S. Malvezzi<sup>a</sup> , A. Massironi<sup>a</sup> , D. Menasce<sup>a</sup> , L. Moroni<sup>a</sup> , M. Paganoni<sup>a,b</sup> , S. Palluotto<sup>a,b</sup> , D. Pedrini<sup>a</sup> , A. Perego<sup>a,b</sup> , B.S. Pinolini<sup>a</sup> , G. Pizzati<sup>a,b</sup> , S. Ragazzi<sup>a,b</sup> , T. Tabarelli de Fatis<sup>a,b</sup> 

#### **INFN Sezione di Napoli<sup>a</sup>, Università di Napoli 'Federico II'<sup>b</sup>, Napoli, Italy; Università della Basilicata<sup>c</sup>, Potenza, Italy; Scuola Superiore Meridionale (SSM)<sup>d</sup>, Napoli, Italy**

S. Buontempo<sup>a</sup> , A. Cagnotta<sup>a,b</sup> , F. Carnevali<sup>a,b</sup> , N. Cavallo<sup>a,c</sup> , F. Fabozzi<sup>a,c</sup> , A.O.M. Iorio<sup>a,b</sup> , L. Lista<sup>a,b,52</sup> , P. Paolucci<sup>a,29</sup> , B. Rossi<sup>a</sup> 

#### **INFN Sezione di Padova<sup>a</sup>, Università di Padova<sup>b</sup>, Padova, Italy; Università degli Studi di Cagliari<sup>c</sup>, Cagliari, Italy**

R. Ardino<sup>a</sup> , P. Azzi<sup>a</sup> , N. Bacchetta<sup>a,53</sup> , A. Bergnoli<sup>a</sup> , D. Bisello<sup>a,b</sup> , P. Bortignon<sup>a</sup> , G. Bortolato<sup>a,b</sup> , A.C.M. Bulla<sup>a</sup> , R. Carlin<sup>a,b</sup> , T. Dorigo<sup>a,54</sup> , F. Gasparini<sup>a,b</sup> , U. Gasparini<sup>a,b</sup> , S. Giorgetti<sup>a</sup> , E. Lusiani<sup>a</sup> , M. Margoni<sup>a,b</sup> , A.T. Meneguzzo<sup>a,b</sup> , J. Pazzini<sup>a,b</sup> , P. Ronchese<sup>a,b</sup> , R. Rossin<sup>a,b</sup> , F. Simonetto<sup>a,b</sup> , M. Tosi<sup>a,b</sup> , A. Triossi<sup>a,b</sup> , S. Ventura<sup>a</sup> , M. Zanetti<sup>a,b</sup> , P. Zotto<sup>a,b</sup> , A. Zucchetta<sup>a,b</sup> , G. Zumerle<sup>a,b</sup> 

#### **INFN Sezione di Pavia<sup>a</sup>, Università di Pavia<sup>b</sup>, Pavia, Italy**

A. Braghieri<sup>a</sup> , S. Calzaferri<sup>a</sup> , D. Fiorina<sup>a</sup> , P. Montagna<sup>a,b</sup> , M. Pelliccioni<sup>a</sup> , V. Re<sup>a</sup> , C. Riccardi<sup>a,b</sup> , P. Salvini<sup>a</sup> , I. Vai<sup>a,b</sup> , P. Vitulo<sup>a,b</sup> 

#### **INFN Sezione di Perugia<sup>a</sup>, Università di Perugia<sup>b</sup>, Perugia, Italy**

S. Ajmal<sup>a,b</sup> , M.E. Ascioti<sup>a,b</sup> , G.M. Bilei<sup>a</sup> , C. Carrivale<sup>a,b</sup> , D. Ciangottini<sup>a,b</sup> , L. Fanò<sup>a,b</sup> , V. Mariani<sup>a,b</sup> , M. Menichelli<sup>a</sup> , F. Moscatelli<sup>a,55</sup> , A. Rossi<sup>a,b</sup> , A. Santocchia<sup>a,b</sup> , D. Spiga<sup>a</sup> , T. Tedeschi<sup>a,b</sup> 

#### **INFN Sezione di Pisa<sup>a</sup>, Università di Pisa<sup>b</sup>, Scuola Normale Superiore di Pisa<sup>c</sup>, Pisa, Italy; Università di Siena<sup>d</sup>, Siena, Italy**

C. Aimè<sup>a</sup> , C.A. Alexe<sup>a,c</sup> , P. Asenov<sup>a,b</sup> , P. Azzurri<sup>a</sup> , G. Bagliesi<sup>a</sup> , R. Bhattacharya<sup>a</sup> , L. Bianchini<sup>a,b</sup> , T. Boccali<sup>a</sup> , E. Bossini<sup>a</sup> , D. Bruschini<sup>a,c</sup> , R. Castaldi<sup>a</sup> , F. Cattafesta<sup>a,c</sup> , M.A. Ciocci<sup>a,b</sup> , M. Cipriani<sup>a,b</sup> , V. D'Amante<sup>a,d</sup> , R. Dell'Orso<sup>a</sup> , S. Donato<sup>a,b</sup> , A. Giassi<sup>a</sup> , F. Ligabue<sup>a,c</sup> , A.C. Marini<sup>a,b</sup> , D. Matos Figueiredo<sup>a</sup> , A. Messineo<sup>a,b</sup> , S. Mishra<sup>a</sup> , V.K. Muraleedharan Nair Bindhu<sup>a,b,41</sup> , M. Musich<sup>a,b</sup> , S. Nandan<sup>a</sup> , F. Palla<sup>a</sup> , A. Rizzi<sup>a,b</sup> , G. Rolandi<sup>a,c</sup> , S. Roy Chowdhury<sup>a,39</sup> , T. Sarkar<sup>a</sup> , A. Scribano<sup>a</sup> , P. Spagnolo<sup>a</sup> , F. Tenchini<sup>a,b</sup> , R. Tenchini<sup>a</sup> , G. Tonelli<sup>a,b</sup> , N. Turini<sup>a,d</sup> , F. Vaselli<sup>a,c</sup> , A. Venturi<sup>a</sup> , P.G. Verдини<sup>a</sup> 

#### **INFN Sezione di Roma<sup>a</sup>, Sapienza Università di Roma<sup>b</sup>, Roma, Italy**

P. Barria<sup>a</sup> , C. Basile<sup>a,b</sup> , F. Cavallari<sup>a</sup> , L. Cunqueiro Mendez<sup>a,b</sup> , D. Del Re<sup>a,b</sup> , E. Di Marco<sup>a,b</sup> , M. Diemoz<sup>a</sup> , F. Errico<sup>a,b</sup> , R. Gargiulo<sup>a,b</sup> , E. Longo<sup>a,b</sup> , L. Martikainen<sup>a,b</sup> , J. Mijuskovic<sup>a,b</sup> , G. Organtini<sup>a,b</sup> , F. Pandolfi<sup>a</sup> , R. Paramatti<sup>a,b</sup> , C. Quaranta<sup>a,b</sup> , S. Rahatlou<sup>a,b</sup> , C. Rovelli<sup>a</sup> , F. Santanastasio<sup>a,b</sup> , L. Soffi<sup>a</sup> , V. Vladimirov<sup>a,b</sup>

**INFN Sezione di Torino<sup>a</sup>, Università di Torino<sup>b</sup>, Torino, Italy; Università del Piemonte Orientale<sup>c</sup>, Novara, Italy**

N. Amapane<sup>a,b</sup> , R. Arcidiacono<sup>a,c</sup> , S. Argiro<sup>a,b</sup> , M. Arneodo<sup>a,c</sup> , N. Bartosik<sup>a,c</sup> , R. Bellan<sup>a,b</sup> , C. Biino<sup>a</sup> , C. Borca<sup>a,b</sup> , N. Cartiglia<sup>a</sup> , M. Costa<sup>a,b</sup> , R. Covarelli<sup>a,b</sup> , N. Demaria<sup>a</sup> , M. Ferrero<sup>a</sup> , L. Finco<sup>a</sup> , M. Grippo<sup>a,b</sup> , B. Kiani<sup>a,b</sup> , F. Legger<sup>a</sup> , F. Luongo<sup>a,b</sup> , C. Mariotti<sup>a</sup> , L. Markovic<sup>a,b</sup> , S. Maselli<sup>a</sup> , A. Mecca<sup>a,b</sup> , L. Menzio<sup>a,b</sup> , P. Meridiani<sup>a</sup> , E. Migliore<sup>a,b</sup> , M. Monteno<sup>a</sup> , R. Mulargia<sup>a</sup> , M.M. Obertino<sup>a,b</sup> , G. Ortona<sup>a</sup> , L. Pacher<sup>a,b</sup> , N. Pastrone<sup>a</sup> , M. Ruspa<sup>a,c</sup> , F. Siviero<sup>a,b</sup> , V. Sola<sup>a,b</sup> , A. Solano<sup>a,b</sup> , C. Tarricone<sup>a,b</sup> , D. Trocino<sup>a</sup> , G. Umoret<sup>a,b</sup> , R. White<sup>a,b</sup> 

**INFN Sezione di Trieste<sup>a</sup>, Università di Trieste<sup>b</sup>, Trieste, Italy**

J. Babbar<sup>a,b</sup> , S. Belforte<sup>a</sup> , V. Candelise<sup>a,b</sup> , M. Casarsa<sup>a</sup> , F. Cossutti<sup>a</sup> , K. De Leo<sup>a</sup> , G. Della Ricca<sup>a,b</sup> 

**Kyungpook National University, Daegu, Korea**

S. Dogra , J. Hong , J. Kim, D. Lee, H. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , M.S. Ryu , S. Sekmen , B. Tae, Y.C. Yang 

**Department of Mathematics and Physics - GWNU, Gangneung, Korea**

M.S. Kim 

**Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea**

G. Bak , P. Gwak , H. Kim , D.H. Moon 

**Hanyang University, Seoul, Korea**

E. Asilar , J. Choi<sup>56</sup> , D. Kim , T.J. Kim , J.A. Merlin, Y. Ryou

**Korea University, Seoul, Korea**

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , S. Lee , J. Yoo 

**Kyung Hee University, Department of Physics, Seoul, Korea**

J. Goh , S. Yang 

**Sejong University, Seoul, Korea**

Y. Kang , H. S. Kim , Y. Kim, S. Lee

**Seoul National University, Seoul, Korea**

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , Y.W. Kim , S. Ko , H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

**University of Seoul, Seoul, Korea**

W. Jang , D.Y. Kang, S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

**Yonsei University, Department of Physics, Seoul, Korea**

S. Ha , K. Hwang , B. Kim , K. Lee , H.D. Yoo 

**Sungkyunkwan University, Suwon, Korea**

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 

**College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait**

T. Beyrouthy , Y. Gharbia 

**Kuwait University - College of Science - Department of Physics, Safat, Kuwait**

F. Alazemi 

**Riga Technical University, Riga, Latvia**

K. Dreimanis , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs, A. Potrebko , M. Seidel , D. Sidiropoulos Kontos 

**University of Latvia (LU), Riga, Latvia**

N.R. Strautnieks 

**Vilnius University, Vilnius, Lithuania**

M. Ambrozias , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 

**National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia**

I. Yusuff<sup>57</sup> , Z. Zolkapli

**Universidad de Sonora (UNISON), Hermosillo, Mexico**

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez, M. León Coello , J.A. Murillo Quijada , A. Sehwat , L. Valencia Palomo 

**Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico**

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, E. De La Cruz-Burelo , I. Heredia-De La Cruz<sup>58</sup> , R. Lopez-Fernandez , J. Mejia Guisao , A. Sánchez Hernández 

**Universidad Iberoamericana, Mexico City, Mexico**

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

**Benemerita Universidad Autonoma de Puebla, Puebla, Mexico**

I. Bautista , F.E. Neri Huerta , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

**University of Montenegro, Podgorica, Montenegro**

I. Bubanja , N. Raicevic 

**University of Canterbury, Christchurch, New Zealand**

P.H. Butler 

**National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**

A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan 

**AGH University of Krakow, Krakow, Poland**

V. Avati, A. Bellora , L. Forthomme , L. Grzanka , M. Malawski , K. Piotrkowski

**National Centre for Nuclear Research, Swierk, Poland**

H. Bialkowska , M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

**Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad 

**Warsaw University of Technology, Warsaw, Poland**

P. Fokow , K. Pozniak , W. Zabolotny 

**Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo 



T. Camporesi , G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff 

**Faculty of Physics, University of Belgrade, Belgrade, Serbia**

P. Adzic , P. Milenovic 

**VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia**

D. Devetak, M. Dordevic , J. Milosevic , L. Nadderd , V. Rekovic, M. Stojanovic 

**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**

J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. Llorente Merino , C. Martin Perez , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , J. Sastre , J. Vazquez Escobar 

**Universidad Autónoma de Madrid, Madrid, Spain**

J.F. de Trocóniz 

**Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain**

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , P. Leguina , E. Palencia Cortezon , J. Prado Pico , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia 

**Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain**

S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasiosa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizan Garcia 

**University of Colombo, Colombo, Sri Lanka**

B. Kailasapathy<sup>59</sup> , D.D.C. Wickramarathna 

**University of Ruhuna, Department of Physics, Matara, Sri Lanka**

W.G.D. Dharmaratna<sup>60</sup> , K. Liyanage , N. Perera 

**CERN, European Organization for Nuclear Research, Geneva, Switzerland**

D. Abbaneo , C. Amendola , E. Auffray , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , A.A. Bin Anuar , A. Bocci , L. Borgonovi , C. Botta , A. Bragagnolo , E. Brondolin , C.E. Brown , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , M. Glowacki, J. Hegeman , J.K. Heikkilä , B. Huber , V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban<sup>27</sup> , G. Karathanasis , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , M. Magherini , L. Malgeri , M. Mannelli , M. Matthewman, A. Mehta , F. Meijers , S. Mersi , E. Meschi , M. Migliorini , V. Milosevic , F. Monti , F. Moortgat , M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer , M. Pierini , M. Pitt , H. Qu , D. Rabadý , B. Ribeiro Lopes , F. Riti , M. Rovere , H. Sakulin , R. Salvatico , S. Sanchez Cruz , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas<sup>61</sup> 

A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea , D. Walter , J. Wanczyk<sup>62</sup> , J. Wang, S. Wuchterl , P. Zehetner , P. Zejdl , W.D. Zeuner

#### **PSI Center for Neutron and Muon Sciences, Villigen, Switzerland**

T. Bevilacqua<sup>63</sup> , L. Caminada<sup>63</sup> , A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli<sup>63</sup> , L. Noehte<sup>63</sup> , T. Rohe , A. Samalan

#### **ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland**

T.K. Aarrestad , M. Backhaus , G. Bonomelli , A. Calandri , C. Cazzaniga , K. Datta , P. De Bryas Dexmiers D'archiac<sup>62</sup> , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer , T.G. Harte, D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani , L. Marchese , A. Mascellani<sup>62</sup> , F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , B. Ristic , R. Seidita , J. Steggemann<sup>62</sup> , A. Tarabini , D. Valsecchi , R. Wallny 

#### **Universität Zürich, Zurich, Switzerland**

C. Amsler<sup>64</sup> , P. Bäertschi , M.F. Canelli , K. Cormier , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechti , A. Macchiolo , P. Meiring , F. Meng , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr, F. Stäger , R. Tramontano 

#### **National Central University, Chung-Li, Taiwan**

C. Adloff<sup>65</sup>, D. Bhowmik, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari<sup>37</sup> 

#### **National Taiwan University (NTU), Taipei, Taiwan**

L. Ceard, K.F. Chen , Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao, S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker , L.s. Tsai, D. Tsiou, H.y. Wu, E. Yazgan 

#### **High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand**

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand 

#### **Tunis El Manar University, Tunis, Tunisia**

Y. Maghrbi 

#### **Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey**

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu<sup>66</sup> , E. Eskut , Y. Guler<sup>67</sup> , E. Gurpinar Guler<sup>67</sup> , C. Isik , O. Kara, A. Kayis Topaksu , Y. Komurcu , G. Onengut , K. Ozdemir<sup>68</sup> , A. Polatoz , B. Tali<sup>69</sup> , U.G. Tok , E. Uslan , I.S. Zorbakir 

#### **Middle East Technical University, Physics Department, Ankara, Turkey**

M. Yalvac<sup>70</sup> 

#### **Bogazici University, Istanbul, Turkey**

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya<sup>71</sup> , O. Kaya<sup>72</sup> , S. Tekten<sup>73</sup> 

#### **Istanbul Technical University, Istanbul, Turkey**

A. Cakir , K. Cankocak<sup>66,74</sup> , S. Sen<sup>75</sup> 

#### **Istanbul University, Istanbul, Turkey**

O. Aydilek<sup>76</sup> , B. Haciasahinoglu , I. Hos<sup>77</sup> , B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , C. Zorbilmez 

#### **Yildiz Technical University, Istanbul, Turkey**

S. Cerci , B. Isildak<sup>78</sup> , D. Sunar Cerci , T. Yetkin 

**Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine**

A. Boyaryntsev , B. Grynyov 

**National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine**

L. Levchuk 

**University of Bristol, Bristol, United Kingdom**

D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran , L. Robertshaw , V.J. Smith , K. Walkingshaw Pass

**Rutherford Appleton Laboratory, Didcot, United Kingdom**

A.H. Ball , K.W. Bell , A. Belyaev<sup>79</sup> , C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , A. Elliot , K.V. Ellis , K. Harder , S. Harper , J. Linacre , K. Manolopoulos , D.M. Newbold , E. Olaiya , D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh , C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams 

**Imperial College, London, United Kingdom**

I. Andreou , R. Bainbridge , P. Bloch , O. Buchmuller , C.A. Carrillo Montoya , G.S. Chahal<sup>80</sup> , D. Colling , J.S. Dancu , I. Das , P. Dauncey , G. Davies , M. Della Negra , S. Fayer , G. Fedi , G. Hall , A. Howard , G. Iles , C.R. Knight , P. Krueper , J. Langford , K.H. Law , J. León Holgado , L. Lyons , A.-M. Magnan , B. Maier , S. Mallios , M. Mieskolainen , J. Nash<sup>81</sup> , M. Pesaresi , P.B. Pradeep , B.C. Radburn-Smith , A. Richards , A. Rose , L. Russell , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , T. Virdee<sup>29</sup> , M. Vojinovic , N. Wardle , D. Winterbottom 

**Brunel University, Uxbridge, United Kingdom**

J.E. Cole , A. Khan , P. Kyberd , I.D. Reid 

**Baylor University, Waco, Texas, USA**

S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish , J. Dittmann , K. Hatakeyama , V. Hegde , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant , C. Sutantawibul , J. Wilson 

**Catholic University of America, Washington, DC, USA**

R. Bartek , A. Dominguez , A.E. Simsek , S.S. Yu 

**The University of Alabama, Tuscaloosa, Alabama, USA**

B. Bam , A. Buchot Perraguin , R. Chudasama , S.I. Cooper , C. Crovella , S.V. Gleyzer , E. Pearson , C.U. Perez , P. Rumerio<sup>82</sup> , E. Usai , R. Yi 

**Boston University, Boston, Massachusetts, USA**

A. Akpinar , G. De Castro , Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , D. Gastler , F. Golf , S. Jeon , J. O'cain , I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , A.G. Zecchinelli 

**Brown University, Providence, Rhode Island, USA**

G. Barone , G. Benelli , D. Cutts , S. Ellis , L. Gouskos , M. Hadley , U. Heintz , K.W. Ho , J.M. Hogan<sup>83</sup> , T. Kwon , G. Landsberg , K.T. Lau , J. Luo , S. Mondal , T. Russell , S. Sagir<sup>84</sup> , X. Shen , M. Stamenkovic , N. Venkatasubramanian



### University of California, Davis, Davis, California, USA

S. Abbott , B. Barton , C. Brainerd , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , S. Ostrom , W. Wei , S. Yoo , F. Zhang 

### University of California, Los Angeles, California, USA

K. Adamidis, M. Bachtis , D. Campos, R. Cousins , A. Datta , G. Flores Avila , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , Y.f. Lo, E. Manca , A. Nunez Del Prado, D. Saltzberg , V. Valuev 

### University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , G. Hanson 

### University of California, San Diego, La Jolla, California, USA

A. Aportela, A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , L. Giannini , Y. Gu, J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , D. Primosch, M. Quinnan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil 

### University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan , C. Campagnari , K. Downham , C. Grieco , M.M. Hussain, J. Incandela , J. Kim , A.J. Li , P. Masterson , H. Mei , J. Richman , S.N. Santpur , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart , T.Á. Vámi , X. Yan , D. Zhang

### California Institute of Technology, Pasadena, California, USA

A. Albert, S. Bhattacharya , A. Bornheim , O. Cerri, J. Mao , H.B. Newman , G. Reales Gutiérrez, M. Spiropulu , J.R. Vlimant , S. Xie , R.Y. Zhu 

### Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , P. Bryant , M. Cremonesi, V. Dutta , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil, M. Kanemura, C. Liu , T. Mudholkar , S. Murthy , P. Palit , K. Park, M. Paulini , A. Roberts , A. Sanchez , W. Terrill 

### University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , N. Manganello , J. Parkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo 

### Cornell University, Ithaca, New York, USA

J. Alexander , X. Chen , D.J. Cranshaw , J. Dickinson , J. Fan , X. Fan , J. Grassi , S. Hogan , P. Kotamnives, J. Monroy , G. Niendorf, M. Oshiro , J.R. Patterson , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou 

### Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , C. Cosby , G. Cummings , I. Dutta , V.D. Elvira , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , T.C. Herwig , J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma 

B. Klima , K.H.M. Kwok , S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , K. Maeshima , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg, V. Papadimitriou , N. Pastika , K. Pedro , C. Pena<sup>85</sup> , F. Ravera , A. Reinsvold Hall<sup>86</sup> , L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , C. Wang , I. Zoi 

#### University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K. Matchev , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , Y. Takahashi , J. Wang 

#### Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhim , A. Askew , S. Bower , R. Hashmi , R.S. Kim , S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, M. Wulansatiti , R. Yohay , J. Zhang

#### Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani , S. Butalla , S. Das , T. Elkafrawy<sup>18</sup> , M. Hohlmann , E. Yanes

#### University of Illinois Chicago, Chicago, Illinois, USA

M.R. Adams , A. Baty , C. Bennett, R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , H. Gupta , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , D. S. Lemos , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , R. Pradhan , E. Prifti, P. Roy, T. Roy , S. Rudrabhatla , N. Singh, M.B. Tonjes , N. Varelas , M.A. Wadud , Z. Ye , J. Yoo 

#### The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini , D. Blend, K. Dilsiz<sup>87</sup> , L. Emediato , G. Karaman , O.K. Köseyan , J.-P. Merlo, A. Mestvirishvili<sup>88</sup> , O. Neogi, H. Ogul<sup>89</sup> , Y. Onel , A. Penzo , C. Snyder, E. Tiras<sup>90</sup> 

#### Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz 

#### The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , S. Popescu<sup>91</sup> , C. Rogan , C. Royon , S. Sanders , C. Smith , G. Wilson 

#### Kansas State University, Manhattan, Kansas, USA

B. Allmond , R. Gujju Gurunadha , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli , D. Roy , G. Sorrentino 

#### University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman, S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim, S. Lascio , A.C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , M.M. Paranjpe, E. Popova<sup>92</sup> , A. Shevelev , L. Wang , L. Zhang 

#### Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

C. Baldenegro Barrera , J. Bendavid , S. Bright-Thonney , I.A. Cali , P.c. Chou 

M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, G. Grosso, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn , A. Novak , M.I. Park , C. Paus , C. Reissel , C. Roland , G. Roland , S. Rothman , G.S.F. Stephans , Z. Wang , B. Wyslouch , T. J. Yang 

#### **University of Minnesota, Minneapolis, Minnesota, USA**

B. Crossman , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , M. Revering , R. Rusack , R. Saradhy , N. Strobbe 

#### **University of Nebraska-Lincoln, Lincoln, Nebraska, USA**

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , A. Rohilla , J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 

#### **State University of New York at Buffalo, Buffalo, New York, USA**

H. Bandyopadhyay , L. Hay , H.w. Hsia , I. Iashvili , A. Kalogeropoulos , A. Kharchilava , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams , P. Young 

#### **Northeastern University, Boston, Massachusetts, USA**

G. Alverson , E. Barberis , J. Bonilla , B. Bylsma, M. Campana , J. Dervan , Y. Haddad , Y. Han , I. Israr , A. Krishna , P. Levchenko , J. Li , M. Lu , R. McCarthy , D.M. Morse , T. Orimoto , A. Parker , L. Skinnari , E. Tsai , D. Wood 

#### **Northwestern University, Evanston, Illinois, USA**

S. Dittmer , K.A. Hahn , D. Li , Y. Liu , M. McGinnis , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliercio , M. Velasco

#### **University of Notre Dame, Notre Dame, Indiana, USA**

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth , K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , C. Moore , Y. Musienko<sup>15</sup> , H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygalá 

#### **The Ohio State University, Columbus, Ohio, USA**

A. Basnet , M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas , K. Wei, D.A. Wenzl, B.L. Winer , B. R. Yates 

#### **Princeton University, Princeton, New Jersey, USA**

H. Bouchamaoui , K. Coldham, P. Das , G. Dezoort , P. Elmer , P. Fackeldey , A. Frankenthal , B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , Y. Lai , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , F. Simpson , D. Stickland , C. Tully , L.H. Vage

#### **University of Puerto Rico, Mayaguez, Puerto Rico, USA**

S. Malik , R. Sharma

#### **Purdue University, West Lafayette, Indiana, USA**

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung , A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , J.F. Schulte , A. K. Viridi , F. Wang , A. Wildridge , W. Xie , Y. Yao 

#### **Purdue University Northwest, Hammond, Indiana, USA**

J. Dolen , N. Parashar , A. Pathak 

**Rice University, Houston, Texas, USA**

D. Acosta , A. Agrawal , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner, F.J.M. Geurts , I. Krommydas , W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 

**University of Rochester, Rochester, New York, USA**

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , N. Parmar , P. Parygin<sup>92</sup> , R. Taus 

**Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA**

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis , M. Heindl , C. Houghton , D. Jaroslowski , S. Konstantinou , I. Laflotte , A. Lath , R. Montalvo, K. Nash, J. Reichert , P. Saha , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora 

**University of Tennessee, Knoxville, Tennessee, USA**

D. Ally , A.G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A.R. Kanuganti , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 

**Texas A&M University, College Station, Texas, USA**

D. Aebi , M. Ahmad , T. Akhter , K. Androsov<sup>62</sup> , O. Bouhali<sup>93</sup> , R. Eusebi , J. Gilmore , T. Huang , T. Kamon<sup>94</sup> , H. Kim , S. Luo , R. Mueller , D. Overton , A. Safonov 

**Texas Tech University, Lubbock, Texas, USA**

N. Akchurin , J. Damgov , Y. Feng , N. Gogate , Y. Kazhykarim, K. Lamichhane , S.W. Lee , C. Madrid , A. Mankel , T. Peltola , I. Volobouev 

**Vanderbilt University, Nashville, Tennessee, USA**

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , D. Rathjens , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

**University of Virginia, Charlottesville, Virginia, USA**

B. Cardwell , H. Chung, B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , C. Mantilla , C. Neu , C. Ramón Álvarez 

**Wayne State University, Detroit, Michigan, USA**

S. Bhattacharya , P.E. Karchin 

**University of Wisconsin - Madison, Madison, Wisconsin, USA**

A. Aravind , S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless , J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , L. Pétré , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

**Authors affiliated with an international laboratory covered by a cooperation agreement with CERN**

S. Afanasiev , V. Alexakhin , Yu. Andreev , T. Aushev , D. Budkouski , A. Ershov , I. Golutvin<sup>†</sup> , I. Gorbunov , V. Karjavine , V. Klyukhin<sup>95</sup> , O. Kodolova<sup>96</sup> , V. Korenkov , A. Lanev , A. Malakhov , V. Matveev<sup>95</sup> , A. Nikitenko<sup>97,96</sup> , V. Palichik , V. Perelygin , S. Petrushanko<sup>95</sup> , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , V. Smirnov , O. Teryaev , A. Toropin , N. Voytishin , B.S. Yuldashev<sup>†98</sup>, A. Zarubin , I. Zhizhin 

**Authors affiliated with an institute formerly covered by a cooperation agreement with CERN**

G. Gavrillov , V. Golovtcov , Y. Ivanov , V. Kim<sup>95</sup> , V. Murzin , V. Oreshkin , D. Sosnov , V. Sulimov , L. Uvarov , A. Vorobyev<sup>†</sup>, A. Dermenev , S. Gninenko , N. Golubev , A. Karneyeu , D. Kirpichnikov , M. Kirsanov , N. Krasnikov , I. Tlisova , K. Ivanov , V. Gavrillov , N. Lychkovskaya , V. Popov , A. Zhokin , M. Chadeeva<sup>95</sup> , R. Chistov<sup>95</sup> , S. Polikarpov<sup>95</sup> , V. Andreev , M. Azarkin , M. Kirakosyan, A. Terkulov , E. Boos , V. Bunichev , M. Dubinin<sup>85</sup> , L. Dudko , O. Lukina , V. Savrin , A. Snigirev , V. Blinov<sup>95</sup>, T. Dimova<sup>95</sup> , A. Kozyrev<sup>95</sup> , O. Radchenko<sup>95</sup> , Y. Skovpen<sup>95</sup> , V. Kachanov , S. Slabospitskii , A. Uzunian , A. Babaev , V. Borshch , D. Druzhkin 

†: Deceased

<sup>1</sup>Also at Yerevan State University, Yerevan, Armenia

<sup>2</sup>Also at TU Wien, Vienna, Austria

<sup>3</sup>Also at Ghent University, Ghent, Belgium

<sup>4</sup>Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

<sup>5</sup>Also at FACAMP - Faculdades de Campinas, Sao Paulo, Brazil

<sup>6</sup>Also at Universidade Estadual de Campinas, Campinas, Brazil

<sup>7</sup>Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

<sup>8</sup>Also at University of Chinese Academy of Sciences, Beijing, China

<sup>9</sup>Also at China Center of Advanced Science and Technology, Beijing, China

<sup>10</sup>Also at University of Chinese Academy of Sciences, Beijing, China

<sup>11</sup>Also at China Spallation Neutron Source, Guangdong, China

<sup>12</sup>Now at Henan Normal University, Xinxiang, China

<sup>13</sup>Also at University of Shanghai for Science and Technology, Shanghai, China

<sup>14</sup>Now at The University of Iowa, Iowa City, Iowa, USA

<sup>15</sup>Also at an institute formerly covered by a cooperation agreement with CERN

<sup>16</sup>Also at Zewail City of Science and Technology, Zewail, Egypt

<sup>17</sup>Also at British University in Egypt, Cairo, Egypt

<sup>18</sup>Now at Ain Shams University, Cairo, Egypt

<sup>19</sup>Also at Purdue University, West Lafayette, Indiana, USA

<sup>20</sup>Also at Université de Haute Alsace, Mulhouse, France

<sup>21</sup>Also at Istinye University, Istanbul, Turkey

<sup>22</sup>Also at Tbilisi State University, Tbilisi, Georgia

<sup>23</sup>Also at The University of the State of Amazonas, Manaus, Brazil

<sup>24</sup>Also at University of Hamburg, Hamburg, Germany

<sup>25</sup>Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

<sup>26</sup>Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

<sup>27</sup>Also at Brandenburg University of Technology, Cottbus, Germany

<sup>28</sup>Also at Forschungszentrum Jülich, Juelich, Germany

<sup>29</sup>Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

<sup>30</sup>Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

<sup>31</sup>Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania

<sup>32</sup>Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

<sup>33</sup>Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

<sup>34</sup>Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt

<sup>35</sup>Also at Punjab Agricultural University, Ludhiana, India

<sup>36</sup>Also at University of Visva-Bharati, Santiniketan, India

<sup>37</sup>Also at Indian Institute of Science (IISc), Bangalore, India



- <sup>38</sup>Also at Amity University Uttar Pradesh, Noida, India
- <sup>39</sup>Also at UPES - University of Petroleum and Energy Studies, Dehradun, India
- <sup>40</sup>Also at IIT Bhubaneswar, Bhubaneswar, India
- <sup>41</sup>Also at Institute of Physics, Bhubaneswar, India
- <sup>42</sup>Also at University of Hyderabad, Hyderabad, India
- <sup>43</sup>Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- <sup>44</sup>Also at Isfahan University of Technology, Isfahan, Iran
- <sup>45</sup>Also at Sharif University of Technology, Tehran, Iran
- <sup>46</sup>Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- <sup>47</sup>Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- <sup>48</sup>Also at Helwan University, Cairo, Egypt
- <sup>49</sup>Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- <sup>50</sup>Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- <sup>51</sup>Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- <sup>52</sup>Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- <sup>53</sup>Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- <sup>54</sup>Also at Lulea University of Technology, Lulea, Sweden
- <sup>55</sup>Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- <sup>56</sup>Also at Institut de Physique des 2 Infinis de Lyon (IP2I ), Villeurbanne, France
- <sup>57</sup>Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- <sup>58</sup>Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- <sup>59</sup>Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- <sup>60</sup>Also at Saegis Campus, Nugegoda, Sri Lanka
- <sup>61</sup>Also at National and Kapodistrian University of Athens, Athens, Greece
- <sup>62</sup>Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- <sup>63</sup>Also at Universität Zürich, Zurich, Switzerland
- <sup>64</sup>Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- <sup>65</sup>Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- <sup>66</sup>Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- <sup>67</sup>Also at Konya Technical University, Konya, Turkey
- <sup>68</sup>Also at Izmir Bakircay University, Izmir, Turkey
- <sup>69</sup>Also at Adiyaman University, Adiyaman, Turkey
- <sup>70</sup>Also at Bozok Universiteleri Rektörlüğü, Yozgat, Turkey
- <sup>71</sup>Also at Marmara University, Istanbul, Turkey
- <sup>72</sup>Also at Milli Savunma University, Istanbul, Turkey
- <sup>73</sup>Also at Kafkas University, Kars, Turkey
- <sup>74</sup>Now at Istanbul Okan University, Istanbul, Turkey
- <sup>75</sup>Also at Hacettepe University, Ankara, Turkey
- <sup>76</sup>Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- <sup>77</sup>Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- <sup>78</sup>Also at Yildiz Technical University, Istanbul, Turkey
- <sup>79</sup>Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- <sup>80</sup>Also at IPPP Durham University, Durham, United Kingdom

---

<sup>81</sup>Also at Monash University, Faculty of Science, Clayton, Australia

<sup>82</sup>Also at Università di Torino, Torino, Italy

<sup>83</sup>Also at Bethel University, St. Paul, Minnesota, USA

<sup>84</sup>Also at Karamanoğlu Mehmetbey University, Karaman, Turkey

<sup>85</sup>Also at California Institute of Technology, Pasadena, California, USA

<sup>86</sup>Also at United States Naval Academy, Annapolis, Maryland, USA

<sup>87</sup>Also at Bingöl University, Bingöl, Turkey

<sup>88</sup>Also at Georgian Technical University, Tbilisi, Georgia

<sup>89</sup>Also at Sinop University, Sinop, Turkey

<sup>90</sup>Also at Erciyes University, Kayseri, Turkey

<sup>91</sup>Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania

<sup>92</sup>Now at another institute formerly covered by a cooperation agreement with CERN

<sup>93</sup>Also at Texas A&M University at Qatar, Doha, Qatar

<sup>94</sup>Also at Kyungpook National University, Daegu, Korea

<sup>95</sup>Also at another institute formerly covered by a cooperation agreement with CERN

<sup>96</sup>Also at Yerevan Physics Institute, Yerevan, Armenia

<sup>97</sup>Also at Imperial College, London, United Kingdom

<sup>98</sup>Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan