CERN-EP-2024-292
2025/06/19

CMS-EXO-21-016

Search for heavy neutral resonances decaying to tau lepton pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV

The CMS Collaboration*

Abstract

A search for heavy neutral gauge bosons (Z') decaying into a pair of tau leptons is performed in proton-proton collisions at $\sqrt{s} = 13$ TeV at the CERN LHC. The data were collected with the CMS detector and correspond to an integrated luminosity of 138 fb^{-1} . The observations are found to be in agreement with the expectation from standard model processes. Limits at 95% confidence level are set on the product of the Z' production cross section and its branching fraction to tau lepton pairs for a range of Z' boson masses. For a narrow resonance in the sequential standard model scenario, a Z' boson with a mass below 3.5 TeV is excluded. This is the most stringent limit to date from this type of search.

Published in Physical Review D as doi:10.1103/PhysRevD.111.112004.

1 Introduction

The CMS experiment [1] at the CERN LHC is conducting an extensive program of searches for evidence of new particles predicted by various extensions of the standard model (SM). One of the simplest extensions postulates an additional $U(1)$ symmetry group that gives rise to the production of a new neutral gauge particle, analogous to the Z boson in the SM, commonly referred to as a Z' boson. There are other more complicated extensions of the SM that also predict the existence of similar mediators, such as grand unified theories [2], $E(6)$ models [3], and the Randall–Sundrum model [4]. Searches for Z' bosons have been carried out by the ATLAS and CMS Collaborations, placing stringent bounds on Z' boson production as a function of its mass, $m_{Z'}$, especially for the $Z' \rightarrow \mu^+\mu^-$ and e^+e^- decay channels. Combining the $\mu^+\mu^-$ and e^+e^- final states and using the sequential SM (SSM) extension, the production of an SSM Z' boson has been excluded at 95% confidence level (CL) for masses below 5.15 (5.10) TeV in searches reported by CMS [1] (ATLAS [5]). The SSM assumes that the Z' boson has the same couplings to SM particles as those of the Z boson, in particular, couplings to SM fermions that are independent of their generation. There are several models that instead propose preferential couplings of the Z' boson to third-generation fermions [6, 7]. In models with nonuniversal lepton couplings, assuming a Z' boson with the same couplings to third generation fermions as the SM Z boson, bounds on $m_{Z'}$ are found for the $Z' \rightarrow \tau^+\tau^-$ decay mode [8, 9] excluding $m_{Z'}$ up to 2.42 TeV and the $Z' \rightarrow b\bar{b}$ decay mode [10, 11] excluding $m_{Z'}$ up to 2.7 TeV.

In recent years, interest in potential effects beyond the SM has emerged as a result of anomalies observed in the precision measurements of B meson decay rates, including $R(D^{(*)})$ [12–22], the ratio of branching fractions with leptons of different flavors. The $R(D^{(*)})$ ratio is sensitive to the violation of lepton flavor universality in $b \rightarrow c l \bar{\nu}_l$ transitions, where l denotes any lepton flavor. These anomalies have led to the proposal of a large number of theoretical models that predict the existence of heavy neutral and charged gauge bosons (Z' and W' , respectively) with nonuniversal fermion couplings within a new $SU(2)$ group. These models further motivate the search for a Z' boson that preferentially couples to third-generation fermions [23, 24]. For this search, we use a simplified phenomenological model, in which the Z' boson mass and couplings are treated as free parameters [25]. In this paper, we consider a Z' boson with the same fermion couplings as the Z boson to compare with previous searches, but also consider scenarios with varied Z' boson couplings to the τ leptons.

This paper presents a search for a Z' boson via the $\tau^+\tau^-$ decays, considering production via quark-antiquark annihilation, in proton-proton (pp) collisions at $\sqrt{s} = 13$ TeV at the LHC. The data were collected with the CMS experiment during 2016–2018, and correspond to an integrated luminosity of 138 fb^{-1} . Tabulated data are provided in the HEPData record [26] for this analysis.

The remaining sections of this paper present the analysis strategy, in Section 2, followed by a short detector description in Section 3. The event reconstruction is detailed in Section 4 and the generation of simulated samples in Section 5. We describe the event selection criteria and background estimation in Sections 6 and 7, respectively, for each of the τ pair decay channels. Taking into account systematic uncertainties as discussed in Section 8, we present the results in Section 9 and a summary in Section 10.

2 Analysis strategy

The τ decay channels can be broadly classified into hadronic (τ_h), $\nu_\tau + \text{hadrons}$, and leptonic (τ_ℓ), $\nu_\tau + \ell + \bar{\nu}_\ell$, with ℓ representing μ or e . The analysis presented here includes the three $Z' \rightarrow \tau^+ \tau^-$ decay channels with the largest τ decay branching fraction products: $Z' \rightarrow \tau^\pm_\mu \tau^\mp_h$, $Z' \rightarrow \tau^\pm_e \tau^\mp_h$, and $Z' \rightarrow \tau^+_h \tau^-_h$, referred to below as $\tau_\mu \tau_h$, $\tau_e \tau_h$, and $\tau_h \tau_h$, respectively.

In the production of Z' bosons via s -channel annihilation of quark-antiquark pairs, the Z' bosons have limited Lorentz boost in the plane transverse to the beam. For the TeV-scale $m_{Z'}$ of interest, the τ decay daughters tend to be clustered within small back-to-back cones. With this picture in mind, we derive an estimate of the Z' boson candidate mass $m_{\text{rec}}(Z')$, making use of the transverse momentum imbalance between the visible daughters of the two τ lepton decays:

$$m_{\text{rec}}(Z') = \sqrt{(E_1^{\tau\text{vis}} + E_2^{\tau\text{vis}} + |\mathbf{p}^{Z'\text{miss}}|)^2 - |\mathbf{p}_1^{\tau\text{vis}} + \mathbf{p}_2^{\tau\text{vis}} + \mathbf{p}^{Z'\text{miss}}|^2}. \quad (1)$$

Here $(E_i^{\tau\text{vis}}, \mathbf{p}_i^{\tau\text{vis}})$ is the four momentum of the reconstructed daughter(s) of the i^{th} τ candidate. The momentum of the invisible neutrinos is represented by $\mathbf{p}^{Z'\text{miss}} = (\vec{p}_T^{Z'\text{miss}}, p_z^{Z'\text{miss}})$, with $\vec{p}_T^{Z'\text{miss}} = -(\vec{p}_{1T}^{\tau\text{vis}} + \vec{p}_{2T}^{\tau\text{vis}})$ and $p_z^{Z'\text{miss}} = 0$. We find that this definition of $\vec{p}_T^{Z'\text{miss}}$ provides better discrimination between the signal and the SM background processes than the event momentum imbalance $\vec{p}_T^{\text{miss}}$, the negative vectorial sum of all reconstructed objects. This is because $\vec{p}_T^{\text{miss}}$ includes contributions from mismeasured jets in the underlying event and from additional pp interactions within the same or nearby bunch crossings (pileup).

We use $m_{\text{rec}}(Z')$ as the primary discriminating variable for extraction of the signal yield. Although the natural decay width of the Z' boson is small compared with its mass ($<3\%$ for the model assumptions in this paper), the relative resolution on $m_{\text{rec}}(Z')$ is between 30 and 45% for the range of $m_{Z'}$ considered. The analysis is thus dominated by experimental resolution effects arising from the undetected neutrinos. Data and simulation are used to obtain estimates of the yields of the SM background processes, and the signal yield is obtained from a maximum likelihood (ML) fit to the $m_{\text{rec}}(Z')$ distributions for each of the three $Z' \rightarrow \tau^+ \tau^-$ decay channels, as described in the following sections.

3 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter, each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are reconstructed in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. More detailed descriptions of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Refs. [27, 28].

Events of interest are selected using a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about 4 μs [29]. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage [30].

4 Object reconstruction

The first stage of particle reconstruction employs the particle-flow (PF) algorithm [31], which aims to reconstruct and identify each individual visible particle in an event (photon, electron, muon, charged hadron, neutral hadron) with an optimized combination of all subdetector information. Further identification (ID) strategies for particular particle types are applied to achieve an appropriate balance between efficiency and the purity of the resulting candidate collections.

In the presence of pileup, the primary vertex, at which the pp collision of interest occurs, is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in Section 9.4.1 of Ref. [32].

Jets are reconstructed offline from PF candidates using the anti- k_T clustering algorithm [33, 34] with a distance parameter of 0.4. Jet momentum is determined as the vectorial sum of all particle momenta in the jet, and is found from simulation to be, on average, within 5 to 10% of the true momentum over the entire p_T spectrum and detector acceptance. Pileup interactions can contribute additional tracks and calorimetric energy depositions, increasing the apparent jet momentum. To mitigate this effect, tracks identified to be originating from pileup vertices are discarded, and an offset correction is applied to correct for remaining contributions [35]. Jet energy corrections are derived from simulation studies so that the average measured energy of jets matches that of particle level jets. In situ measurements of the momentum balance in dijet, photon + jet, Z + jet, and multijet events are used to determine any residual differences between the jet energy scale in data and in simulation, and appropriate corrections are made [36]. Additional selection criteria are applied to each jet to remove jets potentially dominated by instrumental effects or reconstruction failures [35].

The $\vec{p}_T^{\text{miss}}$ vector is computed as the negative vector sum of the transverse momenta of all the PF candidates in an event, modified to account for corrections to the energy scale of the reconstructed jets in the event [37]. Its magnitude is denoted p_T^{miss} .

Muons are measured in the pseudorapidity range $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive-plate chambers. The single-muon trigger efficiency exceeds 90% over the full η range, and the efficiency to reconstruct and identify muons is greater than 96%. Matching muons to tracks measured in the silicon tracker results in a relative transverse momentum resolution of 1% in the barrel and 3% in the endcaps for muons with p_T up to 100 GeV. The p_T resolution in the barrel is better than 7% for muons with p_T up to 1 TeV [38]. This analysis uses the tight working point [38] of the muon ID. To suppress muons from hadron decays and other nonprompt sources, an additional requirement is imposed on the relative isolation, defined as the ratio of the energy from neutral and charged PF candidates in a cone of $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} < 0.4$ around the muon trajectory to that of the muon. A correction for energy mismeasurement from pileup is applied, and we require this corrected relative isolation to be smaller than 0.15.

Electrons are reconstructed using energy deposits in the ECAL detector that have a matching track in (η, ϕ) space in the silicon tracking system, within $|\eta| < 2.4$. Electron candidates that fall in the transition region between the barrel and the endcap of the CMS detector, located at $1.44 < |\eta| < 1.57$, are not considered. The momentum resolution for electrons with $p_T \approx 45$ GeV from $Z \rightarrow ee$ decays ranges from 1.6–5.0%. It is better in the barrel region than in the endcaps, and depends on the bremsstrahlung energy emitted by the electron as it traverses the material in front of the ECAL [39, 40]. To identify electrons, we use the high-energy electron pairs (HEEP) ID for the selection of signal events; for veto and control region (CR) selections,

we use instead the loose working point of the cut-based electron ID [40]. Isolation is defined as the amount of energy in a cone of $\Delta R < 0.3$ around the electron direction. It is an integral part of both electron ID criteria used, as defined in detail in [40], and both IDs utilize electron-energy-dependent isolation criteria with pileup energy corrections that differ between the barrel and endcap regions of the detector.

Hadronic τ lepton decays are reconstructed from jets using the hadrons-plus-strips algorithm [41], which combines 1 or 3 tracks with energy deposits in the calorimeters to form τ_h candidates. Neutral pions are reconstructed as strips with dynamic size in (η, ϕ) from reconstructed electrons and photons, where the strip size varies as a function of the p_T of the electron or photon candidate.

The DEEPTAU deep neural network algorithm [42] is used to distinguish genuine τ_h candidates from jets originating from the hadronization of quarks or gluons, and from electrons or muons. Information from all individually reconstructed particles near the τ_h momentum direction is combined with properties of the τ_h candidate and of the event. We make use of the tight (medium) working point for the discrimination of τ_h from muons (electrons), and of the tight, loose, and very loose working points for discrimination against jets, as described in Sections 6 and 7.

To identify jet candidates from b quark hadronization (hereafter denoted b jets) we use the DEEPCSV b tagging algorithm [43], which is based on a deep neural network. We employ the tight working point of the discriminator, for which the typical efficiency is 50%, while the mistag rate of jets originating from light-flavored quarks or gluons is 0.1% and from c jets is 2.4%.

5 Simulated samples

Monte Carlo simulation is used in the analysis to design the event selection and to model the signal and SM background processes. For estimates of the background yields we use a combination of simulation and data, as described in Section 7.

The background from SM events composed uniquely of jets produced via the strong interaction is referred to as quantum chromodynamics (QCD) multijet events. Events from Drell-Yan (DY), QCD multijet, single t quark, and $W + \text{jets}$ processes are simulated with the MADGRAPH5_aMC@NLO 2.6.0 event generator [44] at leading order (LO) accuracy. The POWHEG 2.0 [45–47] event generator at next-to-LO (NLO) accuracy in QCD is used to simulate $t\bar{t}$ events, while diboson (VV) processes are produced at LO with PYTHIA 8.226 (8.230) for data taken in 2016 (2017 and 2018) [48].

Signal events are generated using MADGRAPH5_aMC@NLO at LO [49]. The influence of interference effects between the Z' signal diagrams and the SM processes are found to be small, resulting in no significant impact on the interpretation of the results, for the range of Z' boson masses and couplings considered in this analysis. A set of samples is produced for Z' boson masses ranging from 250 to 3000 GeV in steps of 250 GeV, as well as at 4000 and 5000 GeV. The primary set of benchmark signal samples for interpretation of the results have the Z' boson couplings set to SM Z values. Additionally, scenarios with varied modifiers g_τ of the couplings to τ leptons are considered. The theoretical decay width of the Z' boson in the production model increases with g_τ but remains smaller than the experimental resolution ($< 30\%$) for the range of g_τ values considered in this study.

The MADGRAPH5_aMC@NLO and POWHEG generators used to simulate signal and background

processes are interfaced with PYTHIA 8.226 (8.230) using the CUETP8M1 (CP5) tunes [50, 51] for parton showering and fragmentation in the 2016 (2017–2018) simulated samples. The NNPDF3.0 (3.1) NNLO [52] parton distribution functions (PDFs) are used in the event generation for 2016 (2017–2018) samples. The cross sections used to normalize the simulated background yields are calculated at next-to-NLO or NLO accuracy [44, 53–63], while those used for the signal yields are calculated at NLO accuracy.

The detector response is modeled with GEANT4 [64]. Pileup is incorporated by simulating additional minimum bias interactions with the multiplicity distribution matching that measured in data.

6 Event selection criteria

The signal region (SR) selection criteria for each $Z' \rightarrow \tau^+ \tau^-$ decay channel are chosen to maximize the expected signal significance by balancing signal efficiency against the suppression of background. The main SM background processes are DY, $W + \text{jets}$, $t\bar{t}$, QCD multijet, and VV (WW , ZZ , and WZ) production. The DY process constitutes an irreducible background since $\tau^+ \tau^-$ pairs arise from the intermediate Z and photon states. However, since the analysis focuses on massive Z' bosons, the visible daughters from the two τ lepton decays typically have higher momentum and larger angular separation than those from DY events. We exploit these characteristics to separate heavy Z' boson signals from background. Events from $W + \text{jets}$ and $t\bar{t}$ can mimic the $\tau_\ell \tau_h$ final states when the W boson, produced directly or from a top quark decay, decays via $W \rightarrow \ell \nu_\ell$, and a jet is misidentified as a τ_h candidate. Events from VV or $t\bar{t}$ production with leptonic decays of both vector bosons produce genuine lepton pairs, leading to final states with a $\tau^+ \tau^-$ pair as in signal events. Multijet events become a background source through the misidentification of jets as τ_h or ℓ candidates.

Events are required to have one τ_h candidate accompanied by a μ , e, or second τ_h candidate of opposite-sign (OS) charge, all within $|\eta| < 2.1$. To avoid possible overlaps among the three channels, we reject events containing additional lepton candidates of any flavor beyond the selected OS lepton candidate pair. Specifically, the additional lepton can be a μ or e candidate with $p_T > 10 \text{ GeV}$ and $|\eta| < 2.1$, or a τ candidate meeting the τ_h requirements of the particular channel. Since the analysis explores the production of Z' bosons through quark-antiquark annihilation, events from vector boson fusion are suppressed by vetoing events with a pair of jets $j^{1,2}$, each of which passes $p_T > 30 \text{ GeV}$ and $|\eta| < 4.7$, with a pseudorapidity separation $|\Delta\eta(j^1, j^2)| > 4.2$ and an invariant mass above 500 GeV .

For the ID of τ_h candidates in the SRs for all decay channels we impose the tight working points of the discriminators against jets and muons, and the medium one against electrons, as defined in Ref. [42]. The probability of misidentifying a jet as a τ_h candidate depends on the p_T and flavor of the jet; for the kinematic conditions of this analysis it has been estimated in a $W + \text{jets}$ sample to be approximately 0.6%, with a genuine τ_h ID efficiency of 60%. The probability for muons (electrons) to be misidentified as τ_h candidates is 0.07% (0.2%) for a genuine τ_h ID efficiency of $>99\%$ (90%).

6.1 The $\tau_\mu \tau_h$ SR

The $\tau_\mu \tau_h$ events are required to satisfy the single-muon trigger, whose efficiency exceeds 90% over the full η range for the selected muon candidates with $p_T > 35 \text{ GeV}$. Muons are also required to be well isolated and to pass the tight ID criteria defined in Ref. [38]. The τ_h candidate is required to satisfy $p_T > 20 \text{ GeV}$. The two candidates are required to be well separated in

(η, ϕ) space by the criterion $\Delta R \equiv \sqrt{(\Delta\phi(\tau_\mu, \tau_h))^2 + (\Delta\eta(\tau_\mu, \tau_h))^2} > 0.3$. We reject events containing b jet candidates with $p_T > 30$ GeV and $|\eta| < 2.4$. Events from DY, W + jets, and QCD multijet production are significantly suppressed by requiring the τ_μ and τ_h candidates to have a large azimuthal separation given by $\cos \Delta\phi(\tau_\mu, \tau_h) < -0.98$. In addition, we require that $\vec{p}_T^{\text{miss}}$ lie in the direction opposite that of the τ_h or of the τ_μ candidate with the highest p_T (leading lepton, ℓ^1), by requiring $\cos \Delta\phi(p_T^{\text{miss}}, \ell^1) < -0.95$. This requirement further reduces the contribution of W + jets and QCD multijet events. The transverse mass of the leading lepton and $\vec{p}_T^{\text{miss}}$, $m_T(p_T^{\text{miss}}, \ell^1) = \sqrt{p_T^{\text{miss}} p_T(\ell^1) (1 - \cos(\Delta\phi(p_T^{\text{miss}}, \ell^1)))}$, is required to be greater than 150 GeV for further suppression of W + jets events.

6.2 The $\tau_e \tau_h$ SR

Similar selection criteria are applied to the $\tau_e \tau_h$ channel, with the following differences. We require these events to satisfy a single-electron trigger that has an efficiency above 90% for electrons after the requirement $p_T > 35$ (55) GeV for data collected in 2016 (2017–2018).

6.3 The $\tau_h \tau_h$ SR

For the $\tau_h \tau_h$ channel we select events that satisfy a dedicated trigger [29] with at least two τ_h candidates. We require each τ_h candidate to have $p_T > 70$ GeV, ensuring a trigger efficiency of at least 90%. The two τ_h candidates must be separated by $\Delta R > 0.3$. Events with any b jet candidate having $p_T > 30$ GeV and $|\eta| < 2.4$ are removed, to suppress top quark backgrounds. To reduce the contribution of DY events, the reconstructed mass of the τ_h pair is required to exceed 100 GeV. To discriminate against W + jets and QCD multijet events, we require the two τ_h candidates to have a large azimuthal separation, $\cos \Delta\phi(\tau_h^1, \tau_h^2) < -0.95$, while the $\vec{p}_T^{\text{miss}}$ and the leading- p_T τ_h candidate τ_h^1 are required to satisfy $|\cos \Delta\phi(p_T^{\text{miss}}, \tau_h^1)| > 0.9$. Further suppression of the contribution of QCD multijet events is achieved with a requirement $p_T^{\text{miss}} > 30$ GeV.

7 Background estimation

The background yields are estimated in discrete intervals (bins) in the variable $m_{\text{rec}}(Z')$, by methods that depend on the SM process and the channel. In most cases, these estimates are based on samples of simulated events. Differences between data and simulation with respect to the efficiency of triggers, particle ID or isolation, and energy scale and resolution of electrons, muons, jets, and p_T^{miss} are taken into account with dedicated correction factors whose uncertainties are propagated throughout. These correction factors are described in the references cited in Section 4. Residual mismodeling can remain due to systematic effects of the theoretical predictions, or to the phase space requirements specific to this analysis or to a specific channel. To ensure that the effect of any such mismodeling and its uncertainty in the background yield is accounted for, we evaluate a data-to-simulation scale factor (SF) in a CR. The systematic uncertainties in the SFs account for correlations between CRs and between the CRs and SRs. Validation tests are performed using additional simulation and data control samples to ensure the correct determination of the SFs and their applicability to the SR background prediction.

The SFs are found to be approximately independent of $m_{\text{rec}}(Z')$, and are thus applied to the $m_{\text{rec}}(Z')$ distributions as bin-independent normalization factors. Uncertainties from the residual shape dependence are evaluated as described in Section 8. The SFs are derived for each data-taking year separately, along with their uncertainties and year-to-year correlations. The values of the SFs and their uncertainties are given in the following subsections. With a few

exceptions, these values are consistent with unity. In the case of the QCD multijet background, the estimate is derived from the yield measured in dedicated orthogonal data control samples using the ABCD method described in Section 7.1. For SM processes that contribute less than 5% to the background yield, the simulation is taken directly.

Each CR is constructed to be enriched in a given SM process and orthogonal to the corresponding SR, but with characteristics similar to those of the SR. We correct for contamination from other SM processes by subtracting their SF-corrected simulated yields. The resulting dependence of the estimate of one background process on the SF for another is treated through the uncertainty correlation terms in the ML fit.

7.1 The $\tau_\ell\tau_h$ CRs

The main sources of background events in the $\tau_\ell\tau_h$ channels are DY, $W + \text{jets}$, and $t\bar{t}$ production.

To derive the SFs for the DY background in the $\tau_\ell\tau_h$ channels we define CRs by selecting events as for the SRs except for an inverted m_T criterion, $m_T < 150 \text{ GeV}$, and a requirement that the reconstructed mass of the $\tau_\ell\tau_h$ pair lie within the Z boson mass window, $70 < m_{\text{rec}}(\tau_\ell\tau_h) < 110 \text{ GeV}$. Since the higher p_T threshold for electrons of 55 GeV in 2017–2018 would deplete the low end of this mass range for a $\tau_e\tau_h$ CR, for those data sets we substitute the SFs measured in the corresponding $\tau_\mu\tau_h$ channels. The values of the DY SFs measured in the $\tau_\mu\tau_h$ channel are 0.98 ± 0.06 , 0.91 ± 0.05 , and 0.90 ± 0.06 in 2016, 2017, and 2018, respectively. The value of the DY SFs for the $\tau_e\tau_h$ channel in 2016 is 1.02 ± 0.07 .

We measure the SF for the $W + \text{jets}$ background in a CR defined to be orthogonal to the SR by inverting the τ_h selection, i.e., by requiring that the τ_h candidates fail the tight discrimination against jets while passing the very loose selection. We also modify the $m_T > 150 \text{ GeV}$ SR requirement so as to include W boson production; we select events with $m_T < 190 \text{ GeV}$, where the extended upper limit serves to probe the $W + \text{jets}$ background in the high mass tail of the $m_{\text{rec}}(Z')$ distribution. The contribution of QCD multijet events in this CR is nonnegligible, amounting to 30–40% depending on the channel and data-taking year; it is estimated from data following the same methodology as that described below in this section for estimating the QCD multijet background in the SR. In the $\tau_\mu\tau_h$ channel, the SFs are 0.98 ± 0.03 , 0.98 ± 0.03 , and 0.92 ± 0.03 in 2016, 2017, and 2018, respectively. For the $\tau_e\tau_h$ channel, the SFs are 1.02 ± 0.07 , 0.94 ± 0.08 , and 0.92 ± 0.09 in 2016, 2017, and 2018, respectively.

For $t\bar{t}$ events, we define the CR by imposing the SR selection criteria but changing the required number of b jet candidates from zero to exactly one. The requirement of exactly one b jet candidate ensures that the CR is kinematically similar to the SR. For 2016, 2017 and 2018, the SFs in the $\tau_\mu\tau_h$ ($\tau_e\tau_h$) channel are 1.02 ± 0.07 , 0.80 ± 0.04 , and 0.82 ± 0.04 (0.90 ± 0.11 , 0.85 ± 0.05 , and 0.85 ± 0.05), respectively. The difference in the scale factors across years is due to a known issue with track reconstruction during part of the 2016 data taking.

In all CRs we find agreement between data and simulation in the $m_{\text{rec}}(Z')$ spectrum, after the corresponding SFs are applied to the simulation.

For the QCD multijet background estimate, we apply an ABCD method, in which four orthogonal regions are constructed with two measured observables. The estimate of the background yield N_A in the SR is then determined from the yields in CRs B, C, and D as $N_A = (N_C/N_D)N_B$. The necessary lack of correlation between the two observables has been validated in a separate ABCD set with $p_T^{\text{miss}} < 30 \text{ GeV}$ in the hadronic di-tau channel. Here the first observable is the charge-sign pairing of the τ candidates: the OS requirement of the SR is reversed to same-sign

(SS) to define CR B. The second observable is the appropriate measure of τ purity, namely lepton ID for τ_ℓ , isolation for τ_h : both tight, as in the SR, for region B, while for regions C (OS) and D (SS) both must fail tight and pass loose ID. Contributions from background sources different from QCD have been removed in all CRs. The calculation is performed for each bin of the $m_{\text{rec}}(Z')$ distribution, separately.

7.2 The $\tau_h\tau_h$ CRs

The main sources of background events in the $\tau_h\tau_h$ channel are DY and QCD multijet events.

For the DY component, the SF is measured in a CR defined by the SR criteria except that the τ_h candidate pair's invariant mass $m(\tau_h, \tau_h)$ is required to satisfy $m(\tau_h, \tau_h) < 100$ GeV. The DY SFs measured for the $\tau_h\tau_h$ channel are 1.11 ± 0.10 , 0.96 ± 0.09 , and 0.90 ± 0.08 in 2016, 2017, and 2018, respectively.

The QCD multijet background is estimated as follows in an ABCD approach: A CR B is selected by the SR criteria except requiring an SS τ_h candidate pair. The common requirements for the events in the C and D CRs are $p_T^{\text{miss}} < 30$ GeV, and that both τ_h candidates fail the tight, while passing the very loose, discrimination against jets. Events entering CR C (D) pass the OS (SS) charge requirements. To ensure that the sideband selection of passing the very loose and failing the tight τ_h discriminator against jets in the low- p_T^{miss} sideband does not introduce any bias, the same method was also tested without the additional requirement of $p_T^{\text{miss}} < 30$ GeV, which was introduced to improve the QCD sample purity. The results of this test, using the τ_h sideband selection, were found to be statistically compatible with those obtained from the nominal selection, but with improved statistical precision in the sideband region. Therefore, the additional discriminator inversion method was adopted for the final result of the QCD estimate.

7.3 The CR mass distributions

Figure 1 shows the $m_{\text{rec}}(Z')$ distributions for the data and simulation in the CRs that are used to compute the SFs. Here the yields are summed over the data-taking years and are measured prior (pre-fit) to the ML fit that is used to extract the signal yields.

8 Systematic uncertainties

We account for systematic uncertainties associated with the various sources as detailed in the following paragraphs.

The integrated luminosities for the 2016, 2017, and 2018 data-taking years have individual uncertainties of 1.2–2.5% [65–67], while the overall uncertainty for the 2016–2018 period is 1.6%. This uncertainty is applied to all background and signal processes.

Uncertainties in the pileup modeling corrections are evaluated by varying the total inelastic pp cross section used in the correction procedure by $\pm 5\%$ [68, 69]. The effect of the resulting change in the distribution of the number of primary vertices is less than 0.01%, and is therefore treated as negligible.

The efficiencies for the electron and muon reconstruction, ID, and isolation are measured with a “tag-and-probe method” [38, 40, 70] with a resulting uncertainty of $\leq 2\%$, depending on p_T and η . The total efficiency for the τ_h ID and isolation requirements is measured from a fit to the $\mu\tau_h$ invariant mass distribution from $Z \rightarrow \tau\tau$ events, not taking into account neutrino hypotheses or p_T^{miss} . This is done in a sample selected with one isolated μ trigger candidate with




Figure 1: Pre-fit reconstructed mass distributions for the (upper row) W + jets, (middle row) $t\bar{t}$, and (lower row) DY CRs, for the (left column) $\tau_\mu\tau_h$, (right column, upper and middle rows) $\tau_e\tau_h$, and (lower right) $\tau_h\tau_h$ channel. The targeted background in each CR, denoted by its CR name, has been corrected with the derived SF from that CR. The background contributions appear as stacked solid histograms, while the data are shown as black solid markers with error bars. Overflow counts are included in the last bin. The gray shadow band represents the statistical uncertainty in the background prediction based on the number of simulated events and SF uncertainties. The lower panel of each plot shows the ratio of data to predicted background yields.

$p_T > 24 \text{ GeV}$, and leads to a relative uncertainty of less than 9% per genuine τ_h candidate [41], dependent on p_T and η . These uncertainties have contributions that are either statistical and thereby uncorrelated, or systematic and therefore correlated across the data-taking years. In the cases where background estimates rely on the modeling of the misidentification of jets as τ_h candidates, additional systematic uncertainties in the corresponding background yields are assigned that range from 20 to 30%.

For the lepton momentum scale, the reconstructed momentum is shifted by 5% and the $\vec{p}_T^{\text{miss}}$ vector recalculated. This results in a change in the expected signal yield of 1–6%, depending on $m_{Z'}$, and negligible change for the backgrounds. The corresponding resolution uncertainty in the simulation is taken as 0.03 times the difference between the reconstructed and true values of the lepton's momentum [38]. It is found to have negligible impact on either signal or background yields.

Similarly, we include the uncertainties in the energy scale and resolution of jets and the propagation thereof to p_T^{miss} [36]. The impact on the SR selection is indirect, affecting only the vetos on b-tagged jets and VBF dijet candidates. Accordingly, the signal event yield is not affected by either uncertainty, while the background event yields vary, depending on the process, by less than 2%.

We propagate the uncertainty in the b jet mistagging SF [43] to its effect on the b jet requirements in the SR selection. The resulting uncertainty in the signal yield is found to be $\sim 1\%$. The effect of the b tagging efficiency uncertainty on the mass distribution in the $t\bar{t}$ CRs for the $\tau_\ell\tau_h$ channels is measured by varying the b tagging SF up and down by its uncertainty; the resulting uncertainty in the $t\bar{t}$ SF is about 2%.

The uncertainties in the cross sections used to predict the rates of various processes, which arise primarily from the factorization and renormalization scale choices and the PDFs, are propagated to the yield estimates. The cross section uncertainties are each separated into their scale (renormalization and factorization scales) and PDF components. They are correlated where appropriate among processes. To derive systematic uncertainties in the predicted signal yields arising from the choice of PDFs used in the simulation, we vary the PDFs following the PDF4LHC prescription [71]. We find that these variations do not affect the shapes of kinematic distributions, but the resulting systematic uncertainty in the overall signal acceptance prediction is 2–6%, depending on the assumed value of $m_{Z'}$.

Uncertainties in the renormalization and factorization scales are evaluated as the differences in predicted signal yields resulting from independent variation of these scales from their default values by factors of 2.0 and 0.5 [72]. The resulting uncertainties in the yields are $< 2\%$, independent of $m_{Z'}$.

The statistical uncertainty associated with the SFs measured in the CRs is propagated to the total uncertainty in the expected yields in the SR. The SFs in the $t\bar{t}$ CRs, where some dependence on $m_{\text{rec}}(Z')$ is observed, are fit with a first-order polynomial as a function of $m_{\text{rec}}(Z')$. The deviation of the resulting function from unity is taken as the systematic uncertainty in the shape. The effect spans the range 1–22%, depending on the $m_{\text{rec}}(Z')$ bin and data-taking year.

For the QCD transfer factors N_C/N_D , the statistical uncertainties are in the ranges 19–23% for $\tau_\mu\tau_h$, 23–26% for $\tau_e\tau_h$, and 1.7–2.1% for $\tau_h\tau_h$. For each of the $\tau_\mu\tau_h$ and $\tau_e\tau_h$ channels, the QCD contribution to the SR is small (6–9%), while it is large for the $\tau_h\tau_h$ channel (27–35%), albeit mostly at low $m_{\text{rec}}(Z')$ and therefore of low relevance to high- $m_{Z'}$ signal scenarios. Closure tests in a low- p_T^{miss} sideband version of the ABCD method for the QCD estimation in the $\tau_h\tau_h$ channel show agreement within statistical uncertainties, and so no additional closure sys-

tematic uncertainty is applied. Instead, for all three channels, the transfer factors are fit with a first-order polynomial as a function of $m_{\text{rec}}(Z')$, and the deviation of this function from unity provides the $m_{\text{rec}}(Z')$ shape uncertainty. The resulting systematic uncertainties for the three channels are 2.0–150% ($\tau_\mu \tau_h$), 2.0–50% ($\tau_e \tau_h$), and 0.2–24% ($\tau_h \tau_h$), depending on the data-taking year and mass value. The large relative values of the shape systematic uncertainties occur at the high end of the $m_{\text{rec}}(Z')$ distribution, especially for the $\tau_\mu \tau_h$ channel, and arise from the limited statistical precision of the extrapolation. Since the QCD multijet background is a very small fraction of the total background in this $m_{\text{rec}}(Z')$ region, the impact of this systematic uncertainty on the results is small.

9 Results

The signal strength is extracted from an ML fit of the probability density function to the data yields in the SRs for all bins of the $m_{\text{rec}}(Z')$ distribution, for all channels, and all data-taking years. The probability density function is a sum of the corresponding predictions of the yields for the signal and for each of the SM background processes, with the signal strength as a free scale parameter modifying the signal prediction. Statistical uncertainties from the CRs and simulated samples, as well as all systematic uncertainties, are introduced as nuisance parameters, including their correlations. The nuisance parameters are constrained by log-normal penalty factors in the likelihood, which are uncorrelated across the $m_{\text{rec}}(Z')$ bins. Any cross-contamination among the CRs is accounted for by the correlation terms between the CRs. The contamination of CRs by the signal is found to be small (order of percent or less), and is neglected in the fit model. The ML fit is implemented with the CMS statistical analysis tool COMBINE [73].

Table 1 shows for each channel the estimated pre-fit event yield for each SM background contribution, together with the total background and the observed yields. The $m_{\text{rec}}(Z')$ distributions for the data and the SM background estimates as determined by the background-only fits (post-fit) are displayed for the three channels in Fig. 2. The lower panels compare the data with the pre-fit and post-fit estimates. The observed yields are consistent with the SM prediction.

Table 1: Pre-fit estimated background and observed event yields for the three channels in the SR. For each SM process the yield in events is shown, along with the percentile contribution to the total. The background SFs derived from the dedicated CRs are used to determine the listed background yields. The uncertainty associated with the event yields is statistical. The uncertainty in the background contributions adjusted by the SFs includes the uncertainty in the corresponding SF.

Process	$\tau_\mu \tau_h$		$\tau_e \tau_h$		$\tau_h \tau_h$	
	Events	%	Events	%	Events	%
W + jets	851 ± 86	24	717 ± 89	30	158 ± 16	10
DY	1315 ± 70	37	661 ± 34	28	920 ± 43	55
$t\bar{t}$	637 ± 21	18	468.1 ± 8.1	20	32.8 ± 1.6	2
Single t	157.4 ± 6.0	5	116.1 ± 5.4	5	5.5 ± 1.2	<1
VV	302.4 ± 7.7	9	247.5 ± 6.0	11	26.8 ± 1.8	2
QCD multijet	251 ± 44	7	135.0 ± 8.0	6	514 ± 27	31
Total background	3514 ± 122	100	2345 ± 96	100	1658 ± 53	100
Data	3737		2269		1549	

Using the CL_s criterion [74, 75], we compute an asymptotic approximation of the upper limit




Figure 2: Post-fit reconstructed mass distributions for the (upper left) $\tau_\mu\tau_h$, (upper right) $\tau_e\tau_h$, and (lower) $\tau_h\tau_h$ channel. The background contributions appear as stacked solid histograms, while the observed data are shown as black solid markers with error bars. Overflow counts are included in the last bin. The gray shadow band represents the statistical plus systematic uncertainty in the post-fit background prediction. The ratio to the pre-fit background prediction is indicated in magenta markers in the ratio. The signal expectation is shown in lines for $m_{Z'}$ of 500 GeV (red dashed) and 2500 GeV (blue dotted), normalized to 1850 and 0.131 pb, respectively.

at 95% CL [76] on the product of the cross section and branching fraction $\sigma\mathcal{B}(Z' \rightarrow \tau^+\tau^-)$, as a function of $m_{Z'}$. Figure 3 shows the limits for each channel and Figure 4 the combination of the three channels.

Because of the large branching fractions of the tau lepton to hadronic modes, the $\tau_h\tau_h$ channel provides the dominant contribution to these limits across most of the Z' boson mass range. However, as $m_{Z'}$ approaches 4-5 TeV, there is a slight decrease in the τ_h identification efficiency, which occurs because hadronic decays of Lorentz-boosted tau leptons lead to merged tracks and neutral hadrons. In contrast, the identification efficiency for light leptons improves as their energies increase to TeV-scale values. As a result, for $m_{Z'} = 4\text{--}5$ TeV, the acceptance for the light lepton channels grows, compensating for their lower branching fractions, and thus these channels slightly outperform the $\tau_h\tau_h$ channel. There is no significant difference in the impact between the mixed tau pair decay channels. The dashed lines represent the theoretical prediction for the branching fraction values of 1, 3, and 10%, where 3% is the approximate value of $\mathcal{B}(Z' \rightarrow \tau^+\tau^-)$ in the sequential standard model (SSM), considering $t\bar{t}$ decays to be kinematically allowed. The corresponding lower bounds on $m_{Z'}$ are 3.0, 3.5, and 4.1 TeV, respectively.

10 Summary

A search has been performed in proton-proton collisions at $\sqrt{s} = 13$ TeV for the production via quark-antiquark annihilation of a heavy neutral gauge boson Z' decaying to $\tau^+\tau^-$. The data were recorded at the LHC during 2016–2018, and correspond to an integrated luminosity of 138 fb^{-1} . As the tau lepton can decay hadronically (τ_h) or leptonically (τ_ℓ), with ℓ representing a muon or electron, the analysis includes the three decay channels with the highest τ lepton branching fraction products: $Z' \rightarrow \tau_\mu^\pm\tau_h^\mp$, $Z' \rightarrow \tau_e^\pm\tau_h^\mp$, and $Z' \rightarrow \tau_h^+\tau_h^-$. An estimator $m_{\text{rec}}(Z')$ is constructed to approximate the Z' boson mass $m_{Z'}$ in the presence of undetected neutrino daughters from the τ lepton decays.

The main background processes are estimated using Monte Carlo simulation adjusted by scale factors derived from data. The observed yields are found to be consistent with the background prediction. A shape-based analysis is performed using the $m_{\text{rec}}(Z')$ distribution as the fit discriminant to determine the likelihood for the observed data in the presence of a signal and the predicted background contributions. The upper limit on the product of the Z' boson production cross section and decay branching fraction is set at 95% confidence level as a function of $m_{Z'}$.

In the scenario considered in this analysis, the data excludes a Z' boson with $m_{Z'}$ less than 3.0, 3.5, or 4.1 TeV for a branching fraction of 1, 3, or 10%, respectively. These exclusion limits are the most stringent to date for a Z' boson decaying to $\tau^+\tau^-$.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centres and personnel of the Worldwide LHC Computing Grid and other centres for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided

by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie programme and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the "Excellence of Science – EOS" – be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy – EXC 2121 "Quantum Universe" – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2021-4.1.2-NEMZ_KI (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Centre for High Performance Computing, Big Data and Quantum Computing and FAIR – Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF "a way of making Europe", and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G670016 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

References

- [1] CMS Collaboration, "Search for resonant and nonresonant new phenomena in high-mass

- dilepton final states at $\sqrt{s} = 13$ TeV", *JHEP* **07** (2021) 208, doi:10.1007/JHEP07(2021)208, arXiv:2103.02708.
- [2] P. K. Mohapatra, R. N. Mohapatra, and P. B. Pal, "Implications of E_6 grand unification", *Phys. Rev. D* **33** (1986) 2010, doi:10.1103/PhysRevD.33.2010.
- [3] D. London and J. L. Rosner, "Extra gauge bosons in E_6 ", *Phys. Rev. D* **34** (1986) 1530, doi:10.1103/PhysRevD.34.1530.
- [4] L. Randall and R. Sundrum, "Large mass hierarchy from a small extra dimension", *Phys. Rev. Lett.* **83** (1999) 3370, doi:10.1103/PhysRevLett.83.3370.
- [5] ATLAS Collaboration, "Search for high-mass dilepton resonances using 139 fb $^{-1}$ of pp collision data collected at $\sqrt{s} = 13$ TeV with the ATLAS detector", *Phys. Lett. B* **796** (2019) 68, doi:10.1016/j.physletb.2019.07.016, arXiv:1903.06248.
- [6] P. Langacker, "The physics of heavy Z' gauge bosons", *Rev. Mod. Phys.* **81** (2009) 1199, doi:10.1103/RevModPhys.81.1199, arXiv:0801.1345.
- [7] K. R. Lynch, E. H. Simmons, M. Narain, and S. Mrenna, "Finding Z' bosons coupled preferentially to the third family at LEP and the Tevatron", *Phys. Rev. D* **63** (2001) 035006, doi:10.1103/PhysRevD.63.035006, arXiv:hep-ph/0007286.
- [8] ATLAS Collaboration, "Search for additional heavy neutral Higgs and gauge bosons in the ditau final state produced in 36 fb $^{-1}$ of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector", *JHEP* **01** (2018) 055, doi:10.1007/JHEP01(2018)055, arXiv:1709.07242.
- [9] CMS Collaboration, "Search for heavy resonances decaying to tau lepton pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV", *JHEP* **02** (2017) 048, doi:10.1007/JHEP02(2017)048, arXiv:1611.06594.
- [10] ATLAS Collaboration, "Search for new resonances in mass distributions of jet pairs using 139 fb $^{-1}$ of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector", *JHEP* **03** (2020) 145, doi:10.1007/JHEP03(2020)145, arXiv:1910.08447.
- [11] CMS Collaboration, "Search for narrow resonances in the b-tagged dijet mass spectrum in proton-proton collisions at $\sqrt{s} = 13$ TeV", *Phys. Rev. D* **108** (2023) 012009, doi:10.1103/PhysRevD.108.012009, arXiv:2205.01835.
- [12] BaBar Collaboration, "Evidence for an excess of $\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau$ decays", *Phys. Rev. Lett.* **109** (2012) 101802, doi:10.1103/PhysRevLett.109.101802, arXiv:1205.5442.
- [13] BaBar Collaboration, "Measurement of an excess of $\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau$ decays and implications for charged Higgs bosons", *Phys. Rev. D* **88** (2013) 072012, doi:10.1103/PhysRevD.88.072012, arXiv:1303.0571.
- [14] Belle Collaboration, "Measurement of the branching ratio of $\bar{B} \rightarrow D^{(*)} \tau^- \bar{\nu}_\tau$ relative to $\bar{B} \rightarrow D^{(*)} \ell^- \bar{\nu}_\ell$ decays with hadronic tagging at Belle", *Phys. Rev. D* **92** (2015) 072014, doi:10.1103/PhysRevD.92.072014, arXiv:1507.03233.
- [15] LHCb Collaboration, "Measurement of the ratio of branching fractions $\mathcal{B}(\bar{B}^0 \rightarrow D^{*+} \tau^- \bar{\nu}_\tau) / \mathcal{B}(\bar{B}^0 \rightarrow D^{*+} \mu^- \bar{\nu}_\mu)$ ", *Phys. Rev. Lett.* **115** (2015) 111803, doi:10.1103/PhysRevLett.115.111803, arXiv:1506.08614. [Erratum: doi:10.1103/PhysRevLett.115.159901].

-
- [16] Belle Collaboration, “Measurement of the branching ratio of $\bar{B}^0 \rightarrow D^{*+}\tau^-\bar{\nu}_\tau$ relative to $\bar{B}^0 \rightarrow D^{*+}\ell^-\bar{\nu}_\ell$ decays with a semileptonic tagging method”, *Phys. Rev. D* **94** (2016) 072007, doi:10.1103/PhysRevD.94.072007, arXiv:1607.07923.
 - [17] Belle Collaboration, “Measurement of the τ lepton polarization and $R(D^*)$ in the decay $\bar{B} \rightarrow D^*\tau^-\bar{\nu}_\tau$ ”, *Phys. Rev. Lett.* **118** (2017) 211801, doi:10.1103/PhysRevLett.118.211801, arXiv:1612.00529.
 - [18] Belle Collaboration, “Measurement of the τ lepton polarization and $R(D^*)$ in the decay $\bar{B} \rightarrow D^*\tau^-\bar{\nu}_\tau$ with one-prong hadronic τ decays at Belle”, *Phys. Rev. D* **97** (2018) 012004, doi:10.1103/PhysRevD.97.012004, arXiv:1709.00129.
 - [19] LHCb Collaboration, “Measurement of the ratio of the $B^0 \rightarrow D^{*-}\tau^+\nu_\tau$ and $B^0 \rightarrow D^{*-}\mu^+\nu_\mu$ branching fractions using three-prong τ -lepton decays”, *Phys. Rev. Lett.* **120** (2018) 171802, doi:10.1103/PhysRevLett.120.171802, arXiv:1708.08856.
 - [20] LHCb Collaboration, “Test of lepton flavor universality by the measurement of the $B^0 \rightarrow D^{*-}\tau^+\nu_\tau$ branching fraction using three-prong τ decays”, *Phys. Rev. D* **97** (2018) 072013, doi:10.1103/PhysRevD.97.072013, arXiv:1711.02505.
 - [21] LHCb Collaboration, “Measurement of the ratios of branching fractions $\mathcal{R}(D^*)$ and $\mathcal{R}(D^0)$ ”, *Phys. Rev. Lett.* **131** (2023) 111802, doi:10.1103/PhysRevLett.131.111802, arXiv:2302.02886.
 - [22] LHCb Collaboration, “Measurement of the branching fraction ratios $R(D^+)$ and $R(D^{*+})$ using muonic τ decays”, *Phys. Rev. Lett.* **134** (2025) 061801, doi:10.1103/PhysRevLett.134.061801, arXiv:2406.03387.
 - [23] R. Diener, S. Godfrey, and T. A. W. Martin, “Unravelling an extra neutral gauge boson at the LHC using third generation fermions”, *Phys. Rev. D* **83** (2011) 115008, doi:10.1103/PhysRevD.83.115008, arXiv:1006.2845.
 - [24] B. C. Allanach, J. M. Butterworth, and T. Corbett, “Collider constraints on Z' models for neutral current B anomalies”, *JHEP* **08** (2019) 106, doi:10.1007/JHEP08(2019)106, arXiv:1904.10954.
 - [25] A. Flórez et al., “Searching for new heavy neutral gauge bosons using vector boson fusion processes at the LHC”, *Phys. Lett. B* **767** (2017) 126, doi:10.1016/j.physletb.2017.01.062, arXiv:1609.09765.
 - [26] HEPData record for this analysis, 2024. doi:10.17182/hepdata.155459.
 - [27] CMS Collaboration, “The CMS Experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
 - [28] CMS Collaboration, “Development of the CMS detector for the CERN LHC Run 3”, *JINST* **19** (2024) P05064, doi:10.1088/1748-0221/19/05/P05064, arXiv:2309.05466.
 - [29] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
 - [30] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.

- [31] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965.
- [32] CMS Collaboration, “Technical proposal for the Phase-II upgrade of the Compact Muon Solenoid”, CMS Technical Proposal CERN-LHCC-2015-010, CMS-TDR-15-02, 2015.
- [33] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- k_T jet clustering algorithm”, *JHEP* **04** (2008) 063, doi:10.1088/1126-6708/2008/04/063, arXiv:0802.1189.
- [34] M. Cacciari, G. P. Salam, and G. Soyez, “FastJet user manual”, *Eur. Phys. J. C* **72** (2012) 1896, doi:10.1140/epjc/s10052-012-1896-2, arXiv:1111.6097.
- [35] CMS Collaboration, “Pileup mitigation at CMS in 13 TeV data”, *JINST* **15** (2020) P09018, doi:10.1088/1748-0221/15/09/P09018, arXiv:2003.00503.
- [36] CMS Collaboration, “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV”, *JINST* **12** (2017) P02014, doi:10.1088/1748-0221/12/02/P02014, arXiv:1607.03663.
- [37] CMS Collaboration, “Performance of missing transverse momentum reconstruction in proton-proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector”, *JINST* **14** (2019) P07004, doi:10.1088/1748-0221/14/07/P07004, arXiv:1903.06078.
- [38] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [39] CMS Collaboration, “Reconstruction of signal amplitudes in the CMS electromagnetic calorimeter in the presence of overlapping proton-proton interactions”, *JINST* **15** (2020) P10002, doi:10.1088/1748-0221/15/10/P10002, arXiv:2006.14359.
- [40] CMS Collaboration, “Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC”, *JINST* **16** (2021) P05014, doi:10.1088/1748-0221/16/05/P05014, arXiv:2012.06888.
- [41] CMS Collaboration, “Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P10005, doi:10.1088/1748-0221/13/10/P10005, arXiv:1809.02816.
- [42] CMS Collaboration, “Identification of hadronic tau lepton decays using a deep neural network”, *JINST* **17** (2022) P07023, doi:10.1088/1748-0221/17/07/P07023, arXiv:2201.08458.
- [43] CMS Collaboration, “Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV”, *JINST* **13** (2018) P05011, doi:10.1088/1748-0221/13/05/P05011, arXiv:1712.07158.
- [44] J. Alwall et al., “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, *JHEP* **07** (2014) 079, doi:10.1007/JHEP07(2014)079, arXiv:1405.0301.
- [45] P. Nason, “A new method for combining NLO QCD with shower Monte Carlo algorithms”, *JHEP* **11** (2004) 040, doi:10.1088/1126-6708/2004/11/040, arXiv:hep-ph/0409146.

-
- [46] S. Frixione, P. Nason, and C. Oleari, “Matching NLO QCD computations with parton shower simulations: the POWHEG method”, *JHEP* **11** (2007) 070, doi:10.1088/1126-6708/2007/11/070, arXiv:0709.2092.
- [47] S. Alioli, P. Nason, C. Oleari, and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX”, *JHEP* **06** (2010) 043, doi:10.1007/JHEP06(2010)043, arXiv:1002.2581.
- [48] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [49] B. Fuks and R. Ruiz, “A comprehensive framework for studying W' and Z' bosons at hadron colliders with automated jet veto resummation”, *JHEP* **05** (2017) 032, doi:10.1007/JHEP05(2017)032, arXiv:1701.05263.
- [50] CMS Collaboration, “Event generator tunes obtained from underlying event and multiparton scattering measurements”, *Eur. Phys. J. C* **76** (2016) 155, doi:10.1140/epjc/s10052-016-3988-x, arXiv:1512.00815.
- [51] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) 4, doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.
- [52] NNPDF Collaboration, “Parton distributions for the LHC run II”, *JHEP* **04** (2015) 040, doi:10.1007/JHEP04(2015)040, arXiv:1410.8849.
- [53] S. Alioli, P. Nason, C. Oleari, and E. Re, “NLO single-top production matched with shower in POWHEG: s - and t -channel contributions”, *JHEP* **09** (2009) 111, doi:10.1088/1126-6708/2009/09/111, arXiv:0907.4076. [Erratum: doi:10.1007/JHEP02(2010)011].
- [54] E. Re, “Single-top Wt -channel production matched with parton showers using the POWHEG method”, *Eur. Phys. J. C* **71** (2011) 1547, doi:10.1140/epjc/s10052-011-1547-z, arXiv:1009.2450.
- [55] R. Gavin, Y. Li, F. Petriello, and S. Quackenbush, “FEWZ 2.0: A code for hadronic Z production at next-to-next-to-leading order”, *Comput. Phys. Commun.* **182** (2011) 2388, doi:10.1016/j.cpc.2011.06.008, arXiv:1011.3540.
- [56] T. Melia, P. Nason, R. Rontsch, and G. Zanderighi, “ W^+W^- , WZ and ZZ production in the POWHEG BOX”, *JHEP* **11** (2011) 078, doi:10.1007/JHEP11(2011)078, arXiv:1107.5051.
- [57] M. Beneke, P. Falgari, S. Klein, and C. Schwinn, “Hadronic top-quark pair production with NNLL threshold resummation”, *Nucl. Phys. B* **855** (2012) 695, doi:10.1016/j.nuclphysb.2011.10.021, arXiv:1109.1536.
- [58] M. Cacciari et al., “Top-pair production at hadron colliders with next-to-next-to-leading logarithmic soft-gluon resummation”, *Phys. Lett. B* **710** (2012) 612, doi:10.1016/j.physletb.2012.03.013, arXiv:1111.5869.
- [59] P. Bärnreuther, M. Czakon, and A. Mitov, “Percent level precision physics at the Tevatron: first genuine NNLO QCD corrections to $q\bar{q} \rightarrow t\bar{t} + X$ ”, *Phys. Rev. Lett.* **109** (2012) 132001, doi:10.1103/PhysRevLett.109.132001, arXiv:1204.5201.

- [60] M. Czakon and A. Mitov, “NNLO corrections to top-pair production at hadron colliders: the all-fermionic scattering channels”, *JHEP* **12** (2012) 054, doi:10.1007/JHEP12(2012)054, arXiv:1207.0236.
- [61] M. Czakon and A. Mitov, “NNLO corrections to top pair production at hadron colliders: the quark-gluon reaction”, *JHEP* **01** (2013) 080, doi:10.1007/JHEP01(2013)080, arXiv:1210.6832.
- [62] R. Gavin, Y. Li, F. Petriello, and S. Quackenbush, “W physics at the LHC with FEWZ 2.1”, *Comput. Phys. Commun.* **184** (2013) 208, doi:10.1016/j.cpc.2012.09.005, arXiv:1201.5896.
- [63] M. Czakon, P. Fiedler, and A. Mitov, “Total top-quark pair-production cross section at hadron colliders through $O(\alpha_s^4)$ ”, *Phys. Rev. Lett.* **110** (2013) 252004, doi:10.1103/PhysRevLett.110.252004, arXiv:1303.6254.
- [64] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [65] CMS Collaboration, “Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS”, *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [66] CMS Collaboration, “CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-17-004, 2018.
- [67] CMS Collaboration, “CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-18-002, 2019.
- [68] CMS Collaboration, “Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV”, *JHEP* **07** (2018) 161, doi:10.1007/JHEP07(2018)161, arXiv:1802.02613.
- [69] ATLAS Collaboration, “Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV with the ATLAS detector at the LHC”, *Phys. Rev. Lett.* **117** (2016) 182002, doi:10.1103/PhysRevLett.117.182002, arXiv:1606.02625.
- [70] CMS Collaboration, “ECAL 2016 refined calibration and Run2 summary plots”, CMS Detector Performance Summary CMS-DP-2020-021, 2020.
- [71] J. Butterworth et al., “PDF4LHC recommendations for LHC Run II”, *J. Phys. G* **43** (2016) 023001, doi:10.1088/0954-3899/43/2/023001, arXiv:1510.03865.
- [72] M. Cacciari et al., “The $t\bar{t}$ cross-section at 1.8 and 1.96 TeV: a study of the systematics due to parton densities and scale dependence”, *JHEP* **04** (2004) 068, doi:10.1088/1126-6708/2004/04/068, arXiv:hep-ph/0303085.
- [73] CMS Collaboration, “The CMS statistical analysis and combination tool: COMBINE”, *Comput. Softw. Big Sci.* **8** (2024) 19, doi:10.1007/s41781-024-00121-4, arXiv:2404.06614.
- [74] T. Junk, “Confidence level computation for combining searches with small statistics”, *Nucl. Instrum. Meth. A* **434** (1999) 435, doi:10.1016/S0168-9002(99)00498-2, arXiv:hep-ex/9902006.

- [75] A. L. Read, “Presentation of search results: the CL_s technique”, *J. Phys. G* **28** (2002) 2693, doi:10.1088/0954-3899/28/10/313.
- [76] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, “Asymptotic formulae for likelihood-based tests of new physics”, *Eur. Phys. J. C* **71** (2011) 1554, doi:10.1140/epjc/s10052-011-1554-0, arXiv:1007.1727. [Erratum: doi:10.1140/epjc/s10052-013-2501-z].




Figure 3: Upper limits at 95% CL on $\sigma B(Z' \rightarrow \tau^+\tau^-)$ for the (upper) $\tau_\mu \tau_h$, (middle) $\tau_e \tau_h$, and (lower) $\tau_h \tau_h$ channel. In each plot, the solid black line represents the observed limit, and the black dashed line with green and yellow bands depicts the expected limit with its one- and two-standard deviation uncertainties, respectively. The red dashed, magenta dotted, and blue dash-dotted lines represent the theoretical predictions for $B(Z' \rightarrow \tau^+\tau^-) = 1, 3, \text{ and } 10\%$, respectively; 3% is the value of $B(Z' \rightarrow \tau^+\tau^-)$ in the SSM. These assume that the $B(Z' \rightarrow \tau^+\tau^-)$ values are fixed for all $m_{Z'}$.




Figure 4: Upper limits at 95% CL on $\sigma\mathcal{B}(Z' \rightarrow \tau^+\tau^-)$ for the combination of all channels. The solid black line represents the observed limit, and the black dashed line with green and yellow bands depicts the expected limit with its one- and two-standard deviation uncertainties, respectively. The red dashed, magenta dotted, and blue dash-dotted lines represent the theoretical predictions for $\mathcal{B}(Z' \rightarrow \tau^+\tau^-) = 1, 3$, and 10% , respectively; 3% is the value of $\mathcal{B}(Z' \rightarrow \tau^+\tau^-)$ in the SSM. These assume that the $\mathcal{B}(Z' \rightarrow \tau^+\tau^-)$ values are fixed for all $m_{Z'}$.

A The CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan¹ 

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, L. Benato , T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic , P.S. Hussain , M. Jeitler² , N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck² , R. Schöfbeck² , D. Schwarz , M. Sonawane , W. Waltenberger , C.-E. Wulz² 

Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , T. Van Laer, P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, S. Lowette , I. Makarenko , D. Müller , S. Tavernier , M. Tytgat³ , G.P. Van Onsem , S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios , J. Jaramillo , A. Khalilzadeh, F.A. Khan , K. Lee , M. Mahdavihorrami , A. Malara , S. Paredes , M.A. Shahzad, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , Y. Hong , J. Knolle , L. Lambrecht , D. Marckx , K. Mota Amarilo , K. Skovpen , N. Van Den Bossche , J. van der Linden , L. Wezenbeek 

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , A. Bethani , G. Bruno , C. Caputo , J. De Favereau De Jeneret , C. Delaere , I.S. Donertas , A. Giammanco , A.O. Guzel , Sa. Jain , V. Lemaitre, J. Lidrych , P. Mastrapasqua , T.T. Tran 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , G. Correia Silva , C. Hensel , T. Menezes De Oliveira , C. Mora Herrera⁴ , P. Rebello Teles , M. Soeiro, A. Vilela Pereira⁴ 

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁵, E.M. Da Costa , G.G. Da Silveira⁶ , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza, T. Laux Kuhn , M. Macedo , J. Martins⁷ , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁶ , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula 

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 

University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , D. Laroze , S. Thakur 

Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu

Institute of High Energy Physics, Beijing, China

G.M. Chen⁸ , H.S. Chen⁸ , M. Chen⁸ , F. Iemmi , C.H. Jiang, A. Kapoor⁹ , H. Liao , Z.-A. Liu¹⁰ , R. Sharma¹¹ , J.N. Song¹⁰, J. Tao , C. Wang⁸, J. Wang , Z. Wang⁸, H. Zhang , J. Zhao 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, X. Sun , D. Wang , H. Yang, L. Zhang , Y. Zhao, C. Zhou 

Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 

Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer¹², B. Li¹³, K. Yi¹⁴ , J. Zhang 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

Y. Li

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao 

Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

F. Ramirez , C. Rendón , M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac 

University of Split, Faculty of Science, Split, Croatia

M. Kovac , A. Petkovic , T. Sculac 

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov¹⁵ , T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , J. Mousa , C. Nicolaou, L. Paizanos, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

Charles University, Prague, Czech Republic

M. Finger , M. Finger Jr. , A. Kveton 

Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

B. El-mahdy , S. Khalil¹⁶ , E. Salama^{17,18} 

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

M. Abdullah Al-Mashad , M.A. Mahmoud 

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht , M. Kadastik, T. Lange , S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken 

Department of Physics, University of Helsinki, Helsinki, Finland

H. Kirschenmann , K. Osterberg , M. Voutilainen 

Helsinki Institute of Physics, Helsinki, Finland

S. Bharthuar , N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew , K.T.S. Kallonen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Myllymäki , M.m. Rantanen , H. Siikonen , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka , H. Petrow 

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi , L. Portales , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro¹⁹ , P. Simkina , M. Titov , M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

F. Beaudette , G. Boldrini , P. Busson , A. Cappati , C. Charlot , M. Chiusi , T.D. Cuisset , F. Damas , O. Davignon , A. De Wit , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , M. Nguyen , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , L. Urda Gómez , E. Vernazza , A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram²⁰ , J. Andrea , D. Apparu , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet , G. Saha , M.A. Sessini , P. Van Hove , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo , P. Depasse , C. Dozen²¹ , H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg , G. Grenier , B. Ille , E. Jourd'huy, I.B. Laktineh, M. Lethuillier , L. Mirabito, S. Perries, A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao 

Georgian Technical University, Tbilisi, Georgia

G. Adamov, I. Lomidze , Z. Tsamalaidze¹⁵ 

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser , A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann , M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi , A. Schmidt , C. Seth, A. Sharma , J.L. Spah , A. Stein , F. Torres Da Silva De Araujo²² , S. Wiedenbeck , S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman²³ , K. Borras²⁴ , A. Campbell , A. Cardini , C. Cheng , F. Colombina , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , O. Filatov , E. Gallo²³ , A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , B. Kaech , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka²⁵ , W. Lohmann²⁶ , F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , Y. Otariid, J. Park , E. Ranken , A. Raspereza , D. Rastorguev , J. Rübenach, L. Rygaard, A. Saggio , M. Scham^{27,24} , S. Schnake²⁴ , P. Schütze , C. Schwanenberger²³ , D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford , F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , Y. Wen , K. Wichmann, L. Wiens²⁴ , C. Wissing , Y. Yang , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , S. Bollweg, M. Bonanomi , P. Connor , K. El Morabit , Y. Fischer , E. Garutti , A. Grohsjean , J. Haller , D. Hundhausen, H.R. Jabusch , G. Kasieczka , P. Keicher , R. Klanner , W. Korcari , T. Kramer , C.c. Kuo, V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoiiu , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, B. Wiederspan, M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , E. Butz , T. Chwalek , A. Dierlamm , A. Droll, U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann²⁸ , R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , R. Koppenhöfer , O. Lavoryk , J.M. Lawhorn , M. Link, A. Lintuluoto , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa, L. Stockmeier, K. Tauqeer, M. Toms , N. Trevisani , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis , A. Kyriakis , A. Papadopoulos²⁸, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

P. Kontaxakis , G. Melachroinos, Z. Painesis , I. Papavergou , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 

National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsipolitis , A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece

I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

C. Hajdu , D. Horvath^{29,30} , K. Márton, A.J. Rádl³¹ , F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkuti³² , M.M.A. Gadallah³³ , Á. Kadlecik , P. Major , G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 

HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo³² , T. Novak 

Panjab University, Chandigarh, India

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra³⁴ , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand, J.B. Singh , A. Singla 

University of Delhi, Delhi, India

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman³⁵ , S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

Indian Institute of Technology Madras, Madras, India

M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , G. Dash , P. Jana 

P. Kalbhor , S. Kamble , J.R. Komaragiri³⁶ , D. Kumar³⁶ , T. Mishra , B. Parida , P.R. Pujahari , N.R. Saha , A. Sharma , A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay 

Tata Institute of Fundamental Research-A, Mumbai, India

S. Dugad, G.B. Mohanty , M. Shelake, P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, S. Kumar , G. Majumder , K. Mazumdar , S. Parolia , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati³⁷ , C. Kar , D. Maity³⁸ , P. Mal , V.K. Muraleedharan Nair Bindhu³⁸ , K. Naskar³⁸ , A. Nayak³⁸ , S. Nayak, K. Pal , P. Sadangi, S.K. Swain , S. Varghese³⁸ , D. Vats³⁸ 

Indian Institute of Science Education and Research (IISER), Pune, India

S. Acharya³⁹ , A. Alpina , S. Dube , B. Gomber³⁹ , P. Hazarika , B. Kansal , A. Laha , B. Sahu³⁹ , S. Sharma , K.Y. Vaish 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi⁴⁰ , A. Jafari⁴¹ , M. Zeinali⁴² 

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Bashiri, S. Chenarani⁴³ , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie , M. Mohammadi Najafabadi , S. Tizchang⁴⁴ 

University College Dublin, Dublin, Ireland

M. Felcini , M. Grunewald 

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia^{a,b} , A. Colaleo^{a,b} , D. Creanza^{a,c} , B. D'Anzi^{a,b} , N. De Filippis^{a,c} , M. De Palma^{a,b} , W. Elmetenawee^{a,b,45} , N. Ferrara^{a,b} , L. Fiore^a , G. Iaselli^{a,c} , L. Longo^a , M. Louka^{a,b} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^a , V. Mastrapasqua^{a,b} , S. My^{a,b} , S. Nuzzo^{a,b} , A. Pellecchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} , R. Radogna^{a,b} , D. Ramos^a , A. Ranieri^a , L. Silvestris^a , F.M. Simone^{a,c} , Ü. Sözbilir^a , A. Stamerra^{a,b} , D. Troiano^{a,b} , R. Venditti^{a,b} , P. Verwilligen^a , A. Zaza^{a,b} 

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , P. Capiluppi^{a,b} , A. Castro^{+a,b} , F.R. Cavallo^a , M. Cuffiani^{a,b} , G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} , D. Fasanella^a , P. Giacomelli^a , L. Giommi^{a,b} , C. Grandi^a , L. Guiducci^{a,b} , S. Lo Meo^{a,46} , M. Lorusso^{a,b} , L. Lunerti^a , S. Marcellini^a , G. Masetti^a , F.L. Navarria^{a,b} , G. Paggi^{a,b} , A. Perrotta^a , F. Primavera^{a,b} , A.M. Rossi^{a,b} , S. Rossi Tisbeni^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b} 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,47} , A. Di Mattia^a , A. Lapertosa^a , R. Potenza^{a,b} , A. Tricomi^{a,b,47} , C. Tuve^{a,b} 

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

P. Assiouras^a , G. Barbagli^a , G. Bardelli^{a,b} , B. Camaiani^{a,b} , A. Cassese^a , R. Ceccarelli^a , V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , E. Focardi^{a,b} 

T. Kello^a , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^a , M. Meschini^a , S. Paoletti^a , A. Papanastassiou^{a,b} , G. Sguazzoni^a , L. Viliani^a 

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁴⁸ , D. Piccolo 

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

M. Alves Gallo Pereira^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} 

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a , F. Brivio^a , F. Cetorelli^{a,b} , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , R. Gerosa^{a,b} , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^a , M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , S. Palluotto^{a,b} , D. Pedrini^a , A. Perego^{a,b} , B.S. Pinolini^a , G. Pizzati^{a,b} , S. Ragazzi^{a,b} , T. Tabarelli de Fatis^{a,b}

INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo^a , A. Cagnotta^{a,b} , F. Carnevali^{a,b} , N. Cavallo^{a,c} , F. Fabozzi^{a,c} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,49} , P. Paolucci^{a,28} , B. Rossi^a

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino^a , P. Azzi^a , N. Bacchetta^{a,50} , M. Bellato^a , D. Bisello^{a,b} , P. Bortignon^a , G. Bortolato^{a,b} , A. Bragagnolo^{a,b} , A.C.M. Bulla^a , P. Checchia^a , T. Dorigo^{a,51} , F. Gasparini^{a,b} , U. Gasparini^{a,b} , S. Giorgetti^a , E. Lusiani^a , M. Margoni^{a,b} , A.T. Meneguzzo^{a,b} , M. Migliorini^{a,b} , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , M. Tosi^{a,b} , A. Triossi^{a,b} , S. Ventura^a , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} , G. Zumerle^{a,b}

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

A. Braghieri^a , S. Calzaferri^a , D. Fiorina^a , P. Montagna^{a,b} , V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b}

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal^{a,b} , M.E. Ascioti^{a,b} , G.M. Bilei^a , C. Carrivale^{a,b} , D. Ciangottini^{a,b} , L. Fanò^{a,b} , M. Magherini^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,52} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b}

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

C. Aimè^a , C.A. Alexe^{a,c} , P. Asenov^{a,b} , P. Azzurri^a , G. Bagliesi^a , R. Bhattacharya^a , L. Bianchini^{a,b} , T. Boccali^a , E. Bossini^a , D. Bruschini^{a,c} , R. Castaldi^a , M.A. Ciocci^{a,b} , M. Cipriani^{a,b} , V. D'Amante^{a,d} , R. Dell'Orso^a , S. Donato^a , A. Giassi^a , F. Ligabue^{a,c} , A.C. Marini^a , D. Matos Figueiredo^a , A. Messineo^{a,b} , S. Mishra^a , M. Musich^{a,b} , F. Palla^a , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^a , T. Sarkar^a , A. Scribano^a , P. Spagnolo^a , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , F. Vaselli^{a,c} , A. Venturi^a , P.G. Verdini^a

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

C. Baldenegro Barrera^{a,b} , P. Barria^a , C. Basile^{a,b} , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , D. Del Re^{a,b} , E. Di Marco^{a,b} , M. Diemoz^a , F. Errico^{a,b} , R. Gargiulo^{a,b} , E. Longo^{a,b} , L. Martikainen^{a,b} , J. Mijuskovic^{a,b} , G. Organtini^{a,b} , F. Pandolfi^a , R. Paramatti^{a,b} , C. Quaranta^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a

F. Santanastasio^{a,b} , L. Soffi^a , V. Vladimirov^{a,b}

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^a , R. Bellan^{a,b} , A. Bellora^{a,b} , C. Biino^a , C. Borca^{a,b} , N. Cartiglia^a , M. Costa^{a,b} , R. Covarelli^{a,b} , N. Demaria^a , L. Finco^a , M. Grippo^{a,b} , B. Kiani^{a,b} , F. Legger^a , F. Luongo^{a,b} , C. Mariotti^a , L. Markovic^{a,b} , S. Maselli^a , A. Mecca^{a,b} , L. Menzio^{a,b} , P. Meridiani^a , E. Migliore^{a,b} , M. Monteno^a , R. Mulargia^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , M. Pelliccioni^a , M. Ruspa^{a,c} , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} , A. Staiano^a , C. Tarricone^{a,b} , D. Trocino^a , G. Umoret^{a,b} , R. White^{a,b} 

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar^{a,b} , S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , K. De Leo^a , G. Della Ricca^{a,b} 

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , B. Kim , J. Kim, D. Lee, H. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , M.S. Ryu , S. Sekmen , B. Tae, Y.C. Yang 

Department of Mathematics and Physics - GWNu, Gangneung, Korea

M.S. Kim 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon 

Hanyang University, Seoul, Korea

E. Asilar , J. Choi⁵³ , D. Kim , T.J. Kim , J.A. Merlin, Y. Ryou

Korea University, Seoul, Korea

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , S. Lee , J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh , S. Yang 

Sejong University, Seoul, Korea

H. S. Kim , Y. Kim, S. Lee

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , Y.W. Kim , S. Ko , H. Kwon , H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, Y. Kang , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea

S. Ha , K. Hwang , H.D. Yoo 

Sungkyunkwan University, Suwon, Korea

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy , Y. Gharbia 

Kuwait University - College of Science - Department of Physics, Safat, Kuwait

F. Alazemi 

Riga Technical University, Riga, Latvia

K. Dreimanis , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs, A. Potrebko ,
M. Seidel , D. Sidiropoulos Kontos 

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Yusuff⁵⁴ , Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,
M. León Coello , J.A. Murillo Quijada , A. Sehwat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, E. De La Cruz-Burelo , I. Heredia-
De La Cruz⁵⁵ , R. Lopez-Fernandez , J. Mejia Guisao , C.A. Mondragon Herrera,
A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

University of Montenegro, Podgorica, Montenegro

I. Bujanja , N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan 

AGH University of Krakow, Krakow, Poland

V. Avati, L. Grzanka , M. Malawski 

National Centre for Nuclear Research, Swierk, Poland

H. Bialkowska , M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski ,
A. Muhammad 

Warsaw University of Technology, Warsaw, Poland

P. Fokow , K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo ,
T. Camporesi , G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo 

G.B. Marozzo , T. Niknejad , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff 

Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , P. Milenovic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

D. Devetak , M. Dordevic , J. Milosevic , L. Nadderd , V. Rekovic

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. Llorente Merino , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , S. Sánchez Navas , J. Sastre , J. Vazquez Escobar 

Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , P. Leguina , E. Palencia Cortezon , J. Prado Pico , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Bhowmik , S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasasoa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

B. Kailasapathy⁵⁶ , D.D.C. Wickramarathna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna⁵⁷ , K. Liyanage , N. Perera 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler , D. Barney , A. Bermúdez Martínez , M. Bianco , A.A. Bin Anuar , A. Bocci , L. Borbonovi , C. Botta , E. Brondolin , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , G. Franzoni , W. Funk , S. Giani , D. Gigi , K. Gill , F. Glege , J. Hegeman , J.K. Heikkilä , B. Huber , V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban²⁶ , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , L. Malgeri , M. Mannelli , M. Matthewman , A. Mehta , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Monti , F. Moortgat , M. Mulders , I. Neutelings , S. Orfanelli , F. Pantaleo , G. Petrucciani , A. Pfeiffer , M. Pierini , H. Qu , D. Rabady , B. Ribeiro Lopes , F. Riti , M. Rovere , H. Sakulin , S. Sanchez Cruz , S. Scarfi , C. Schwick , M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas⁵⁸ , A.G. Stahl Leitner , A. Steen , S. Summers , D. Treille , P. Tropea , D. Walter , J. Wanczyk⁵⁹ , J. Wang , K.A. Wozniak⁶⁰ , S. Wuchterl , P. Zehetner , P. Zejdl 

W.D. Zeuner

PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T. Bevilacqua⁶¹ , L. Caminada⁶¹ , A. Ebrahimi , W. Erdmann , R. Horisberger ,
Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli⁶¹ , L. Noehte⁶¹ ,
T. Rohe , A. Samalan

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , M. Backhaus , G. Bonomelli , A. Calandri , C. Cazzaniga ,
K. Datta , P. De Bryas Dexmiers D'archiac⁵⁹ , A. De Cosa , G. Dissertori , M. Dittmar,
M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer ,
T.G. Harte, D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani ,
L. Marchese , C. Martin Perez , A. Mascellani⁵⁹ , F. Nessi-Tedaldi , F. Pauss ,
V. Perovic , S. Pigazzini , B. Ristic , R. Seidita , J. Steggemann⁵⁹ , A. Tarabini ,
D. Valsecchi , R. Wallny 

Universität Zürich, Zurich, Switzerland

C. Amsler⁶² , P. Bärttschi , M.F. Canelli , K. Cormier , M. Huwiler , W. Jin ,
A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechti , A. Macchiolo , P. Meiring ,
F. Meng , U. Molinatti , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr,
F. Stäger , R. Tramontano 

National Central University, Chung-Li, Taiwan

C. Adloff⁶³, D. Bhowmik, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari³⁶ 

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao,
S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker ,
L.s. Tsai, D. Tsionou, H.y. Wu, E. Yazgan 

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu⁶⁴ , E. Eskut , Y. Guler⁶⁵ ,
E. Gurpınar Guler⁶⁵ , C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , Y. Komurcu ,
G. Onengut , K. Ozdemir⁶⁶ , A. Polatoz , B. Tali⁶⁷ , U.G. Tok , S. Turkcapar ,
E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

G. Sokmen, M. Yalvac⁶⁸ 

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya⁶⁹ , O. Kaya⁷⁰ , S. Tekten⁷¹ 

Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak^{64,72} , G.G. Dincer⁶⁴ , S. Sen⁷³ 

Istanbul University, Istanbul, Turkey

O. Aydılek⁷⁴ , B. Hacisahinoglu , I. Hos⁷⁵ , B. Kaynak , S. Ozkorucuklu , O. Potok ,
H. Sert , C. Simsek , C. Zorbilmez 

Yildiz Technical University, Istanbul, Turkey

S. Cerci , B. Isildak⁷⁶ , D. Sunar Cerci , T. Yetkin 

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , B. Grynyov 

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 

University of Bristol, Bristol, United Kingdom

D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , M. Glowacki , J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran , L. Robertshaw , V.J. Smith , K. Walkingshaw 

Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball , K.W. Bell , A. Belyaev⁷⁷ , C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , A. Elliot , K.V. Ellis , K. Harder , S. Harper , J. Linacre , K. Manolopoulos , D.M. Newbold , E. Olaiya , D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh , C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams 

Imperial College, London, United Kingdom

I. Andreou , R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller , C.A. Carrillo Montoya , G.S. Chahal⁷⁸ , D. Colling , J.S. Dancu , I. Das , P. Dauncey , G. Davies , M. Della Negra , S. Fayer , G. Fedi , G. Hall , A. Howard , G. Iles , C.R. Knight , P. Krueper , J. Langford , K.H. Law , J. León Holgado , L. Lyons , A.-M. Magnan , B. Maier , S. Mallios , M. Mieskolainen , J. Nash⁷⁹ , M. Pesaresi , P.B. Pradeep , B.C. Radburn-Smith , A. Richards , A. Rose , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , L.H. Vage , T. Virdee²⁸ , M. Vojinovic , N. Wardle , D. Winterbottom 

Brunel University, Uxbridge, United Kingdom

J.E. Cole , A. Khan , P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish , J. Dittmann , K. Hatakeyama , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant , C. Sutantawibul , J. Wilson 

Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , A.E. Simsek , S.S. Yu 

The University of Alabama, Tuscaloosa, Alabama, USA

B. Bam , A. Buchot Perraguin , R. Chudasama , S.I. Cooper , C. Crovella , S.V. Gleyzer , E. Pearson , C.U. Perez , P. Rumerio⁸⁰ , E. Usai , R. Yi 

Boston University, Boston, Massachusetts, USA

A. Akpinar , C. Cosby , G. De Castro , Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , D. Gastler , F. Golf , S. Jeon , J. O'cain , I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , A.G. Zecchinelli 

Brown University, Providence, Rhode Island, USA

G. Benelli , D. Cutts , L. Gouskos , M. Hadley , U. Heintz , J.M. Hogan⁸¹ , T. Kwon , G. Landsberg , K.T. Lau , D. Li , J. Luo , S. Mondal , T. Russell , S. Sagir⁸² , X. Shen , F. Simpson , M. Stamenkovic , N. Venkatasubramanian , X. Yan 

University of California, Davis, Davis, California, USA

S. Abbott , C. Brainerd , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , S. Ostrom , W. Wei , S. Yoo , F. Zhang

University of California, Los Angeles, California, USA

K. Adamidis , M. Bachtis , D. Campos , R. Cousins , A. Datta , G. Flores Avila , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , Y.f. Lo , E. Manca , A. Nunez Del Prado , D. Saltzberg , V. Valuev

University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , M. Gordon , G. Hanson , W. Si 

University of California, San Diego, La Jolla, California, USA

A. Aportela , A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , L. Giannini , Y. Gu , J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , M. Quinnan , B.V. Sathia Narayanan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil

University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan , C. Campagnari , K. Downham , C. Grieco , J. Incandela , J. Kim , A.J. Li , P. Masterson , H. Mei , J. Richman , S.N. Santpur , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart , T.Á. Vámi , S. Wang , D. Zhang

California Institute of Technology, Pasadena, California, USA

S. Bhattacharya , A. Bornheim , O. Cerri , A. Latorre , J. Mao , H.B. Newman , G. Reales Gutiérrez , M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu

Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , P. Bryant , M. Cremonesi , V. Dutta , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil , C. Liu , T. Mudholkar , S. Murthy , P. Palit , K. Park , M. Paulini , A. Roberts , A. Sanchez , W. Terrill

University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , G. Karathanasis , N. Manganelli , J. Parkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo

Cornell University, Ithaca, New York, USA

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan , S. Hogan , P. Kotamnives , J. Monroy , M. Oshiro , J.R. Patterson , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou

Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , G. Cummings , J. Dickinson , I. Dutta , V.D. Elvira , Y. Feng , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green , A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig , J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima , K.H.M. Kwok , S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , C. Madrid , K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg , V. Papadimitriou

N. Pastika , K. Pedro , C. Pena⁸³ , F. Ravera , A. Reinsvold Hall⁸⁴ , L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , I. Zoi 

University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K. Matchev , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , Y. Takahashi , J. Wang 

Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhimi , A. Askew , S. Bower , R. Hashmi , R.S. Kim , S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, M. Wulansatiti , R. Yohay , J. Zhang

Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani , M.M. Baarmand , S. Butalla , S. Das , T. Elkafrawy¹⁸ , M. Hohlmann , E. Yanes

University of Illinois Chicago, Chicago, Illinois, USA

M.R. Adams , A. Baty , C. Bennett, R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , M. Hawsworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , D. S. Lemos , A.H. Merrit , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , R. Pradhan , E. Prifti, T. Roy , S. Rudrabhatla , N. Singh, M.B. Tonjes , N. Varelas , M.A. Wadud , Z. Ye , J. Yoo 

The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini , D. Blend, K. Dilsiz⁸⁵ , L. Emediato , G. Karaman , O.K. Köseyan , J.-P. Merlo, A. Mestvirishvili⁸⁶ , O. Neogi, H. Ogul⁸⁷ , Y. Onel , A. Penzo , C. Snyder, E. Tiras⁸⁸ 

Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz 

The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , M. Pitt , S. Popescu⁸⁹ , C. Rogan , C. Royon , R. Salvatico , S. Sanders , C. Smith , G. Wilson 

Kansas State University, Manhattan, Kansas, USA

B. Allmond , R. Gujju Gurunadha , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli , D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman, Y.M. Chen , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim, Y. Lai , S. Lascio , A.C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , M.M. Paranjpe, E. Popova⁹⁰ , A. Shevelev , L. Wang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

J. Bendavid , I.A. Cali , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, G. Grosso, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn , A. Novak , M.I. Park 

C. Paus , C. Reissel , C. Roland , G. Roland , S. Rothman , G.S.F. Stephans , Z. Wang , B. Wyslouch , T. J. Yang 

University of Minnesota, Minneapolis, Minnesota, USA

B. Crossman , B.M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , M. Revering , R. Rusack , R. Saradhy , N. Strobbe 

University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 

State University of New York at Buffalo, Buffalo, New York, USA

H. Bandyopadhyay , L. Hay , H.w. Hsia , I. Iashvili , A. Kalogeropoulos , A. Kharchilava , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams , P. Young 

Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , J. Bonilla , B. Bylsma, M. Campana , J. Dervan , Y. Haddad , Y. Han , I. Israr , A. Krishna , J. Li , M. Lu , G. Madigan , R. McCarthy , D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari , D. Wood 

Northwestern University, Evanston, Illinois, USA

J. Bueghly, S. Dittmer , K.A. Hahn , Y. Liu , M. McGinnis , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliercio , M. Velasco

University of Notre Dame, Notre Dame, Indiana, USA

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth , K.W. Ho , K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , C. Moore , Y. Musienko¹⁵ , H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygala 

The Ohio State University, Columbus, Ohio, USA

A. Basnet , M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas , K. Wei, D.A. Wenzl, B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, USA

H. Bouchamaoui , K. Coldham, P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , R. Sharma

Purdue University, West Lafayette, Indiana, USA

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung , A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , V. Scheurer, J.F. Schulte , M. Stojanovic , J. Thieman , A. K. Viridi , F. Wang , A. Wildridge , W. Xie , Y. Yao 

Purdue University Northwest, Hammond, Indiana, USA

J. Dolen , N. Parashar , A. Pathak 

Rice University, Houston, Texas, USA

D. Acosta , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner, F.J.M. Geurts , I. Krommydas , W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 

University of Rochester, Rochester, New York, USA

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , N. Parmar , P. Parygin⁹⁰ , R. Taus 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis , M. Heindl , C. Houghton , D. Jaroslowski , S. Konstantinou , I. Laflotte , A. Lath , R. Montalvo, K. Nash, J. Reichert , H. Routray , P. Saha , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora , H. Wang 

University of Tennessee, Knoxville, Tennessee, USA

D. Ally , A.G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A.R. Kanuganti , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , T. Akhter , K. Androsov⁵⁹ , O. Bouhali⁹¹ , R. Eusebi , J. Gilmore , T. Huang , T. Kamon⁹² , H. Kim , S. Luo , R. Mueller , D. Overton , D. Rathjens , A. Safonov 

Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , N. Gogate , V. Hegde , A. Hussain , Y. Kazhykarim, K. Lamichhane , S.W. Lee , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

University of Virginia, Charlottesville, Virginia, USA

B. Cardwell , H. Chung, B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , C. Neu 

Wayne State University, Detroit, Michigan, USA

S. Bhattacharya , P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind , S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless , J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , L. Pétré , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

Authors affiliated with an international laboratory covered by a cooperation agreement with CERN

G. Gavrilov , V. Golovtcov , Y. Ivanov , V. Kim⁹³ , P. Levchenko⁹⁴ , V. Murzin , V. Oreshkin , D. Sosnov , V. Sulimov , L. Uvarov , A. Vorobyev[†]

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , D. Budkouski , I. Golutvin[†] , I. Gorbunov , V. Karjavine , V. Korenkov , A. Lanev , A. Malakhov , V. Matveev⁹³ , V. Palichik , V. Perelygin , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , V. Smirnov , O. Teryaev , N. Voytishin , B.S. Yuldashev⁹⁵, A. Zarubin , I. Zhizhin , Yu. Andreev , A. Dermenev , S. Gninenko , N. Golubev , A. Karneyeu , D. Kirpichnikov 

M. Kirsanov , N. Krasnikov , I. Tlisova , A. Toropin , T. Aushev , V. Gavrilov , N. Lychkovskaya , A. Nikitenko^{96,97} , V. Popov , A. Zhokin , R. Chistov⁹³ , M. Danilov⁹³ , S. Polikarpov⁹³ , V. Andreev , M. Azarkin , M. Kirakosyan, A. Terkulov , E. Boos , V. Bunichev , M. Dubinin⁸³ , L. Dudko , A. Ershov , V. Klyukhin , O. Kodolova⁹⁷ , A. Markina, S. Obraztsov , M. Perfilov , V. Savrin , G. Vorotnikov , V. Blinov⁹³, T. Dimova⁹³ , A. Kozyrev⁹³ , O. Radchenko⁹³ , Y. Skovpen⁹³ , V. Kachanov , D. Konstantinov , S. Slabospitskii , A. Uzunian , A. Babaev , V. Borshch , D. Druzhkin , V. Chekhovsky, V. Makarenko

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Ghent University, Ghent, Belgium

⁴Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁵Also at Universidade Estadual de Campinas, Campinas, Brazil

⁶Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁷Also at UFMS, Nova Andradina, Brazil

⁸Also at University of Chinese Academy of Sciences, Beijing, China

⁹Also at China Center of Advanced Science and Technology, Beijing, China

¹⁰Also at University of Chinese Academy of Sciences, Beijing, China

¹¹Also at China Spallation Neutron Source, Guangdong, China

¹²Now at Henan Normal University, Xinxiang, China

¹³Also at University of Shanghai for Science and Technology, Shanghai, China

¹⁴Now at The University of Iowa, Iowa City, Iowa, USA

¹⁵Also at an institute formerly covered by a cooperation agreement with CERN

¹⁶Also at Zewail City of Science and Technology, Zewail, Egypt

¹⁷Also at British University in Egypt, Cairo, Egypt

¹⁸Now at Ain Shams University, Cairo, Egypt

¹⁹Also at Purdue University, West Lafayette, Indiana, USA

²⁰Also at Université de Haute Alsace, Mulhouse, France

²¹Also at Istinye University, Istanbul, Turkey

²²Also at The University of the State of Amazonas, Manaus, Brazil

²³Also at University of Hamburg, Hamburg, Germany

²⁴Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²⁵Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

²⁶Also at Brandenburg University of Technology, Cottbus, Germany

²⁷Also at Forschungszentrum Jülich, Juelich, Germany

²⁸Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

²⁹Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

³⁰Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania

³¹Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

³²Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

³³Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt

³⁴Also at Punjab Agricultural University, Ludhiana, India

³⁵Also at University of Visva-Bharati, Santiniketan, India

³⁶Also at Indian Institute of Science (IISc), Bangalore, India

³⁷Also at IIT Bhubaneswar, Bhubaneswar, India

³⁸Also at Institute of Physics, Bhubaneswar, India

³⁹Also at University of Hyderabad, Hyderabad, India

- ⁴⁰ Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ⁴¹ Also at Isfahan University of Technology, Isfahan, Iran
- ⁴² Also at Sharif University of Technology, Tehran, Iran
- ⁴³ Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴⁴ Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁴⁵ Also at Helwan University, Cairo, Egypt
- ⁴⁶ Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁴⁷ Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁴⁸ Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁴⁹ Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- ⁵⁰ Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- ⁵¹ Also at Lulea University of Technology, Lulea, Sweden
- ⁵² Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵³ Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
- ⁵⁴ Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁵ Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- ⁵⁶ Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁵⁷ Also at Saegis Campus, Nugegoda, Sri Lanka
- ⁵⁸ Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁵⁹ Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶⁰ Also at University of Vienna, Vienna, Austria
- ⁶¹ Also at Universität Zürich, Zurich, Switzerland
- ⁶² Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶³ Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- ⁶⁴ Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶⁵ Also at Konya Technical University, Konya, Turkey
- ⁶⁶ Also at Izmir Bakircay University, Izmir, Turkey
- ⁶⁷ Also at Adiyaman University, Adiyaman, Turkey
- ⁶⁸ Also at Bozok Universiteleri Rektörlüğü, Yozgat, Turkey
- ⁶⁹ Also at Marmara University, Istanbul, Turkey
- ⁷⁰ Also at Milli Savunma University, Istanbul, Turkey
- ⁷¹ Also at Kafkas University, Kars, Turkey
- ⁷² Now at Istanbul Okan University, Istanbul, Turkey
- ⁷³ Also at Hacettepe University, Ankara, Turkey
- ⁷⁴ Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- ⁷⁵ Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁶ Also at Yildiz Technical University, Istanbul, Turkey
- ⁷⁷ Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁷⁸ Also at IPPP Durham University, Durham, United Kingdom
- ⁷⁹ Also at Monash University, Faculty of Science, Clayton, Australia
- ⁸⁰ Also at Università di Torino, Torino, Italy
- ⁸¹ Also at Bethel University, St. Paul, Minnesota, USA
- ⁸² Also at Karamanoğlu Mehmetbey University, Karaman, Turkey

⁸³Also at California Institute of Technology, Pasadena, California, USA

⁸⁴Also at United States Naval Academy, Annapolis, Maryland, USA

⁸⁵Also at Bingol University, Bingol, Turkey

⁸⁶Also at Georgian Technical University, Tbilisi, Georgia

⁸⁷Also at Sinop University, Sinop, Turkey

⁸⁸Also at Erciyes University, Kayseri, Turkey

⁸⁹Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania

⁹⁰Now at another institute formerly covered by a cooperation agreement with CERN

⁹¹Also at Texas A&M University at Qatar, Doha, Qatar

⁹²Also at Kyungpook National University, Daegu, Korea

⁹³Also at another institute formerly covered by a cooperation agreement with CERN

⁹⁴Also at Northeastern University, Boston, Massachusetts, USA

⁹⁵Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan

⁹⁶Also at Imperial College, London, United Kingdom

⁹⁷Now at Yerevan Physics Institute, Yerevan, Armenia