

Echinodermata of Kaštela Bay (Adriatic Sea, Croatia)

Dušan ZAVODNIK

*Center for Marine Research, "Ruđer Bošković" Institute, 52210 Rovinj, Croatia,
E-mail: dzavodnik@cim.irb.hr*

Records of the Kaštela Bay benthic fauna are reviewed with special regard to Echinodermata. Thirty three echinoderm species and one ecophenotype are listed among which eight species are newly recorded in the area. Morphological and ecological notes are provided for few selected taxa.

Key words: Echinodermata, Kaštela Bay, Adriatic Sea

INTRODUCTION

The Kaštela Bay is one of the most studied coastal areas in the eastern Adriatic Sea. The area became a subject of the special interest about seventy years ago when, at that time named as the Oceanographical Institute was built at the cap of Marjan promontory (ZORE-ARMANDA and ALAJBEG, 1995). Initial basic hydrographic and biological research was soon enlarged by fishery, and recently pollution and nature preservation studies.

Many informations are available about the overall characteristics and seasonal dynamics of physical and chemical parameters, phyto- and zooplankton, and fish in the Bay (i.e. GAMULIN, 1938; ERCEGOVIĆ, 1940; JUKIĆ and ŽUPANOVIĆ, 1965; VUČETIĆ, 1965; ALFIREVIĆ, 1966; PUCHER-PETKOVIĆ, 1966, 1971; KAČIĆ, 1972; JUKIĆ, 1973; ZORE-ARMANDA, 1976; KRŠINIĆ, 1980; PUCHER-PETKOVIĆ and MARASOVIĆ, 1980; REGNER and VUČETIĆ, 1980; REGNER, 1980; ZORE-ARMANDA *et al.*,

1991; etc.). On the contrary, in spite of a fair diversity of bottom habitats (GRUBELIĆ and ŠIMUNOVIĆ, 1997), information about the benthos is rather scarce. Benthic macroflora was studied in particular by ERCEGOVIĆ (1932, 1952), ŠPAN (1964), ANTOLIĆ (1996) and ŠPAN and ANTOLIĆ (1997). Only the few heigh groups of benthic invertebrates attracted special attention of scientists: protozoans (HADŽI, 1938 a; STILLER, 1946), sponges (BREITFUSS, 1938; HADŽI, 1938 b), cnidarians (BROCH, 1933, 1935; PAX and MÜLLER, 1953, 1962; KORNMAN, 1968), molluscs (HORVATH, 1963; SALVINI - PLAIVEN, 1972; ŠIMUNOVIĆ and GRUBELIĆ, 1992; ŠIMUNOVIĆ, 1992, 1995) and arthropods (VIETS, 1939, 1940; ZAVODNIK, 1960; KARLOVAC, 1969, 1970; PETKOVSKI and APOSTOLOV, 1974). Only the few records of echinoderms were published (ZAVODNIK, 1960; GAMULIN-BRIDA *et al.*, 1971; ŠIMUNOVIĆ, 1992, 1995; GRUBELIĆ and ŠIMUNOVIĆ, 1997; ŠIMUNOVIĆ and GRUBELIĆ, 1998).

With regard to Kaštela Bay benthic communities above all, attention was paid to rocky littoral (ŠIMUNOVIĆ, 1970), coarse sand and fine gravel (ŠIMUNOVIĆ, 1992), and terrigenous muddy habitats (GAMULIN-BRIDA *et al.*, 1971; GAMULIN-BRIDA, 1974, 1979; ŠIMUNOVIĆ, 1992). No attempt has ever been made to map the distributional pattern of benthic communities in the area.

As the echinoderm fauna in Kaštela Bay was never reviewed, with regard to biodiversity studies I think it is opportune to complete the list of GRUBELIĆ and ŠIMUNOVIĆ (1997) by my personal data not published so far.

MATERIAL AND METHODS

In the Kaštela Bay I collected echinoderms at occasional excursions in 1956-1988 (Fig. 1). At coastal stations, down to about 8 meters

depth, echinoderms were collected by skin diving. Samplings by PETERSEN grab (0.1 m²), the dredge of the "mušular" type (GAMULIN-BRIDA *et al.*, 1968), and the commercial bottom-trawl were applied at greater depth. All major types of the bottom were surveyed: bedrock covered with algal turf, loose stones, cobbles, gravel, rough sand and sand mixed with shell debris and detritic clumps, silty sand and terrigenous ooze. The material was processed by standard methods and preserved in 70 % alcohol or 4 % buffered formol solution, or specimens were dried. The M.V. HVAR cruises protocol records for the station Nr. 83 (KARLOVAC, 1956) have also been considered. Voucher specimens are kept in reference collections of the Institute of Oceanography and Fisheries, Split, the Center for Marine Research of the "Ruder Bošković" Institute, Rovinj, and the Croatian Natural History Museum, Zagreb, Croatia.




Fig. 1. Research area and stations

RESULTS

In the material studied 31 echinoderm species and one ecophenotype were identified (Table 1). Most of them are well known and

quite common in the shallow water of the Adriatic and Mediterranean benthic communities (TORTONESE, 1965). Therefore, with an aim to economize the space comments are made only on selected taxa.

Table 1. Echinoderms recorded in Kaštela Bay. Taxa not noted previously in the area are marked with an asterisk ()*

Records	HVAR Stn 83	ŠIMUNOVIĆ & GRUBELIĆ (1997)	Author's stations	Bottom Habitat	Ecological significance
CRINOIDEA					
<i>Antedon mediterranea</i> (LAMARCK, 1816)	+	+	D2,3; T1,2, 3,4,5,6	D,M,MD,S	Wed.
HOLOTHUROIDEA					
<i>Holothuria forskali</i> DELLE CHIAJE, 1823	+	+	D2; T5	M,MD	Wed.
* <i>Holothuria impatiens</i> (FORSSKÅL, 1775)			S9; T1	R, S	Wed.
<i>Holothuria polii</i> DELLE CHIAJE, 1823	+	+	S2,9	S	Sand tol.
* <i>Holothuria</i> sp.			T1,2,4,5	MD,MS,S	
<i>Holothuria tubulosa</i> GMELIN, 1788	+	+	D1,2,3;S1,2, 3,4,5,6,7,9, 10,11; T1,2, 3,4,5,6	M,S,R	Wed.
<i>Eostichopus regalis</i> CUVIER, 1817	+	+	T1,2,3,4,5,6	MD,S,SM	Wed.
<i>Oenus planci</i> (BRANDT, 1835)	+	+	D1,2; T1,3,4	MS,S	pr. CD
<i>Leptopentacta elongata</i> (DÜBEN&KOREN, 1844)					Silt tol.
<i>Leptopentacta tergestina</i> (M.SARS, 1857)					pr. CTO
<i>Labidoplax digitata</i> (MONTAGU, 1815)			G1,3,5,7	M,SM	ex. CTO
* <i>Leptosynapta inhaerens</i> (O.F.MÜLLER, 1776)			G1	SM	pr. CTO
ASTEROIDEA					
<i>Astropecten aranciacus</i> (LINNAEUS, 1758)	+	+	D2; S7; T1,3, 4,5,6	MD,MS,S	Wed.
* <i>Astropecten hispinosus</i> (OTTO, 1823)			D2	S	Sand tol.
<i>Astropecten irregularis</i> (PENNANT, 1777)	+	+	D1; T1	MS,S	Min.
* <i>Astropecten platyacanthus</i> (PHILIPPI, 1837)			S7	S	pr. SG
<i>Asterina gibbosa</i> (PENNANT, 1777)					Crypt.
<i>Anseropoda placenta</i> (PENNANT, 1777)	+	+	T1,3,4,5,6	MD,MS,S,SM	ex. CD
<i>Echinaster sepositus</i> (RETZIUS, 1783)	+	+	D1; S1,5,9;T1,6	MD,MS,R,S	Wed.
<i>Coscinasterias tenuispina</i> (LAMARCK, 1816)			S1,2,9	R	pr. PSW
<i>Marthasterias glacialis</i> (LINNAEUS, 1758)	+	+	S1,2,5,10; T1,2,3,5,6	D,G,M,R,S	Wed.
OPHIUROIDEA					
* <i>Ophiomyxa pentagona</i> (LAMARCK, 1816)			T1,2,3	MD,MS,S	pr. CD
* <i>Amphiura chiajei</i> FORBES, 1843			G7,8	M,MS	Silt tol.

Table 1. Cont'd

<i>*Amphiura filiformis</i> (O.F.MÜLLER, 1776)		G1,3,5,7	M,SM	Min.
<i>Amphipholis squamata</i> (DELLE CHIAJE, 1828)		S1,2,4,6,9,10,12	A,R	Wed.
<i>Ophiothrix fragilis</i> (ABILDGAARD, 1789)	+	D2; S1,2,4,5, 6,8,9,11	A,R,S	Wed.
<i>Ophiothrix quinquemaculata</i> (DELLE CHIAJE, 1828)		T1,2,3,4,5,6	MD,MS,S,SM	Mix.
<i>Ophioderma longicaudum</i> (RETZIUS, 1805)	+	S1,5,9,11; T3	MS,R	pr. PSW
<i>Ophiura ophiura</i> (LINNAEUS, 1758)	+	T1,3,4,5,6	MD,MS,S,SM	Wed.
ECHINOIDEA				
<i>Arbacia lixula</i> (LINNAEUS, 1758)	+	S2,5	R	ex. PSW
<i>Sphaerechinus granularis</i> (LAMARCK, 1816)	+	D2,3; S1,7,9;T1	D,G,R,S	Wed.
<i>Echinus acutus</i> LAMARCK, 1816	+	T4	SM	Wed.
<i>Psammechinus microtuberculatus</i> (BLAINVILLE, 1825)+	+	D1,2; T1,2	MD,MS,S	pr. CD
<i>Paracentrotus lividus</i> (LAMARCK, 1816)	+	S1,2,5,6,7,8, 9,10,11,12; T1,4	R,G,S,SM	pr. PSW
<i>*Echinocyamus pusillus</i> (O.F.MÜLLER, 1776)		D3	DS	Grav.

ABBREVIATIONS

Bottom - Habitat:

- A = Algal thalli
- D = Organogenic detritic deposits
- G = Gravel, cobbles and loose stones
- M = Muddy bottom
- R = Bedrock with algal turf
- S = Sandy deposits

Ecology:

- CD = Community of coastal detritic bottom
- CTO = Community of coastal terrigenous ooze
- SG = Meadows of sea-grasses
- PSW = Community of photophilic seaweeds
- ex. = Species exclusive to a specific community
- pr. = Community preferential species
- Crypt. = Species preferring concealed habitats
- Grav. = Species living in rough sand and fine gravels
- Min. = Species living in fine sand and silts
- Mix. = Species living on mixed bottoms
- Sand tol. = Species living in sands and tolerant to other fractions
- Silt tol. = Species living in silts and tolerant to other fractions
- Wed. = Species with a wide ecological distribution

Holothuria impatiens (FORSSKÅL, 1775)

South of the Kaštel Novi, one specimen was collected from the sandy bottom by dredge, at about 25 meters depth. This is the maximum depth noted in the Adriatic Sea (BELAMARIĆ and ŠERMAN, 1989). Usually, specimens were collected in the shallow water at only 0-4 meters depth (MAYER, 1937; ZAVODNIK, 1997).

Holothuria sp.

By their body and coloration characteristics, some holothurians recorded in trawl catches long ago conformed with the description of *Holothuria stellata* DELLE CHIAJE, 1923 by KOEHLER (1921, 1927). Regrettably, the identification could not be proved before the ossi-

cles had deteriorated. However, due to wide ranged variation in external features and ossicles in closely related species of *tubulosa* group (PANNING, 1939; MASSIN, personal communication), in spite of GUSTATO and VILLARI (1980) standpoint the systematic position of *H. stellati* is still dubious (ROWE, 1969, personal communication).

Labidoplax digitata (MONTAGU, 1815)

Not rare in the Kaštela Bay. In eight specimens studied no giant anchors were noted, the anchor flukes were not serrated, and a concave shaped anchor apex rarely occurred. One specimen rich in abnormal anchor ossicles was collected by grab at station G1 at 25 metres depth. Among 219 anchors analysed 56 (i.e. about 25 %) were peculiar for their variously shaped

apical part (Fig. 2). The B type anchors were most common. The C and D types were recorded only in one and two anchors, respectively. The additional two pointed apical flukes (E type) may join creating thus an intermediate drop-like hole (F type). Only one anchor ossicle was distinguished by its abnormally formed stock (G type).

Amphiura filiformis (O.F. MÜLLER, 1776)

In three specimens small rudimentary scales were visible on the disc oral side - an occurrence rarely noted in this species. It seems that in the area studied the population density does not exceed about 10 individuals per square metre. Apparently this species is more common in the Bay than is the congeneric *Amphiura chiajei*.




Fig. 2. Ossicles in *Labidoplax digitata*. A = normal anchor and anchor plates, B-G = aberrant ossicles. For explanation see text

***Amphipholis squamata* (DELLE CHIAJE, 1828)**

Occasionally noted in algal turf. One specimen parasited by an ovigerous female of *Cancerilla tubulata* DALYELL, 1851, (Crustacea, Copepoda) was collected in the neighbourhood of the Institute of Oceanography and Fisheries campus (ZAVODNIK, 1960).

***Ophiothrix quinquemaculata*
(DELLE CHIAJE, 1828)**

According to GUILLE (1965) and BARIĆ and STURMBAUER (1999) *Ophiothrix quinquemaculata* is an ecologically well defined ecophenotype of *Ophiothrix fragilis* characteristic in mediterranean coastal detritic communities (PÉRÈS and PICARD, 1964). It is very common in the central area of Kaštela Bay.

DISCUSSION AND CONCLUSIONS

Among 31 species and one ecophenotype recorded at present research, 8 taxa were not noted previously in the Kaštela Bay. With addition of three species (*Leptopentacta elongata*, *Leptopentacta tergestina* and *Asterina gibbosa*) listed by GRUBELIĆ and ŠIMUNOVIĆ (1997) but not recorded at present research, the total of 33 echinoderm species and one ecophenotype was established (Table 1). This is about 33% of the Adriatic Sea echinoderm fauna, and just 39% of taxa distributed within the depth range of the Bay, i.e. 0-51 meters. The absence of typical thermophilic and deep sea species is conceivable.

However, the species list composition proved insufficient endeavours in the research of cryptic habitats. Quite curious is the absence of irregular echinoid records. Until now, according to information available, only one live specimen and many *Echinocyamus pusillus* tests were noted whilst neither live individuals nor tests of greater species have ever been collected in the area. The phenomenon cannot be explained.

In previous taxonomic and biogeographical echinological papers the species locality area was sometimes noted by a toponomy "Split" (Ital. "Spalato") (HELLER, 1868; MARENZELLER, 1874; STOSSICH, 1883; CARUS, 1885; TORTONESE, 1935, 1965; MAYER, 1937; KOLOSVÁRY, 1940a, 1940b; ZAVODNIK, 1971; VIDOVIC-MATVEJEV, 1978). Because of a geographically uncertain position of the record, such notices were not considered in the present paper. However, there exists a possibility of collection inside the Kaštela Bay. The same procedure was applied for voucher specimens deposited in museum collections (TORTONESE, 1984; Zoological Museum in Zagreb - not published). One cannot exclude the possibility of future findings of at least some of the pertinent species in the Kaštela Bay. The voucher specimens are badly needed.

Little is known on echinoderm abundance in the area. A great share of *Eostichopus regalis* in trawl catches was noted by GAMULIN-BRIDA (1974). GRUBELIĆ and ŠIMUNOVIĆ (1998) noted great *Holothuria tubulosa* shoals in the northern part of the Bay at less than 10 meters depth. In ophiurans, perhaps the most important in turn over is the filter-feeder *Ophiothrix quinquemaculata*, which in some trawl catches was recorded as a dominant taxon.

ACKNOWLEDGEMENTS

Author is greatly obliged to the Director of the Institute of Oceanography and Fisheries, Split for a kind permission to consult protocols of the M. V. HVAR cruises, and to staff of the research vessels BIOS, PREDVODNIK and VILA VELEBITA for help in collections. Thanks are due also to Miss V. EMRIĆ, who put at disposal echinoderms she has collected at three coastal stations, and to two anonymous referees who contributed to the improvement of the manuscript. The paper was prepared in the frames of the biodiversity programme Nr. 00981302 supported by the Ministry of Science and Technology of the Republic of Croatia.

REFERENCES

- ALFIREVIĆ, S. 1966. Les sources sous-marines de la baie de Kaštela. *Acta Adriat.*, 10(12): 39 pp.
- ANTOLIĆ, B. 1996. Floristički sastav i struktura zajednice posidonije, *Posidonia oceanica* (L.) Delile, u srednjem Jadranu. Ph.D. Thesis, Sveučilište u Zagrebu, 349 pp.
- BARIĆ, S. and C. STURMBAUER. 1999. Ecological parallelism and cryptic species in the genus *Ophiothrix* derived from mitochondrial DNA sequences. *Moll. Phylog. Evol.*, 11: 157-162.
- BELAMARIĆ, J. and D. ŠERMAN. 1989. Ekološka studija podmorja Lokruma. in: Otok Lokrum, HED. Ekološke monografije, 1: 361-411.
- BREITFUSS, L. 1938. Zur Kenntnis der Kalkschwammfauna Jugoslawiens. *God. Oceanogr. Inst. Split*, 1: 178-181.
- BROCH, H. 1933. Zur Kenntnis der adriatischen Hydroidenfauna von Split. *Skr. norske Vidensk.-Akad., Mat.-Naturw. Kl.*, 1933 (4): 115 pp.
- BROCH, H. 1935. Beobachtungen an einigen Seichtwasser-Anthozoen von Split. *Biologia Generalis*, 11: 1-14.
- CARUS, J.V. 1885. Prodromus faunae mediterraneae. I. E. Schweizerbart, Stuttgart, 524 pp.
- ERCEGOVIĆ, A. 1932. Wellengang und Lithophytenzone an der ostadiatischen Küste. *Acta Adriat.*, 1(3): 20 pp.
- ERCEGOVIĆ, A. 1940. Ishrana srdele (*Clupea pilchardus* Walb.) u stadiju metamorfoze. *God. Oceanogr. Inst. Split*, 2: 26-44.
- ERCEGOVIĆ, A. 1952. Jadranske cistozire. Fauna i flora Jadrana, 2. Institut za oceanografiju i ribarstvo, Split, 212 pp.
- GAMULIN, T. 1938. Izvještaj o istraživanjima pojavljivanja planktonskih kopepoda u okolini Splita u godini 1936-37. *God. Oceanogr. Inst. Split*, 1: 52-61.
- GAMULIN-BRIDA, H. 1974. Biocoenoses benthiques de la Mer Adriatique. *Acta Adriat.*, 15(9): 3-102.
- GAMULIN-BRIDA, H. 1979. Litoralne biocenoze na području srednjodalmatinskih otoka. *Acta biol. Jugosl. akad. zn. umjet.*, 8: 37-63.
- GAMULIN-BRIDA, H., S. ALFIREVIĆ, D. CRNKOVIC, A. ŠIMUNOVIĆ, S. JUKIĆ, A. POŽAR, and M. LEGAC. 1971. Contribution à l'étude de la dynamique de certaines communi- nautés benthiques en Adriatique. *Thalassia Jugosl.*, 7: 57-66.
- GAMULIN-BRIDA, H., A. POŽAR and D. ZAVODNIK. 1968. Contribution aux recherches sur la bionomie benthique des fonds meubles de l'Adriatique du nord (II). *Biol. Glas.*, 21: 157-201.
- GRUBELIĆ, I. and A. ŠIMUNOVIĆ. 1997. Prilog poznavanju bentoske faune šireg područja Splita. in: Finka B. (Editor). *Tisuću godina prvoga spomena ribarstva u Hrvata*. Hrvatska akademija znanosti i umjetnosti etc., Zagreb, 613-627.
- GUILLÉ, A. 1965. Contribution à l'étude de la systématique et de l'écologie d'*Ophiothrix quinquemaculata* D.Ch. *Vie Milieu*, 15: 243-301.
- GUSTATO, G. and A. VILLARI. 1980. About the question of the taxonomic status of *Holothuria stellati* Delle Chiaje. in: M. Jangoux (Editor). *Echinoderms: Present and Past. Proc. European Colloquium on Echinoderms*, Brussels, 3-8 September 1979. Balkema, Rotterdam, The Netherlands, pp. 107-110.
- HADŽI, J. 1938 a. Beitrag zur Kenntnis der adriatischen Folliculiniden (Inf., Heterotricha). 1. Subfamilie: Eufolliculininae. *Acta Adriat.*, 2(1): 46 pp.
- HADŽI, J. 1938 b. Zoološke beleške. *God. Oceanogr. Inst. Split*, 1: 181-193.
- HELLER, C. 1868. Zoophyten und Echinodermen des Adriatischen Meeres. Wien, 86 pp.
- HORVATH, A. 1963. Adriatic molluscs of the Split area. *Acta Adriat.*, 10(4): 12 pp.
- JUKIĆ, S. 1973. The contribution to the knowledge of bogue's (*Boops boops* L.) nourishment in the bay of Kaštela. *Acta Adriat.*, 14(11): 15 pp.
- JUKIĆ, S. and Š. ŽUPANOVIĆ. 1965. Relations entre la température et l'intensité de l'alimentation chez *Mullus barbatus* L. et *Pagellus erythrinus* L. dans la baie de Kaštela. *FAO Proc. gen. Fish. Coun. Medit.*, 8: 173-177.
- KAČIĆ, I. 1972. The behaviour, distribution and quantity of sardines in the bay of Kaštela. *Acta Adriat.*, 14(10): 32 pp.
- KARLOVAC, O. 1956. Station list of the M. V. "Hvar" fishery-biological cruises 1948-1949. Izvješća-Reports. The M.V. "Hvar" cruises-researches into fisheries biology, 1948-49, 1(3): 177 pp.

- KARLOVAC, O. 1969. Prilog poznavanju faune rakova desetonožaca u priobalnim vodama srednjeg Jadrana. I. Pomorski zbornik, 7: 967-974.
- KARLOVAC, O. 1970. Prilog poznavanju faune rakova desetonožaca u priobalnim vodama srednjeg Jadrana. II. Pomorski zbornik, 8: 883-899.
- KOEHLER, R. 1921. Echinodermes. Faune de France, I. Lechevalier, Paris, 210 pp.
- KOEHLER, R. 1927. Les Echinodermes des mers d'Europe. II. Doin, Paris, 339 pp.
- KOLOSVÁRY, G. v. 1940a. Variationsstudien über *Becheria adriatica* (Heding) und *Cucumaria elongata* Düben et Koren aus Dalmatien. Folia zool. hydrobiol., 10: 109-111.
- KOLOSVÁRY, G. v. 1940b. Über die geographische Verbreitung einiger adriatischen Echinodermen. Folia zool. hydrobiol., 10: 371-381.
- KORNMANN, J. 1968. Ergänzende Mitteilungen über die Anthozoenfauna der Bucht von Kaštela bei Split. Bilješke-Notes. Inst. oceanogr. i rib., Split, (22): 4 pp.
- KRŠINIĆ, F. 1980. Kvalitativna i kvantitativna istraživanja tintinida uz istočnu obalu Jadranskog mora. Acta Adriat., 21(1): 19-104.
- MARENZELLER, E. v. 1874. Kritik adriatischer Holothurien. Verh. zool. bot. Ges. Wien, 24: 299-320.
- MAYER, B. 1937. Die Holothurien der Adria. Thalassia, 2(9): 54 pp.
- PANNING, A. 1939. Holothurien von den Kanaren und von Dakar. Vidensk. Medd. dansk naturh. Foren., 103: 523-546.
- PAX, F. and I. MÜLLER. 1953. Die Anthozoenfauna der Bucht von Kaštela bei Split. Acta Adriat., 5(1): 3-35.
- PAX, F. and I. MÜLLER. 1962. Die Anthozoenfauna der Adria. Fauna et Flora adriatica, 3. Institut za oceanografiju i ribarstvo, Split, 343 pp.
- PÉRÈS, J.-M. and J. PICARD. 1964. Nouveau manuel de bionomie benthique de la Méditerranée. Recl Trav. Stn mar. Endoume, 31(47): 5-137.
- PETKOVSKI, T. and A. APOSTOLOV. 1974. Zweite *Nannomesochra*-Art (Copepoda, Harpacticoida) aus der Adria, *N. zavodniki* n.sp. Fragmenta Balcanica, 10(1-221): 1-9.
- PUCHER-PETKOVIĆ, T. 1966. Végétation des Diatomées pélagiques de l'Adriatique moyenne. Acta Adriat., 13(1): 98 pp.
- PUCHER-PETKOVIĆ, T. 1971. Recherches sur la production primaire et la densité des populations du phytoplancton en Adriatique moyenne (1962-1967). Rapp. Comm. int. Mer Médit., 20: 339-343.
- PUCHER-PETKOVIĆ, T. and I. MARASOVIĆ. 1980. Développement des populations phytoplanctoniques caractéristiques pour un milieu eutrophisé (baie de Kaštela, Adriatique centrale). Acta Adriat., 21(2): 79-93.
- REGNER, D. and T. VUČETIĆ. 1980. Seasonal and multiannual fluctuations of copepods in the Kaštela Bay (1960-1969). Acta Adriat., 21(2): 101-122.
- REGNER, S. 1980. The larval stages of fish in the Kaštela Bay. Acta Adriat., 21(2): 123-136.
- ROWE, F.W.E. 1969. A review of the family Holothuriidae (Holothurioidea : Aspidochirotida). Bull. Br. Mus. nat. Hist. (Zool.), 18: 119-170.
- SALVINI-PLAWEN, L.v. 1972. Die Caudofoveata des Mittelmeeres und das Genus *Scutopus* (Mollusca, Aculifera). Proc. 5th Europ. Mar. Biol. Symp., Piccin, Padova: 27-51.
- STILLER, J. 1946. Beitrag zur Kenntnis der Peritrichen-Fauna der Adria bei Split (Spalato). Ann. Hist. nat. Mus. natn. Hung. Zool., 39(3): 59-74.
- STOSSICH, M. 1883. Prospetto della fauna del Mare Adriatico. V. Echinodermata. Boll. Soc. adr. Sci. nat. Trieste, 8: 90-110.
- ŠIMUNOVIĆ, A. 1970. Recherches écologiques sur les peuplements faunistiques de l'étage supralittoral des côtes rocheuses de l'Adriatique moyenne. Acta Adriat., 14(2): 39 pp.
- ŠIMUNOVIĆ, A. 1992. Zoobenthos of Kaštela Bay - state and problems. Acta Adriat., 33: 163-168.
- ŠIMUNOVIĆ, A. 1995. Ecological study of Prosobranchiata (Gastropoda) in the eastern part of the Adriatic Sea and their relationship to benthic biocoenoses. Acta Adriat., 36: 3-162.
- ŠIMUNOVIĆ, A. and I. GRUBELIĆ. 1992. Biological and ecological studies of the dateshell (*Lithophaga lithophaga* L.) from the eastern Adriatic Sea. Period. biol., 94: 187-192.
- ŠIMUNOVIĆ, A. and I. GRUBELIĆ. 1998. A contribution to the knowledge of the species *Holothuria tubulosa* Gmelin, 1788 (Holothuria, Echinodermata) in the coastal area of the eastern Adriatic. Acta Adriat., 39: 13-23.
- ŠPAN, A. 1964. Preliminarna kvantitativna ispitivanja cistozira u okolini Splita. Acta Adriat., 11: 255-260.

- ŠPAN, A. and B. ANTOLIĆ. 1997. Sastav, rasprostranjenost i stanje bentoske flore na širem području Splita. in: Finka B. (Editor). Tisuću godina prvoga spomena ribarstva u Hrvata. Hrvatska akademija znanosti i umjetnosti etc., Zagreb, pp. 493-513.
- TORTONESE, E. 1935. Contributo alla conoscenza degli Echinodermi mediterranei. Annali Mus. civ. Stor. nat. Genova, 57: 219-272.
- TORTONESE, E. 1965. Echinodermata. Fauna d'Italia, 6. Calderini, Bologna, 419 pp.
- TORTONESE, E. 1984. Echinodermi del Museo di Storia naturale di Verona. Mus. civ. Stor. nat. Verona, Ser. Cataloghi, 2: 73 pp.
- VIDOVIĆ-MATVEJEV, A. 1978: Catalogue of the Adriatic Echinoderms. Acta Adriat., 17(15): 3-24.
- VIETS, K. 1939-1940. Meeresmilben aus der Adria (Halacaridae und Hydrachnella, Acari). Arch. Naturgesch., N.F., 8: 518-550 (1939); 9: 1-135 (1940).
- VUČETIĆ, T. 1965. Long-term observations of zooplankton fluctuation in the bay of Kaštela. FAO Proc. gen. Fish. Coun. Medit., 8: 261-265.
- ZAVODNIK, D. 1960. O kopepodu *Cancerilla tubulata* Dal., ektoparazitu na kačjerepu *Amphipholis squamata* (D.Ch.). Biol. Vestn., 7: 81-83.
- ZAVODNIK, D. 1971. Osrt na poznavanje iregularnih ježinaca (Echinoidea, Irregularia) Jadranskog mora. Zbornik I. Simp. sistemat. Jugosl., Sarajevo, 215-219.
- ZAVODNIK, D. 1997. Echinodermata of the marine National Park "Kornati" (Adriatic Sea). Period. biol., 99: 367-380.
- ZORE-ARMANDA, M. 1976. Kaštelski zaljev. Pomorska enciklopedija (2.izd.). Leksikografski zavod, Zagreb. 3: 533-534.
- ZORE-ARMANDA, M. and M. ALAJBEG. 1995. 65 godina Instituta za oceanografiju i ribarstvo, Split. Institut za oceanografiju i ribarstvo, Split, 144 pp.
- ZORE-ARMANDA, M., M. BONE, V. DADIĆ, M. MOROVIĆ, D. RATKOVIĆ, L. STOJANOSKI and I. VUKADIN. 1991. Hydrographic properties of the Adriatic Sea in the period from 1971 through 1983. Acta Adriat., 32: 1-547.

Accepted: 28 September 1999

Bodljikaši (Echinodermata) Kaštelskog zaljeva

Dušan ZAVODNIK

*Zavod za istraživanje mora, Institut "Ruder Bošković", 52210 Rovinj, Hrvatska
E-mail: dzavodnik@cim.irb.hr*

SAŽETAK

Dat je pregled izvornih podataka o bentoskoj fauni Kaštelskog zaljeva s posebnim osvrtom na bodljikaše. Utvrđena je nazočnost 33 vrste i jednog ekofenotipa bodljikaša, od kojih osam do sada nije bilo spomenuto za to područje. Navode se i značajniji podaci o ekologiji i morfologiji izabranih svojti.