



# The polymorphisms in Toll-like receptor genes and cancer risk

ANA GOMAZ  
JASMINKA PAVELIĆ  
TANJA MATIJEVIĆ GLAVAN

Ruder Bošković Institute  
Division of Molecular Medicine  
Laboratory of Molecular Oncology  
Bijenička cesta 54, 10 000 Zagreb, Croatia

#### Correspondence:

Tanja Matijević Glavan  
Ruder Bošković Institute  
Division of Molecular Medicine  
Laboratory of Molecular Oncology  
Bijenička cesta 54, 10 000 Zagreb, Croatia  
E-mail: tmatijev@irb.hr

**Key words:** Toll-like receptor, polymorphism, cancer, immune response, inflammation

## Abstract

*Toll-like receptors (TLRs) are a family of pattern recognition receptors and have important role in pathogen recognition including activation of innate and acquired immune responses. They are mainly expressed in epithelium but have also been found in cancer cells, where they can have pro-tumorigenic and anti-tumorigenic effect. Single nucleotide polymorphisms (SNPs) may change protein structure and influence receptor function. Polymorphisms in TLR genes have been linked with various cancer types and reports were often contradictory. The strongest association with increased cancer risk was found for TLR2 polymorphism –196 to –174 del ( $\Delta 22$ ) (gastric, gallbladder and cervical cancer) and TLR4 polymorphisms rs4986790 (gastrointestinal cancers and lymphoma) and rs4986791 (gastrointestinal cancers, nasopharyngeal and gallbladder cancer). Other polymorphisms associated with cancer risk include TLR10-TLR1-TLR6 gene cluster polymorphisms rs10008492 (non-Hodgkin's lymphoma) and rs7696175 (breast cancer), TLR6 rs5743815 (non-Hodgkin's lymphoma), TLR10 rs11466657 (meningioma), TLR2 GT microsatellite repeat number polymorphism (colorectal cancer) and rs3804100 (MALT lymphoma), TLR3 829A>C (nasopharyngeal cancer), TLR4 11350G>C (nasopharyngeal cancer), rs1927911 and rs10759931 (prostate cancer), TLR5 rs5744174 (gastric cancer) and TLR9 rs5743836 (Hodgkin's lymphoma) and rs352140 (Hodgkin's lymphoma and cervical cancer). Discrepancies in results of different studies regarding TLR polymorphisms and cancer risk may have arisen due to insufficient sample size, differences in ethnicity or age, undetected infections or other environmental factors.*

## 1. INTRODUCTION

Toll-like receptors (TLRs) are a family of innate immune receptors known as pattern recognition receptors (PRRs) and they recognize molecular motifs related to pathogens or pathogen associated molecular patterns (PAMPs) (1). TLRs have a crucial role in recognition of many different pathogens, including viruses, bacteria, protozoa and fungi, and due to their important function, they are evolutionary conserved (2). They were first described in *Drosophila* in 1984, and were later discovered in vertebrates, including humans (1, 3). To date, 13 TLRs were discovered in mammals, and 10 are functional in humans (2). Human genes encoding TLRs reside on chromosomes 1 (*TLR5*), 3 (*TLR9*), 4 (*TLR1*, *TLR2*, *TLR3*, *TLR6* and *TLR10*), 9 (*TLR4*) and X (*TLR7* and *TLR8*) (1, 4). TLR signaling begins with binding of specific ligand, continues with dimerization and signal transduction into the cell followed by

activation of three main transcription factors: interferon-regulatory factor (IRF3, IRF5 and IRF7), NF- $\kappa$ B and AP1. They then promote the transcription of inflammation mediators: cytokines, chemokines and interferons (2, 5). TLRs are also important in activation of adaptive immunity: downstream signaling and expression of cytokines leads to maturation of dendritic cells and activation of B and T cells (5).

## 2. TOLL-LIKE RECEPTORS: STRUCTURE, LOCALIZATION AND FUNCTION

TLRs are type I transmembrane glycoproteins consisting of three domains: leucine rich ligand binding ectodomain, transmembrane region, and cytoplasmic TIR-domain (the Toll/interleukin-1 receptor domain). Ectodomain contains multiple leucine rich repeats (LRRs) which are crucial in recognition and binding of pathogen motifs and also in forming characteristic horseshoe shape of TLRs, a structure important in ligand binding and downstream signaling (6). TIR-domain dimerizes after the ligand binding in the ectodomain enabling the recruitment of TIR domain containing adaptor molecules which transduce the signal downstream: MyD88 (myeloid differentiation factor 88), MAL (MyD88 adaptor-like), TRIF (Toll receptor-associated activator of interferon) and TRAM (Toll receptor-associated molecule) (2, 6). All TLRs except TLR3 use MyD88 for signal transduction, and TLR3 depends on TRIF. TLR4 can activate both MyD88 and TRIF. TLRs can form homodimers (like TLR3-TLR3) or heterodimers (such as TLR1-TLR2 and TLR2-TLR6) (2). Due to differences in ectodomain, different TLRs bind different PAMPs, such as lipids and lipoproteins (TLR1, TLR2, TLR4 and TLR6), proteins (TLR5 and mouse TLR11), and microbial nucleic acids (TLR3, TLR7, TLR8 and TLR9), while ligands of TLR10 and mouse TLR12 and TLR13 have not yet been identified (7). Nature of ligands they bind determines the location of TLRs in the cell: TLR1, TLR2, TLR4, TLR5 and TLR6 bind components of pathogen envelope or cell wall and are expressed on the surface of the cell (as well as TLR10), while TLR3, TLR7, TLR8 and TLR9 bind nucleic acids and are mainly located in the endoplasmic reticulum membrane or endosomal membrane (2). TLRs are mainly expressed in lymphoid tissues, innate immune cells (dendritic cells, macrophages, monocytes, polymorphonuclear leukocytes, NK cells) and lymphocytes T and B. However, they can also be expressed in different cell types, including cancer cells and cancer cell lines (5, 8). In normal tissues, TLRs are highly expressed in many epithelial cells (9).

## 3. TOLL-LIKE RECEPTORS IN CANCER

TLRs participate in many important cellular processes: cell proliferation, survival, apoptosis, cell migration, metastasis and angiogenesis. Due to their important role in inflammation, it is considered that they can have a pro-tumorigenic and anti-tumorigenic function, an effect


known as a „double edged sword“ (10, 11). In some cases enhanced TLR activation can lead to apoptosis of cancer cells and more efficient elimination of tumor cells by immune system, while in others it may support tumorigenesis by promoting chronic inflammation (8, 12, 13). TLR signaling has been implicated in various autoimmune and inflammatory diseases and chronic inflammation is known to create a microenvironment rich in growth and survival factors and has been linked with development of many types of cancer (13). TLR expression was reported in several cancer types and cancer cell lines. It was found that TLR4 and TLR5 are expressed in gastric epithelium infected with *Helicobacter pylori* as well as in precursor lesions and in gastric cancer cells (14). It is considered that TLRs enable cells to interact with *H. pylori* which can induce tumorigenic factors and may promote cancer development. TLR expression has also been found in colon cancer, hepatocellular carcinoma, ovarian and cervical cancers, breast and prostate cancers, lung cancer, melanoma and neuroblastoma (9). TLR expression in cancer cells has been linked with tumor progression, evasion of immune system surveillance, apoptosis and survival. TLR4 expression in lung cancer cells is linked with production of immunosuppressive cytokines (TGF $\beta$ ), angiogenic factors VEGF and IL-8, and with increased resistance to apoptosis (15). Silencing of TLR4 expression in breast cancer cell line MDA-MB-231 significantly decreased cell proliferation and production of proinflammatory cytokines IL-6 and IL-8 (16). In ovarian cancer cells and cell lines TLR4 activation by lipopolysaccharide or Paclitaxel activated NF- $\kappa$ B, promoted production of IL-8, IL-6, VEGF and MCP-1, and inhibited Paclitaxel mediated apoptosis, while TLR4 silencing lead to loss of resistance to Paclitaxel (17). TLR2 mRNA expression was significantly higher in sporadic colorectal cancer cells than in noncancerous cells which could implicate its role in cancerogenesis (18).

## 4. THE POLYMORPHISMS OF TLR GENES

Single nucleotide polymorphisms (SNPs) may often cause no change in protein function and pass unnoticed, but can also lead to amino acid substitutions and changes in protein function and stability, splicing, mRNA stability, post-translational modifications or regulation of expression (Figure 1A and 1B) (19). SNPs in TLR ectodomain (in LRRs) can affect binding of ligands and SNPs in transmembrane domain may alter receptor localization and trafficking in the cell. SNPs in the cytoplasmic domain may lead to defects in recruitment of adapter proteins, receptor dimerization and internalization after ligand binding (20, 21). TLR polymorphisms have been linked with susceptibility or protective effect in various diseases, mainly infective and inflammatory, but also in cancer (Table 1) (20).

### 4.1. TLR1, TLR6 and TLR10 polymorphisms

Receptors TLR1, TLR6 and TLR10 have similar structures and genes *TLR1*, *TLR6* and *TLR10* form a single



**Figure 1.** Position of cancer risk associated TLR polymorphisms within genes (A) and proteins (B). A) Polymorphisms (four-angled star) in TLR genes. B) Location of polymorphisms (black triangle) in TLR proteins: ectodomain leucine rich repeats (light grey), transmembrane region (black wavy line) or Toll/interleukin-1 receptor (TIR) domain (dark grey box). After ligand binding all TLRs interact with myeloid differentiation factor 88 (MyD88) except TLR3 who interacts with Toll receptor-associated activator of interferon (TRIF). TLR4 can interact with MyD88 and TRIF. Downstream signaling leads to the activation of transcription factors: activator protein 1 (AP1), nuclear factor  $\kappa$ B (NF- $\kappa$ B) and interferon-regulatory factors (IRFs).

gene cluster on the chromosome 4p14 (22). Receptor TLR1 recognizes triacylated lipoprotein, TLR6 binds diacylated lipoprotein and so far no specific ligand has been linked with TLR10 (7). To bind ligands TLR1 and TLR6 must form dimers with TLR2, while TLR10 homodimerizes or heterodimerizes with TLR1 or TLR2. There have been several association studies between SNPs in *TLR10-TLR1-TLR6* gene cluster and prostate cancer

risk, but with highly contradictory results regarding even the same investigated polymorphisms (23, 24, 25). Finally, pooled analysis on genotype data from three case control studies with more than 3,000 prostate cancer patients and more than 2,500 controls of European ancestry found no overall association between polymorphisms in *TLR10-TLR1-TLR6* gene cluster or *TLR4* gene with prostate cancer risk (26). However, polymorphisms in *TLR10-*

**TABLE 1**  
The association of polymorphisms in *TLR* genes with cancer risk.

Polymorphism	Gene (protein) region	Cancer type	Association*	Sample size, population	References
<b><i>TLR10-TLR1-TLR6</i></b>					
rs483103 (C>A)	In proximity of <i>TLR1</i>	Non-Hodgkin's lymphoma	Negative for allele A	1,946 patients, 1,808 controls, Caucasian	(28)
rs7696175 (C>T)	In proximity of <i>TLR1</i> and <i>TLR6</i>	Breast cancer	Positive for allele T	742 patients, 658 controls, African American; 1,230 patients, 1,118 controls Caucasian	(30)
rs10008492 (C>T)	In proximity of <i>TLR10</i>	Non-Hodgkin's lymphoma	Positive for allele T	1,946 patients, 1,808 controls, Caucasian	(28)
<b><i>TLR6</i></b>					
rs743815 (c.1280 T>C, Val427Ala)	Exon 2 (LRR)	Non-Hodgkin's lymphoma	Positive for allele C	458 patients, 484 controls, Caucasian	(27)
<b><i>TLR10</i></b>					
rs11466657 (Ex3-1099 T>C (reverse strand), Ile473Thr)	Exon 3 (LRR)	Meningioma	Negative for heterozygous genotype AG	101 patients, 330 controls, Caucasian (U.S.)	(29)
<b><i>TLR2</i></b>					
		Gastric cancer	Positive for 'del' allele	174 gastric cancer patients, 208 chronic gastritis patients, 225 controls, Brazilian Caucasian	(34)
		HCV associated hepatocellular carcinoma	Positive for 'del' allele	189 HCV associated hepatocellular carcinoma patients, 192 HCV infected patients, 347 controls, Germany	(35)
c. -196 to -174 del ( $\Delta$ 22nt)	Exon 1 (5'UTR of <i>TLR2</i> )	Breast cancer	Positive for 'del' allele	261 patient, 480 controls, Greece	(36)
		Prostate cancer	Positive for 'del' allele	195 patients, 250 controls, North India	(37)
		Gallbladder cancer	Positive for 'del' allele	230 patients, 230 controls, North India	(38)
		Cervical cancer	Positive for 'del' allele	150 patients, 150 controls, North India	(39)
	Intron 2	Colorectal cancer	Negative for 'M' allele, positive for 'S' allele and 'L' allele	89 patients, 88 controls, Croatia	(40)
Number of GT microsatellite repeats					
rs3804099 (c.597 T>C, Asn199Asn)	Exon 3 (LRR)	Hepatocellular carcinoma	Negative for genotype CT	211 patients, 232 controls, China	(41)
rs3804100 (c.1350 T>C, Ser450Ser)	Exon 3 (LRR)	MALT lymphoma	Positive for allele C	1,946 patients, 1,808 controls, Caucasian	(28)
		Hepatocellular carcinoma	Negative for genotype CT	211 patients, 232 controls, China	(41)
<b><i>TLR3</i></b>					
829 A>C	Intron 1	Nasopharyngeal carcinoma	Positive for allele C	434 patients, 512 controls, Southern China	(43)

TLR4					
	Colorectal cancer	Positive for allele G	89 patients, 88 controls, Croatia	(40)	
	Non-cardia gastric cancer	Positive for allele G	45 hypochlorhydria patients, 149 controls; 312 non-cardia gastric cancer patients, 419 controls; 184 non-cardia gastric cancer patients, 123 cardia cancer patients, 211 esophageal cancer, Caucasian population	(45)	
rs4986790 (c.896 A>G, Asp299Gly)	Gastric cancer	Positive for genotype AG	174 gastric cancer patients, 208 chronic gastritis patients, 225 controls, Brazilian Caucasian	(34)	
Exon 4 (LRR)	Gastrointestinal cancer	Positive for allele G	9,463 patients, 10,825 controls	(46)	
	Prostate cancer	Negative for genotype GG	9,463 patients, 10,825 controls	(46)	
	Breast cancer	Positive for allele Gly	261 patient, 480 controls, Greece	(36)	
	MALT lymphoma, Hodgkin's lymphoma	Positive for allele G	710 patients, 710 controls, Germany	(47)	
	Gastrointestinal cancer	Positive for allele T	9,463 patients, 10,825 controls	(46)	
rs4986791 (c.1196 C>T, Thr399Ile)	Nasopharyngeal carcinoma	Positive for allele T	236 patients, 287 controls, China	(49)	
	Gallbladder cancer	Positive for allele T	230 patients, 230 controls, North India	(48)	
11350 G>C	Nasopharyngeal carcinoma	Positive for genotype GC	486 patients, 529 controls, China	(50)	
rs1927911 (T>C)	Prostate cancer	Positive for allele C	157 patients, 143 controls, Korea	(51)	
rs10759931 (G>A)	Prostate cancer	Positive for genotype GG	157 patients, 143 controls, Korea	(51)	
rs10759932 (T>C)	Gastric cancer	Negative for allele C	1,053 patients, 1,100 controls, China	(52)	
<b>TLR5</b>					
rs5744174 (c.1846 T>C, Phe616Leu)	Gastric cancer	Positive for allele C	248 gastric cancer patients, 846 gastric dysplasia and metaplasia patients, 496 controls with gastritis, China	(33)	
<b>TLR9</b>					
rs5743836 (-1237 T>C)	Hodgkin's lymphoma	Positive for allele C	90 patients, 92 controls, Greece	(54)	
rs187084 (-1486 T>C)	Cervical cancer	Positive for allele C	426 patients, 460 controls, Poland	(55)	
rs352140 (2848 A>G, Pro545Pro)	Hodgkin's lymphoma	Positive for allele A	90 patients, 92 controls, Greece	(54)	
rs352140 (2848 C>T, Pro545Pro)	Cervical cancer	Positive for genotypes CT and TT	426 patients, 460 controls, Poland	(55)	

\*Negative association: polymorphism is linked with decreased cancer risk (odds ratio (OR) < 1)

\*Positive association: polymorphism is linked with increased cancer risk (OR > 1)

5'UTR: 5' untranslated region

3'UTR: 3' untranslated region

LRR: leucine rich repeats

LRRCT: leucine rich repeats C-terminal domain

HCV: hepatitis C virus

MALT: mucosa associated lymphoid tissue

'L' allele: 26 – 43 GT repeats in intron 2 of TLR2 (40)

'M' allele: 19 – 25 GT repeats (40)

'S' allele: < 18 GT repeats (40)



-*TLR1-TLR6* gene cluster have been associated with lymphoma, meningioma and breast cancer risk. *TLR6* polymorphism rs5743815 (c.1280T>C, Val427Ala) was found to be associated with significantly increased non-Hodgkin's lymphoma risk (27). A polymorphism close to *TLR10* gene, rs10008492 (C>T), was associated with significantly increased risk of non-Hodgkin's lymphoma, while polymorphism close to *TLR1* gene, rs4833103 (C>A), was associated with decreased risk of this tumor type development (28). Both SNPs are in linkage disequilibrium with a *TLR1* polymorphism rs5743618 (c.1805G>T, Ser602Ile) which is known to affect ligand binding (28). *TLR10* polymorphism rs11466657 (Ex3-1099T>C, reverse strand, Ile473Thr) was associated with decreased meningioma risk (29) and polymorphism between genes *TLR1* and *TLR6*, rs7696175 (C>T), with increased breast cancer risk in African American population (30).

## 4.2. *TLR2* gene polymorphisms

### 4.2.1. *TLR2* -196 to -174 del polymorphism

*TLR2* gene maps to 4q32 and *TLR2* receptor binds lipoproteins, peptidoglycans and glycoproteins (22). Special attention has been focused on *TLR2* -196 to -174 del ( $\Delta 22$ ) functional polymorphism in 5' untranslated region known to reduce *TLR2* transcriptional activity. There have been several studies regarding impact of  $\Delta 22$  polymorphism on gastric cancer development, but with contradictory results, especially in Japanese and Chinese populations (31, 32, 33). Recently,  $\Delta 22$  polymorphism was associated with increased gastric cancer risk in Brazilian population (34) and with higher hepatitis C viral load and was more frequent in hepatitis C associated hepatocellular carcinoma patients than controls in Germany (35). Also,  $\Delta 22$  polymorphism was found to be more frequent in breast cancer patients than in healthy controls in Greece (36). It was also associated with increased prostate (37), gallbladder (38) and cervical cancer risk (39) in India.

### 4.2.2. Other *TLR2* polymorphisms

Microsatellite GT number repeat polymorphism is located in intron 2 of *TLR2* gene. Middle (M) size 'allele' with 20 and 21 GT repeats is found to be associated with decreased colorectal cancer risk in Croatian population when compared with shorter (S) and larger (L) number of repeats (40), suggesting possible impact of polymorphism on receptor transcriptional activity. Also, *TLR2* polymorphism rs3804100 (c.1350T>C, Ser450Ser) was associated with increased mucosa associated lymphoid tissue (MALT) lymphoma risk (28), and together with rs3804099 (c.597T>C, Asn199Asn) with decreased hepatocellular carcinoma risk (41), but not with breast cancer risk (42).

## 4.3. *TLR3* gene polymorphisms

*TLR3* maps to 4q35 and binds double stranded RNA (22). There have been few reports of *TLR3* polymorphisms in cancer and mainly with no cancer risk association. Those which were proven to be associated in-

clude: (1) *TLR3* polymorphism 829A>C, associated with increased nasopharyngeal carcinoma risk in Chinese population (43), and (2) *TLR3* polymorphism rs3775291 (c.1234 C>T, Leu412Phe) which was implicated as a negative prognostic factor for survival of stage II colorectal cancer patients (44).

## 4.4. *TLR4* gene polymorphisms

*TLR4* gene maps to 9q32-33 and *TLR4* receptor binds lipopolysaccharide (22). *TLR4* polymorphisms have been investigated in numerous studies and in various cancer types, with main focus on polymorphisms rs4986790 (c.896A>G, Asp299Gly) and rs4986791 (c.1196 C>T, Thr399Ile). Both polymorphisms lead to amino acid substitution in extracellular domain of the *TLR4* receptor and are known to diminish response to lipopolysaccharide. Polymorphism Asp299Gly was found to be more frequent in colorectal cancer patients than healthy controls (40). It was also later associated with increased non-cardia gastric cancer risk, but not with gastric cardia carcinoma or esophageal cancer (45). Recently, association with increased gastric cancer risk was confirmed in Brazilian population (34) and meta analysis based on data from 22 studies confirmed association with increased gastrointestinal cancer risk, but with decreased prostate cancer risk (46). Also, Asp299Gly was more frequent in breast cancer patients than healthy controls in Greece (36). On the contrary, no such association was found in breast cancer patients from Croatian population (42). Therefore it is still under the question whether this polymorphism could be used as a marker for breast cancer susceptibility. Polymorphism Asp299Gly was also associated with increased MALT lymphoma and Hodgkin's lymphoma risk (47), but not with non-Hodgkin's lymphoma risk (28). Interestingly, Asp299Gly was associated with improved survival in melanoma patients (48). Polymorphism Thr399Ile was found to be associated with increased gastrointestinal cancer risk and increased overall cancer risk in recent meta analysis (46), but there have also been reports of no association with gastric cancer (34), possibly due to small sample size. Also, polymorphism Thr399Ile was associated with increased nasopharyngeal carcinoma risk in China (49) and increased gallbladder cancer risk in India (38).

### 4.4.1. Other *TLR4* polymorphisms

There have been reports regarding several *TLR4* polymorphisms and cancer risk in China. *TLR4* polymorphism 11350G>C was associated with increased nasopharyngeal carcinoma risk (50). rs1927911 (T>C) and rs10759931 (G>A) were associated with increased prostate cancer risk (51) and rs10759932 (T>C) with reduced gastric cancer risk (52).

## 4.5. *TLR5* gene polymorphisms

*TLR5* gene maps to 1q33.3 and *TLR5* receptor binds flagellin (22). *TLR5* polymorphisms have mainly not been associated with cancer risk. However, *TLR5* polymorphism rs5744174 (c.1846T>C, Phe616Leu), which

affects ligand binding, was recently associated with increased gastric cancer risk in China, with especially high risk in patients infected with *H. pylori* (33).

#### 4.6. TLR7 and TLR8 gene polymorphisms

*TLR7* and *TLR8* share a common locus on chromosome Xp22 and their receptors bind single stranded RNA (22). So far polymorphisms in *TLR7* and *TLR8* have not been firmly associated with cancer risk.

#### 4.7. TLR9 gene polymorphisms

*TLR9* gene maps to 3p21.3, and *TLR9* receptor binds CpG rich bacterial DNA (22). Recently, it was reported that *TLR9* polymorphism Arg892Trp causes hyporesponsiveness to CpG oligodeoxynucleotides (53). *TLR9* polymorphism rs5743836 (-1237T>C) was associated with significantly increased Hodgkin's lymphoma risk, as well as A allele of rs352140 (2848A>G, Pro545Pro) (54). Recently, *TLR9* rs187084 (-1486T>C) and rs352140 (2848C>T, Pro545Pro) polymorphisms were found to be associated with increased cervical cancer risk in Poland (55).

### 5. CONCLUSION

Toll-like receptors have numerous important roles in innate and acquired immunity. Wide spectrum of their signaling pathways and downstream effectors makes them crucial in development of infective diseases, chronic inflammation and cancer. Polymorphisms in *TLR* genes can reduce receptor function, cause over excessive signaling or change the level of transcription, and therefore influence disease development and progression. Although there have been numerous studies regarding impact of polymorphisms on TLR function and disease development, there is a lot of controversial and yet uncertain data. Contradictory results in different studies may reflect insufficient sample size, ethnicity or age differences, infection with undetected pathogens or other environmental factors. The role of *TLR10-TLR1-TLR6* polymorphisms in prostate cancer is controversial and so far those polymorphisms cannot be firmly associated with prostate cancer risk.

Epstein-Bar virus (EBV) infection plays important role in development of non-Hodgkin's lymphoma as well as nasopharyngeal carcinoma. The association of polymorphisms in *TLR* genes with increased cancer risk has been shown in lymphoma (*TLR6* rs5743815) and nasopharyngeal carcinoma (*TLR3* 829A>C and *TLR4* rs4986791). This may affect virus recognition and immune response to pathogens leading to increased disease susceptibility.

Discrepancies regarding *TLR2* -196 to -174 del polymorphism and gastric cancer risk are similar to those in *TLR4* (rs4986790 and rs4986791). Both receptors are important in *H. pylori* infection, which has been implicated in gastric cancer development, and it is possible that they influence cancer development only in presence

of *H. pylori* infection. Conflicting outcome may also result from different types of cancer investigated, only noncardia gastric cancer or all gastric cancer types. Despite somewhat contradictory results, *TLR2* -196 to -174 del polymorphism and *TLR4* rs4986790 and rs4986791 have been linked with various cancer types (gastric, hepatocellular, breast and prostate cancer) in various populations (Japanese, European, Indian) and their role in carcinogenesis and modulation of immune response to pathogens should be further investigated.

Little and less is known about effects of polymorphisms in *TLR3*, *TLR5*, *TLR7* and *TLR8* genes on cancer risk and development. *TLR3* polymorphism 829A>C has been implicated in nasopharyngeal carcinoma risk and *TLR5* polymorphism rs5744174 in gastric cancer risk, but possible impact of *TLR7* and *TLR8* polymorphisms on cancer risk still remains a mystery.

*Acknowledgement:* This work has been funded by the grant number 098-0982464-2394 of the Ministry of Science, Education and Sports, Republic of Croatia.

### REFERENCES

1. ROCK F L, HARDIMAN G, TIMANS J C, KASTELEIN R A, BAZAN J F 1998 A family of human receptors structurally related to *Drosophila* Toll. *Proc Natl Acad Sci U S A* 95: 588-593
2. KAWAI T, AKIRA S 2010 The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol* 11: 373-384
3. ANDERSON K V, JÜRGENS G, NÜSSEIN-VOLHARD C 1985 Establishment of dorsal-ventral polarity in the *Drosophila* embryo: genetic studies on the role of the Toll gene product. *Cell* 42: 779-789
4. ARMANT M A, FENTON M J 2002 Toll-like receptors: a family of pattern recognition receptors in mammals. *Genome Biol* 3: reviews 3011.1- reviews 3011.6
5. MEDZHITOV R, PRESTON HURLBURT P, JANEWAY C A JR 1997 A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature* 388: 394-397
6. HUYNH T, ROSSJOHN J, WILCE M 2007 Toll-like receptors: structural pieces of a curve-shaped puzzle. *Immunol Cell Biol* 85: 406-410
7. YAMAMOTO M, TAKEDA K, AKIRA S 2004 TIR domain-containing adaptors define the specificity of TLR signaling. *Mol Immunol* 40: 861-868
8. MATIJEVIC T, MARJANOVIC M, PAVELIC J 2009 Functionally active toll-like receptor 3 on human primary and metastatic cancer cells. *Scand J Immunol* 70: 18-24
9. YU L, CHEN S 2008 Toll-like receptors expressed in tumor cells: targets for therapy. *Cancer Immunol Immunother* 57: 1271-1278
10. HUANG B, ZHAO J, UNKELESS J C, FENG Z H, XIONG H 2008 TLR signaling by tumor and immune cells: a double-edged sword. *Oncogene* 27: 218-224
11. MATIJEVIC T, PAVELIC J 2010 Toll-like receptors: cost or benefit for cancer? *Curr Pharm Des* 16: 1081-1090
12. MATIJEVIC T, KIRINEC G, PAVELIC J 2011 Antitumor activity from the combined application of poly(I:C) and chemotherapeutics in human metastatic pharyngeal cell lines. *Chemotherapy* 57: 460-467
13. GRIVENNIKOV S I, GRETEN F R, KARIN M 2010 Immunity, inflammation, and cancer. *Cell* 140: 883-899
14. SCHMAUSSER B, ANDRULIS M, ENDRICH S, MÜLLER-HERMELINK H K, ECK M 2005 Toll-like receptors TLR4, TLR5 and TLR9 on gastric carcinoma cells: an implication for interaction with *Helicobacter pylori*. *Int J Med Microbiol* 295: 179-185
15. HE W, LIU Q, WANG L, CHEN W, LI N, CAO X 2007 TLR4 signaling promotes immune escape of human lung cancer cells by

- inducing immunosuppressive cytokines and apoptosis resistance. *Mol Immunol* 44: 2850-2859
16. YANG H, ZHOU H, FENG P, ZHOU X, WEN H, XIE X, SHEN H, ZHU X 2010 Reduced expression of Toll-like receptor 4 inhibits human breast cancer cells proliferation and inflammatory cytokines secretion. *J Exp Clin Cancer Res* 29: 92
  17. SZAJNIK M, SZCZEPANSKI M J, CZYSTOWSKA M, ELI-SHAEV E, MANDAPATHIL M, NOWAK-MARKWITZ E, SPACZYNSKI M, WHITESIDE T L 2009 TLR4 signaling induced by lipopolysaccharide or paclitaxel regulates tumor survival and chemoresistance in ovarian cancer. *Oncogene* 28: 4353-4363
  18. NIHON-YANAGI Y, TERAI K, MURANO T, MATSUMOTO T, OKAZUMI S 2012 Tissue expression of Toll-like receptors 2 and 4 in sporadic human colorectal cancer. *Cancer Immunol Immunother* 61: 71-77
  19. ALLALI-HASSANI A, WASNEY G A, CHAU I, HONG B S, SENISTERRA G, LOPPNAU P, SHI Z, MOULT J, EDWARDS A M, ARROWSMITH C H, PARK H W, SCHAPIRA M, VEDADI M 2009 A survey of proteins encoded by non-synonymous single nucleotide polymorphisms reveals a significant fraction with altered stability and activity. *Biochem J* 424: 15-26
  20. JOHNSON C M, LYLE E A, OMUETI K O, STEPENSKY V A, YEGIN O, ALPSOY E, HAMANN L, SCHUMANN R R, TAPPING R I 2007 Cutting edge: A common polymorphism impairs cell surface trafficking and functional responses of TLR1 but protects against leprosy. *J Immunol* 178: 7520-7524
  21. ETOKEBE G E, SKJELDAL F, NILSEN N, RODIONOV D, KNEZEVIC J, BULAT-KARDUM L, ESPEVIK T, BAKKE O, DEMBIC Z 2010 Toll-like receptor 2 (P631H) mutant impairs membrane internalization and is a dominant negative allele. *Scand J Immunol* 71: 369-381
  22. TAKEDA K, KAISHO T, AKIRA S 2003 Toll-like receptors. *Annu Rev Immunol* 21: 335-376
  23. SUN J, WIKLUND F, ZHENG S L, CHANG B, BÄLTER K, LI L, JOHANSSON J E, LI G, ADAMI H O, LIU W, TOLIN A, TURNER A R, MEYERS D A, ISAACS W B, XU J, GRÖNBERG H 2005 Sequence variants in Toll-like receptor gene cluster (TLR6-TLR1-TLR10) and prostate cancer risk. *J Natl Cancer Inst* 97: 525-532
  24. CHEN Y C, GIOVANNUCCI E, KRAFT P, LAZARUS R, HUNTER D J 2007 Association between Toll-like receptor gene cluster (TLR6, TLR1, and TLR10) and prostate cancer. *Cancer Epidemiol Biomarkers Prev* 16: 1982-1989
  25. STEVENS V L, HSING A W, TALBOT J T, ZHENG S L, SUN J, CHEN J, THUN M J, XU J, CALLE E E, RODRIGUEZ C 2008 Genetic variation in the toll-like receptor gene cluster (TLR10-TLR1-TLR6) and prostate cancer risk. *Int J Cancer* 123: 2644-2650
  26. LINDSTRÖM S, HUNTER D J, GRÖNBERG H, STATTIN P, WIKLUND F, XU J, CHANOCK S J, HAYES R, KRAFT P 2010 Sequence variants in the TLR4 and TLR6-1-10 genes and prostate cancer risk. Results based on pooled analysis from three independent studies. *Cancer Epidemiol Biomarkers Prev* 19: 873-876
  27. CERHAN J R, ANSELL S M, FREDERICKSEN Z S, KAY N E, LIEBOW M, CALL T G, DOGAN A, CUNNINGHAM J M, WANG A H, LIU-MARES W, MACON W R, JELINEK D, WITZIG T E, HABERMANN T M, SLAGER S L 2007 Genetic variation in 1253 immune and inflammation genes and risk of non-Hodgkin lymphoma. *Blood* 110: 4455-4463
  28. PURDUE M P, LAN Q, WANG S S, KRICKER A, MENASHE I, ZHENG T Z, HARTGE P, GRULICH A E, ZHANG Y, MORTON L M, VAJDIC C M, HOLFORD T R, SEVERSON R K, LEADERER B P, CERHAN J R, YEAGER M, COZEN W, JACOBS K, DAVIS S, ROTHMAN N, CHANOCK S J, CHATTERJEE N, ARMSTRONG B K 2009 A pooled investigation of Toll-like receptor gene variants and risk of non-Hodgkin lymphoma. *Carcinogenesis* 30: 275-281
  29. RAJARAMAN P, BRENNER A V, NETA G, PFEIFFER R, WANG S S, YEAGER M, THOMAS G, FINE H A, LINET M S, ROTHMAN N, CHANOCK S J, INSKIP P D 2010 Risk of meningioma and common variation in genes related to innate immunity. *Cancer Epidemiol Biomarkers Prev* 19: 1356-1361
  30. BARNHOLTZ-SLOAN J S, SHETTY P B, GUAN X, NYANTE S J, LUO J, BRENNAN D J, MILLIKAN R C 2010 *FGFR2* and other loci identified in genome-wide association studies are associated with breast cancer in African-American and younger women. *Carcinogenesis* 31: 1417-1423
  31. TAHARA T, ARISAWA T, WANG F, SHIBATA T, NAKAMURA M, SAKATA M, HIRATA I, NAKANO H 2007 Toll-like receptor 2 -196 to 174del polymorphism influences the susceptibility of Japanese people to gastric cancer. *Cancer Sci* 98: 1790-1794
  32. HISHIDA A, MATSUO K, GOTO Y, NAITO M, WAKAI K, TAJIMA K, HAMAJIMA N 2010 No associations of Toll-like receptor 2 (TLR2) -196 to -174del polymorphism with the risk of *Helicobacter pylori* seropositivity, gastric atrophy, and gastric cancer in Japanese. *Gastric Cancer* 13: 251-257
  33. ZENG H M, PAN K F, ZHANG Y, ZHANG L, MA J L, ZHOU T, SU H J, LI W Q, LI J Y, GERHARD M, CLASSEN M, YOU W C 2011 Genetic variants of toll-like receptor 2 and 5, *Helicobacter pylori* infection, and risk of gastric cancer and its precursors in a Chinese population. *Cancer Epidemiol Biomarkers Prev* 20: 2594-2602
  34. DE OLIVEIRA J G, SILVA A E 2012 Polymorphisms of the *TLR2* and *TLR4* genes are associated with risk of gastric cancer in Brazilian population. *World J Gastroenterol* 18: 1235-1242
  35. NISCHALKE H.-D., COENEN M, BERGER C, ALDENHOFF K, MÜLLER T, BERG T, KRÄMER B, KÖRNER C, ODENTHAL M, SCHULZE F, GRÜNHAGE F, NATTERMANN J, SAUERBRUCH T, SPENGLER U 2012 The toll-like receptor 2 (TLR2) -196 to -174 del/ins polymorphism affects viral loads and susceptibility to hepatocellular carcinoma in chronic hepatitis C. *Int J Cancer* 130: 1470-1475
  36. THEODOROPOULOS G E, SARIDAKIS V, KARANTANOS T, MICHALOPOULOS N V, ZAGOURI F, KONTOGIANNI P, LYMPERI M, GAZOULI M, ZOGRAFOS G C 2012 Toll-like receptors gene polymorphisms may confer increased susceptibility to breast cancer development. *Breast* 21: 534-538
  37. MANDAL R K, GEORGE G P, MITTAL R D 2012 Association of Toll-like receptor (TLR) 2, 3 and 9 genes polymorphism with prostate cancer risk in North Indian population. *Mol Biol Rep* 39: 7263-7269
  38. SRIVASTAVA K, SRIVASTAVA A, KUMAR A, MITTAL B 2011 Gallbladder cancer predisposition: a multigenic approach to DNA-repair, apoptotic and inflammatory pathway genes. *PLoS ONE* 6: e16449
  39. PANDEY S, MITTAL R D, SRIVASTAVA M, SRIVASTAVA K, SINGH S, SRIVASTAVA S, MITTAL B 2009 Impact of Toll-like receptors  $\delta$ TLR $\delta$  2 (-196 to -174 del) and TLR 4 (Asp299Gly, Thr399Ile) in cervical cancer susceptibility in North Indian women. *Gynecol Oncol* 114: 501-505
  40. BORASKA JELAVIC T, BARISIC M, DRMIC HOFMAN I, BORASKA V, VRDOLJAK E, PERUZOVIC M, HOZO I, PULJIZ Z, TERZIC J 2006 Microsatellite GT polymorphism in the toll-like receptor 2 is associated with colorectal cancer. *Clin Genet* 70: 156-160
  41. JUNJIE X, SONGYAO J, MINMIN S, YANYAN S, BAIYONG S, XIAXING D, JIABIN J, XI Z, HAO C 2012 The association between Toll-like receptor 2 single-nucleotide polymorphisms and hepatocellular carcinoma susceptibility. *BMC Cancer* 12: 57
  42. ETOKEBE G E, KNEZEVIC J, PETRICEVIC B, PAVELIC J, VRBANEC D, DEMBIC Z 2009 Single-nucleotide polymorphisms in genes encoding toll-like receptor -2, -3, -4, and -9 in case-control study with breast cancer. *Genet Test Mol Biomarkers* 13: 729-734
  43. HE J-F, JIA W-H, FAN Q, ZHOU X-X, QIN H-D, SHUGART Y-Y, ZENG Y-X 2007 Genetic polymorphisms of *TLR3* are associated with nasopharyngeal carcinoma risk in Cantonese population. *BMC Cancer* 7: 194
  44. CASTRO F A, FÖRSTIA, BUCH S, KALTHOFF H, KRAUSS C, BAUER M, EGBERTS J, SCHNIEWIND B, BROERING D C, SCHREIBER S, SCHMITT M, HAMPE J, HEMMINCKI K, SCHAFMAYER C 2011 TLR-3 polymorphism is an independent prognostic marker for stage II colorectal cancer. *Eur J Cancer* 47: 1203-1210
  45. HOLD G L, RABKIN C S, CHOW W H, SMITH M G, GAMMON M D, RISCH H A, VAUGHAN T L, MCCOLL K E, LISSOWSKAJ, ZATONSKI W, SCHOENBERG J B, BLOT W J, MOWAT N A, FRAUMENI J F JR, EL-OMAR E M 2007 A functional polymorphism of toll-like receptor 4 gene increases risk of gastric carcinoma and its precursors. *Gastroenterology* 132: 905-912
  46. JING J J, LI M, YUAN Y 2012 Toll-like receptor 4 Asp299Gly and Thr399Ile polymorphisms in cancer: a meta-analysis. *Gene* 499: 237-242
  47. NIETERS A, BECKMANN L, DEEG E, BECKER N 2006 Gene polymorphisms in Toll-like receptors, interleukin-10, and interleukin-10 receptor alpha and lymphoma risk. *Genes Immun* 7: 615-624



48. GAST A, BERMEJO J L, CLAUS R, BRANDT A, WEIRES M, WEBERA, PLASS C, SUCKERA, HEMMINKI K, SCHADEN-DORF D, KUMAR R 2011 Association of inherited variation in Toll-like receptor genes with malignant melanoma susceptibility and survival. *PLoS ONE* 6: e24370
49. YANG Z H, DAI Q, GU Y J, GUO Q X, GONG L 2012 Cytokine and chemokine modification by Toll-like receptor polymorphisms is associated with nasopharyngeal carcinoma. *Cancer Sci* 103: 653-658
50. SONG C, CHEN L Z, ZHANG R H, YU X J, ZENG YX 2006 Functional variant in the 3'-untranslated region of Toll-like receptor 4 is associated with nasopharyngeal carcinoma risk. *Cancer Biol Ther* 5: 1285-1291
51. SONG J, KIM D Y, KIM C S, KIM H J, LEE D H, LEE H M, KO W, LEE G 2009 The association between Toll-like receptor 4 (TLR4) polymorphisms and the risk of prostate cancer in Korean men. *Cancer Genet Cytogenet* 190: 88-92
52. HUANG H, WU J, JIN G, ZHANG H, DING Y, HUA Z, ZHOU Y, XUE Y, LU Y, HU Z, XU Y, SHEN H 2010 A 5'-flanking region polymorphism in toll-like receptor 4 is associated with gastric cancer in a Chinese population. *J Biomed Res* 24: 100-106
53. KNEZEVIC J, PAVLINIC D, ROSE W A 2ND, LEIFER C A, BENDELJA K, GABRILOVAC J, PARCINA M, LAUC G, KUBARENKO A V, PETRICEVIC B, VRBANEC D, BULAT-KARDUM L, BEKEREDJIAN-DING I, PAVELIC J, DEMBIC Z, WEBER A N 2012 Heterozygous carriage of a dysfunctional Toll-like receptor 9 allele affects CpG oligonucleotide responses in B cells. *J Biol Chem* 287: 24544-24553
54. MOLLAKI V, GEORGIADIS T, TASSIDOU A, IOANNOU M, DANIL Z, KOUTSOKERA A, PAPANASSIOU A A, ZINTZARAS E, VASSILOPOULOS G 2009 Polymorphisms and haplotypes in TLR9 and MYD88 are associated with the development of Hodgkin's lymphoma: a candidate-gene association study. *J Hum Genet* 54: 655-659
55. ROSZAK A, LIANERI M, SOWIŃSKA A, JAGODZIŃSKI P P 2012 Involvement of toll-like receptor 9 polymorphism in cervical cancer development. *Mol Biol Rep* 39: 8425-8430

