

# Quantum Twist-Deformed $D = 4$ Phase Spaces with Spin Sector and Hopf Algebroid Structures

Jerzy Lukierski<sup>1</sup>, Stjepan Meljanac<sup>2</sup>, Mariusz Woronowicz<sup>1</sup>

<sup>1</sup> Institute for Theoretical Physics, University of Wrocław,  
pl. Maxa Borna 9, 50-205 Wrocław, Poland

<sup>2</sup>Division of Theoretical Physics, Rudjer Bošković Institute,  
Bijenička c. 54, HR-10002 Zagreb, Croatia

## Abstract

We consider the generalized  $(10 + 10)$ -dimensional  $D = 4$  quantum phase spaces containing translational and Lorentz spin sectors associated with the dual pair of twist-quantized Poincare Hopf algebra  $\mathbb{H}$  and quantum Poincare Hopf group  $\widehat{\mathbb{G}}$ . Two Hopf algebroid structures of generalized phase spaces with spin sector will be investigated: first one  $\mathcal{H}^{(10,10)}$  describing dynamics on quantum group algebra  $\widehat{\mathbb{G}}$  provides by the Heisenberg double algebra  $\mathcal{HD} = \mathbb{H} \rtimes \widehat{\mathbb{G}}$ , and second, denoted by  $\tilde{\mathcal{H}}^{(10,10)}$ , describing twisted Hopf algebroid with base space containing twisted noncommutative Minkowski space  $\hat{x}_\mu$ . We obtain the first explicit example of Hopf algebroid structure of relativistic quantum phase space which contains quantum-deformed Lorentz spin sector.

## 1 Introduction

Following recent description of noncommutative quantum phase spaces as bialgebroids and Hopf algebroids (see [1]-[10]) we shall study in this paper such algebraic structures in quantum phase spaces derived from twisted quantum Poincare symmetries. In comparison with recent studies [5]-[10] the novelty in our approach is the appearance of Hopf algebroid description of  $D = 4$  generalized relativistic quantum-deformed phase spaces with additional coordinates and momenta describing Lorentz spin sector.

The quantum-deformed relativistic phase spaces  $\mathcal{H}^{(4,4)}$ , spanned by the degrees of freedom  $(\hat{x}_\mu, \hat{p}_\mu)$ , are described by quantum deformations of canonical relativistic Heisenberg algebra<sup>1</sup>

<sup>1</sup>We denote the undeformed canonical phase space variables and canonical Poincare algebra generators by letters without hats.

$$[p_\mu, x_\nu] = -i\eta_{\mu\nu}, \quad [x_\mu, x_\nu] = [p_\mu, p_\nu] = 0. \quad (1)$$

Such choice of phase space is suitable only for the description of standard spinless dynamics. In this paper we shall consider the generalized  $D = 4$  quantum phase spaces  $\mathcal{H}^{(10,10)}$  and  $\tilde{\mathcal{H}}^{(10,10)}$ . The additional generators  $\hat{\Lambda}_{\mu\nu}$  will be given by quantum counterpart of six Lorentzian angles ( $\hat{\Lambda}\hat{\Lambda}^T = \hat{\Lambda}^T\hat{\Lambda} = I$ ) dual to the generators  $M_{\mu\nu}$  describing quantum-deformed Lorentz algebra.

The phase space description of spin dynamics as Heisenberg double has its roots in the half century old idea of Souriau [11] and Kostant [12] who described symplectic dynamics of point-like objects by the geometry of cosets  $\mathcal{K} = G/S$ , where  $G$  is the space-time symmetry group, and  $S$  its so-called stability group. If  $G$  is the Poincare group  $\mathcal{P}^4$ , for providing the dynamics of spin degrees of freedom the coset  $\mathcal{K}$  should include besides translations as well some Lorentz group parameters (see e.g. [13]-[17]). In such a way one can describe e.g. the infinite-dimensional spin multiplets by adding to coordinate sector the spinorial Weyl spinor coordinates  $\eta_\alpha, \eta_{\dot{\alpha}} = \bar{\eta}_\alpha (\alpha = 1, 2)$  defined by fourdimensional coset of  $SL(2; \mathbb{C})$ . In this way one can introduce the generalized wave functions (classical fields)  $\Psi(x; \eta_\alpha, \eta_{\dot{\alpha}})$  obtained if  $S$  is a suitable two-dimensional subgroup of Lorentz group<sup>2</sup>.

The Souriau-Kostant approach as well as its generalization to Poisson manifolds [23],[24] and Poisson-Lie theory [25]-[27] can be also extended to models based on noncommutative (NC) geometry. The construction of quantum-deformed NC space-times extended by the additional quantum Lorentz group parameters has been recently also proposed in the case of quantum-deformed Poincare group (see e.g. [28]-[29] where the case  $D = 2 + 1$  was considered).

In this paper we shall study firstly the generalized quantum-deformed phase space  $\mathcal{H}^{(10,10)} = (\hat{\xi}_\mu, \hat{\Lambda}_{\mu\nu}; p_\mu, M_{\mu\nu})$  with noncommutative generalized coordinates described by the algebraic manifold of full quantum  $D = 4$  Poincare group (i.e.  $S = 1$ ), and supplemented with dual algebra of generalized momenta defined by ten generators of quantum-deformed Poincare algebra. The quantization will be obtained by the twisting procedure considered recently in [10] (see also [30]), which generate the Lie-algebraic deformation of space-time translations algebra and the quadratic algebra describing the commutators of all ten quantum Poincare group generators  $(\hat{\xi}_\mu, \hat{\Lambda}_{\mu\nu})$ . Subsequently, following [1] we anticipate the Hopf algebroid structure describing the dynamics on algebraic quantum Poincare group manifold.

Further, following [2],[9] we introduce  $(10 + 10)$ -dimensional quantum phase space  $\tilde{\mathcal{H}}^{(10,10)} = (\hat{x}_\mu, \tilde{\Lambda}_{\mu\nu}; p_\mu, M_{\mu\nu})$  with Hopf algebroid structure generated by twist-deformed  $D = 4$  Poincare-Hopf algebra  $\mathbb{H}$ , with base space described

---

<sup>2</sup>It can be shown ( see e.g. [18],[15]) that for  $D = 4$  the generalized field  $\psi(x; \eta, \bar{\eta})$  can be defined as introduced on 8-dimensional coset  $\mathcal{P}^4/H_2$ , where  $H_2$  is generated by Lorentz generators  $(M_{01} + M_{31}, M_{02} + M_{32})$  and  $SL(2; \mathbb{C})/H_2$  is parametrized by a pair  $(\eta_\alpha, \bar{\eta}_{\dot{\alpha}})$  of complex-conjugated  $D = 4$  Weyl spinors. The idea of studying dynamics on 10-dimensional Poincare group manifold was early advocated by Lurcat [19], [14]. Other way of introducing additional variables describing spin is to supplement space-time by two-parameter auxiliary manifold described by the sphere  $S^2$  [20]-[22].

by  $\mathbb{H}$ -module  $\hat{X}_A = (\hat{x}_\mu, \tilde{\Lambda}_{\mu\nu})$ , ( $A = 1 \dots 10$ ), its noncommutativity structure determined by twist-deformed star product multiplication

$$\hat{X}_A \cdot \hat{X}_B \simeq X_A \star_{\mathcal{F}} X_B = m[\mathcal{F}^{-1} \circ (X_A \otimes X_B)] = (\mathcal{F}_{(1)}^{-1} \triangleright X_A)(\mathcal{F}_{(2)}^{-1} \triangleright X_B) \quad (2)$$

where the Drinfeld twist factor  $\mathcal{F} = \sum \mathcal{F}_{(1)} \otimes \mathcal{F}_{(2)} \in \mathbb{H} \otimes \mathbb{H}$  satisfies the two-cocycle condition [2],[31]. In nondeformed relativistic theory one can identify the translations fourvectors  $\hat{\xi}_\mu$  and Minkowski space-time coordinates  $\hat{x}_\mu$ ; in the presence of quantum deformations the algebraic properties of base spaces  $(\hat{\xi}_\mu, \hat{\Lambda}_{\mu\nu})$  and  $(\hat{x}_\mu, \tilde{\Lambda}_{\mu\nu})$  are usually different<sup>3</sup>.

We shall consider therefore two possible types of Hopf algebroids describing quantum phase spaces which are defined by twist-quantized Hopf algebra  $\mathbb{H}$  (quantum deformed Poincare algebra) defining generalized momenta and generalized coordinates, given by two choices of the  $\mathbb{H}$ -module:

- i) described by Heisenberg double  $\mathcal{HD} = \mathbb{H} \rtimes \hat{\mathbb{G}}$  defined by smash product<sup>4</sup> of dual Hopf algebras  $\mathbb{H}, \hat{\mathbb{G}}$ , with built-in Hopf action  $h \blacktriangleright \hat{g} = \hat{g}_{(1)} \langle h, \hat{g}_{(2)} \rangle$  and  $h \blacktriangleright (\hat{g}\hat{g}') = (h_{(1)} \blacktriangleright \hat{g})(h_{(2)} \blacktriangleright \hat{g}')$  ( $h \in \mathbb{H}; \hat{g}, \hat{g}' \in \hat{\mathbb{G}}; \Delta(h) = h_{(1)} \otimes h_{(2)}$ ). The associative multiplication formula in  $\hat{\mathbb{G}} \otimes \mathbb{H}$  algebra is given by

$$(\hat{g} \otimes h)(\hat{g}' \otimes h') = \hat{g}(h_{(1)} \blacktriangleright \hat{g}') \otimes h_{(2)}h' \quad (3)$$

where  $\hat{\mathbb{G}}$  describes generalized coordinates.

Heisenberg double data are specified by the pair of dual Hopf algebras  $(\mathbb{H}, \hat{\mathbb{G}})$  with the pairing  $\langle \cdot, \cdot \rangle$  (see also <sup>4</sup>), which expresses the duality of  $\mathbb{H}$  and  $\hat{\mathbb{G}}$  by the formulae

$$\langle h, \hat{g}\hat{g}' \rangle = \langle \Delta(h), \hat{g} \otimes \hat{g}' \rangle \quad (4)$$

$$\langle hh', \hat{g} \rangle = \langle h \otimes h', \Delta(\hat{g}) \rangle. \quad (5)$$

Heisenberg doubles provide the associative algebras of quantum phase spaces endowed with quantum-deformed NC symplectic structure (see e.g. [27]), however without the Hopf-algebraic coalgebra sector<sup>5</sup>.

- ii) We introduce the twist-deformed coordinate sector described by  $\hat{X}_A \in \hat{\mathbb{X}}$ , where  $\hat{\mathbb{X}}$  is the  $\mathbb{H}$ -module algebra, however not obtained from  $\mathbb{H}$  via Hopf-algebraic duality<sup>6</sup>. We shall attach to the quantum phase space algebra

<sup>3</sup>In the canonical case of  $\theta_{\mu\nu}$ -deformation this difference is exposed e.g. in [30], [32].

<sup>4</sup>Smash product algebra (see e.g. [33]) is a special kind of cross-product algebra  $\mathbb{H} \rtimes \mathbb{V}$  when the left  $\mathbb{H}$ -module  $\mathbb{V}$  is provided by dual Hopf algebra  $\hat{\mathbb{G}}$ , with the action  $\mathbb{H} \otimes \hat{\mathbb{G}} \rightarrow \hat{\mathbb{G}}$  defined with the help of bilinear pairing  $\langle \cdot, \cdot \rangle : \mathbb{H} \otimes \hat{\mathbb{G}} \rightarrow \mathbb{C}$ . In the derivation of quantum-mechanical Heisenberg algebra (see (1)) as Heisenberg double the  $\mathbb{C}$ -number pairing is assumed to be proportional to  $\hbar$ ; in this paper we shall put  $\hbar = 1$ .

<sup>5</sup>Partial coalgebraic structure can be introduced only separately in generalized coordinate and generalized momenta sectors, described respectively by dual Hopf algebras. If in quantum phase space we introduce the bialgebroid coproducts (see Sect. 4), the group coproducts are changed, but the partial coalgebraic structure in generalized momenta sector can be preserved.

<sup>6</sup>In distinction to the action  $h \blacktriangleright g$ , the action  $h \triangleright X_A$  is described by a white triangle.

$\tilde{\mathcal{H}}^{(10,10)} = (\hat{x}_\mu, \tilde{\Lambda}_{\mu\nu}; p_\mu, M_{\mu\nu})$  the coalgebraic sector and antipodes, which define the twisted Hopf algebroid structure (see [2], [9]). For twisted Hopf algebroid  $\tilde{\mathcal{H}}^{(10,10)}$  the base algebra  $(\hat{\mathbb{X}}, \star_{\mathcal{F}})$  is endowed with the multiplication defined by the star product (2), which describes the generalized NC coordinates  $\hat{X}_A = (\hat{x}_\mu, \tilde{\Lambda}_{\mu\nu})$ , which will describe the Lie-algebraic type of noncommutativity.

The plan of our paper is the following: In Sect. 2 following [10] and [30] we shall describe the pair of dual twist-deformed Poincare-Hopf algebras  $\mathbb{H}, \hat{\mathbb{G}}$ . In Sect. 3 we shall introduce the generalized  $D = 4$  quantum phase spaces  $\mathcal{H}^{(10,10)}$  containing space-time translations generators  $\hat{\xi}_\mu$  as well as  $\tilde{\mathcal{H}}^{(10,10)}$  with the quantum Minkowski space-time coordinates  $\hat{x}_\mu$ . In Sect. 4 by following Xu twisted bialgebroids framework [2] we will show how to get for the quantum phase space  $\tilde{\mathcal{H}}^{(10,10)}$  the explicit Hopf algebroid structure. In Sect. 5 we present final remarks and comments on possible extensions of presented results.

## 2 Dual pair of twisted Hopf algebras

### 2.1 Twist-deformed Poincare algebra $\mathbb{H}$

The classical  $D = 4$  Poincare-Hopf algebra looks as follows

$$[p_\mu, p_\nu] = 0 \quad (6)$$

$$[M_{\mu\nu}, p_\rho] = \eta_{\nu\rho} p_\mu - \eta_{\mu\rho} p_\nu \quad (7)$$

$$[M_{\mu\nu}, M_{\rho\sigma}] = \eta_{\nu\rho} M_{\mu\sigma} - \eta_{\mu\rho} M_{\nu\sigma} - \eta_{\nu\sigma} M_{\mu\rho} + \eta_{\mu\sigma} M_{\nu\rho} \quad (8)$$

where  $\eta_{\mu\nu} = \text{diag}(-1, 1, \dots, 1)$  and

$$\Delta_0(p_\mu) = p_\mu \otimes 1 + 1 \otimes p_\mu \quad \Delta_0(M_{\mu\nu}) = M_{\mu\nu} \otimes 1 + 1 \otimes M_{\mu\nu} \quad (9)$$

$$S_0(p_\mu) = -p_\mu \quad S_0(M_{\mu\nu}) = -M_{\mu\nu} \quad (10)$$

$$\epsilon_0(p_\mu) = 0 \quad \epsilon_0(M_{\mu\nu}) = 0. \quad (11)$$

We define twist  $\mathcal{F}$  as an element of  $\mathbb{H} \otimes \mathbb{H}$  ( $\mathbb{H} = \mathcal{U}(\mathcal{P})$ ) which has an inverse, satisfies the cocycle condition

$$\mathcal{F}_{12} (\Delta_0 \otimes 1) \mathcal{F} = \mathcal{F}_{23} (1 \otimes \Delta_0) \mathcal{F} \quad (12)$$

and the normalization condition

$$(\epsilon \otimes 1) \mathcal{F} = (1 \otimes \epsilon) \mathcal{F} = 1 \quad (13)$$

where  $\mathcal{F}_{12} = \mathcal{F} \otimes 1$  and  $\mathcal{F}_{23} = 1 \otimes \mathcal{F}$ . It is known, that  $\mathcal{F}$  does not modify the algebraic part and the counit, but changes the coproducts and the antipodes in the following way

$$\Delta_{\mathcal{F}}(h) = \mathcal{F} \circ \Delta_0(h) \circ \mathcal{F}^{-1} \quad (14)$$

$$S_{\mathcal{F}}(h) = U S_0(h) U^{-1} \quad (15)$$

where due to (12) the coproduct  $\Delta_{\mathcal{F}}$  is coassociative,  $U = \sum f_{(1)}S(f_{(2)})$  (we use Sweedler's notation  $\mathcal{F} = \sum f_{(1)} \otimes f_{(2)}$ ). In such a way we obtain for any choice of the twist  $\mathcal{F}$  the quasitriangular Hopf algebra.

Our family of Abelian twists is <sup>7</sup>

$$\begin{aligned} \mathcal{F}_u &:= \exp(f_u) = \exp\left(\left(1-u\right)\frac{a \cdot p}{2\kappa} \otimes \theta^{\alpha\beta} M_{\alpha\beta} - u\theta^{\alpha\beta} M_{\alpha\beta} \otimes \frac{a \cdot p}{2\kappa}\right) \\ &= \exp\left(\frac{1-u}{2}\mathcal{K}^{\alpha\beta} \otimes M_{\alpha\beta} - \frac{u}{2}M_{\alpha\beta} \otimes \mathcal{K}^{\alpha\beta}\right) \end{aligned} \quad (16)$$

where  $a \cdot p = a^\mu \eta_{\mu\nu} p^\nu$ , The parameter  $u \in [0, 1]$ ,  $\kappa$  is the deformation parameter with dimension of mass,  $a^2 \equiv a_\mu a^\mu \in \{-1, 0, 1\}$ ,

$$\mathcal{K}^{\mu\nu} = \frac{a \cdot p}{\kappa} \theta^{\mu\nu} \quad (17)$$

and further

$$\theta^{\mu\nu} = -\theta^{\nu\mu}, \quad a_\mu \theta^{\mu\nu} = 0. \quad (18)$$

From twists (16) one gets the coproducts

$$\Delta_{\mathcal{F}}(p_\mu) = p_\alpha \otimes (e^{-u\mathcal{K}})^\alpha_\mu + (e^{(1-u)\mathcal{K}})^\alpha_\mu \otimes p_\alpha \quad (19)$$

$$\begin{aligned} \Delta_{\mathcal{F}}(M_{\mu\nu}) &= M_{\alpha\beta} \otimes (e^{-u\mathcal{K}})^\alpha_\mu (e^{-u\mathcal{K}})^\beta_\nu + (e^{(1-u)\mathcal{K}})^\alpha_\mu (e^{(1-u)\mathcal{K}})^\beta_\nu \otimes M_{\alpha\beta} \\ &\quad + \frac{\theta^{\alpha\beta}}{2\kappa} (a_\mu \delta_\nu^\gamma - a_\nu \delta_\mu^\gamma) \left[ (1-u)p_\delta \otimes M_{\alpha\beta} (e^{-u\mathcal{K}})^\delta_\gamma - uM_{\alpha\beta} (e^{(1-u)\mathcal{K}})^\delta_\gamma \otimes p_\delta \right] \end{aligned} \quad (20)$$

where  $\mathcal{K}_{\mu\nu}$  is given by equation (17), and corresponding antipodes are:

$$S_{\mathcal{F}}(p_\mu) = -(e^{-(1-2u)\mathcal{K}})^\alpha_\mu p_\alpha \quad (21)$$

$$\begin{aligned} S_{\mathcal{F}}(M_{\mu\nu}) &= -M_{\alpha\beta} (e^{-(1-2u)\mathcal{K}})^\alpha_\mu (e^{-(1-2u)\mathcal{K}})^\beta_\nu \\ &\quad + \frac{1}{\kappa} (a_\mu \delta_\nu^\alpha - a_\nu \delta_\mu^\alpha) [S(p_\alpha) + (1-2u)\theta_{\alpha\beta} S(p^\beta)] \end{aligned} \quad (22)$$

The counit is trivial:

$$\epsilon(p_\mu) = 0, \quad \epsilon(M_{\mu\nu}) = 0. \quad (23)$$

## 2.2 RTT matrix quantum Poincare group $\widehat{\mathbb{G}}$

The universal  $\mathcal{R}$ -matrix ( $(a \wedge b = a \otimes b - b \otimes a)$ )

$$\mathcal{R} = \mathcal{F}_u^T \mathcal{F}_u^{-1} = \exp\left[\frac{1}{2}(M_{\alpha\beta} \wedge \mathcal{K}^{\alpha\beta})\right] \quad (a \otimes b)^T = b \otimes a \quad (24)$$

---

<sup>7</sup>See [30] and [10]; the twists corresponding to special cases  $u = 0$  and  $u = \frac{1}{2}$  were considered also respectively in [35] and [36]. The twists (16) with different  $u$  can be related by coboundary twist (see e.g. [34], Sect. 3.1) which does not modify the universal  $\mathcal{R}$ -matrix.

can be used for the description of 10-generator deformed  $D = 4$  Poincaré group. Using the  $5 \times 5$  - matrix realization of the Poincaré generators

$$(M_{\mu\nu})^A_B = \delta^A_\mu \eta_{\nu B} - \delta^A_\nu \eta_{\mu B} \quad (P_\mu)^A_B = -i\delta^A_\mu \delta^4_B \quad (25)$$

we can show that in (24) only the linear term is non-vanishing

$$\mathcal{R} = 1 \otimes 1 + \frac{1}{2}(M_{\alpha\beta} \wedge \mathcal{K}^{\alpha\beta}). \quad (26)$$

To find the matrix quantum group which is dual to our Hopf algebra  $\mathbb{H}$  we introduce the following  $5 \times 5$  - matrices

$$\hat{\mathcal{T}}_{AB} = \begin{pmatrix} \hat{\Lambda}_{\mu\nu} & \hat{\xi}_\mu \\ 0 & 1 \end{pmatrix} \quad (27)$$

where  $\hat{\Lambda}_{\mu\nu}$  parametrizes the quantum Lorentz rotation and  $\hat{\xi}_\mu$  denotes quantum translations. In the framework of the  $\mathcal{FRT}$  procedure [37], the algebraic relations defining such a quantum group  $\hat{\mathbb{G}}$  is described by the following relation

$$\mathcal{R}\hat{\mathcal{T}}_1\hat{\mathcal{T}}_2 = \hat{\mathcal{T}}_2\hat{\mathcal{T}}_1\mathcal{R} \quad (28)$$

while the composition law for the coproduct remains classical

$$\Delta(\hat{\mathcal{T}}_{AB}) = \hat{\mathcal{T}}_{AC} \otimes \hat{\mathcal{T}}_B^C \quad (29)$$

with  $\hat{\mathcal{T}}_1 = \hat{\mathcal{T}} \otimes 1$ ,  $\hat{\mathcal{T}}_2 = 1 \otimes \hat{\mathcal{T}}$  and quantum  $\mathcal{R}$ -matrix given in the representation (25).

In terms of the operator basis  $(\hat{\Lambda}_{\mu\nu}, \hat{\xi}_\mu)$  the algebraic relations (28) describing quantum group algebra can be written as follows

$$[\hat{\xi}_\mu, \hat{\xi}_\nu] = \frac{i}{\kappa}(a_\nu \theta_\mu^\alpha - a_\mu \theta_\nu^\alpha) \hat{\xi}_\alpha \quad (30)$$

$$[\hat{\xi}_\mu, \hat{\Lambda}_{\rho\sigma}] = \frac{i}{\kappa}(\hat{\Lambda}_{\rho\alpha} \theta_\sigma^\alpha a^\gamma \hat{\Lambda}_{\mu\gamma} - a_\mu \theta_\rho^\alpha \hat{\Lambda}_{\alpha\sigma}) \quad (31)$$

$$[\hat{\Lambda}_{\mu\nu}, \hat{\Lambda}_{\rho\sigma}] = 0 \quad (32)$$

while the coproduct (29) takes the well known primitive forms

$$\Delta(\hat{\Lambda}_{\mu\nu}) = \hat{\Lambda}_{\mu\rho} \otimes \hat{\Lambda}_\nu^\rho \quad \Delta(\hat{\xi}_\mu) = \hat{\Lambda}_{\mu\nu} \otimes \hat{\xi}^\nu + \hat{\xi}_\mu \otimes 1. \quad (33)$$

One can check that coproducts (33) are consistent with the commutators (30)-(32).

### 2.3 Duality between quantum algebra $\mathbb{H}$ and quantum group $\widehat{\mathbb{G}}$

Two Hopf algebras  $\mathbb{H}, \widehat{\mathbb{G}}$  are said to be dual if there exists a nondegenerate bilinear form

$$\langle \cdot, \cdot \rangle : \mathbb{H} \times \widehat{\mathbb{G}} \longrightarrow \mathbb{C} \quad (h, \hat{g}) \longrightarrow \langle h, \hat{g} \rangle \quad (34)$$

such that the duality relations (4)-(5) are satisfied. One can check, using the following pairing relations

$$\langle p_\mu, \hat{\xi}_\nu \rangle = -i\eta_{\mu\nu} \quad \langle M_{\mu\nu}, \hat{\Lambda}_{\alpha\beta} \rangle = -(\eta_{\mu\alpha}\eta_{\nu\beta} - \eta_{\nu\alpha}\eta_{\mu\beta}) \quad \langle 1, \hat{\Lambda}_{\mu\nu} \rangle = \eta_{\mu\nu} \quad (35)$$

that we have in our case  $(h, h' = \{M_{\mu\nu}, p_\mu\}; \hat{g}, \hat{g}' = \{\hat{\Lambda}_{\mu\nu}, \hat{\xi}_\mu\})$

$$\langle h, [\hat{g}, \hat{g}'] \rangle = \langle \Delta(h), \hat{g} \otimes \hat{g}' - \hat{g}' \otimes \hat{g} \rangle \quad (36)$$

$$\langle [h, h'], \hat{g} \rangle = \langle h \otimes h' - h' \otimes h, \Delta(\hat{g}) \rangle. \quad (37)$$

The basic action of  $\mathbb{H}$  on  $\widehat{\mathbb{G}}$  promoting  $\widehat{\mathbb{G}}$  to the  $\mathbb{H}$ -module is given by the relation

$$h \blacktriangleright \hat{g} = \hat{g}_{(1)} \langle h, \hat{g}_{(2)} \rangle. \quad (38)$$

After using (4) one gets the relation

$$\begin{aligned} h \blacktriangleright (\hat{g}\hat{g}') &= \hat{g}\hat{g}'_{(1)} \langle \Delta h, \hat{g}_{(2)} \otimes \hat{g}'_{(2)} \rangle = \hat{g}\hat{g}'_{(1)} \langle h_{(1)}, \hat{g}_{(2)} \rangle \langle h_{(1)}, \hat{g}'_{(2)} \rangle \\ &= (h_{(1)} \blacktriangleright \hat{g})(h_{(2)} \blacktriangleright \hat{g}') \end{aligned} \quad (39)$$

what establishes that algebra  $\widehat{\mathbb{G}}$  is the  $\mathbb{H}$ -module.

## 3 Heisenberg double $\mathcal{HD}$ and generalized $D = 4$ quantum phase spaces

Using the formula (3), in Heisenberg double framework we can obtain cross commutators between the algebra  $\mathbb{H}$  and group  $\widehat{\mathbb{G}}$  by the following relation

$$[h, \hat{g}] = \hat{g}_{(2)} \langle h_{(1)}, \hat{g}_{(1)} \rangle h_{(2)} - \hat{g}h \quad h = \{M_{\mu\nu}, p_\mu\}; \hat{g} = \{\hat{\Lambda}_{\mu\nu}, \hat{\xi}_\mu\}. \quad (40)$$

Using pairing (35) and coproducts (19), (20) and (33) we get

$$[p_\mu, \hat{\xi}_\nu] = -i(e^{-u\mathcal{K}})_{\nu\mu} + i(1-u)\frac{a_\nu}{\kappa}\theta_\mu^\alpha p_\alpha \quad (41)$$

$$[p_\mu, \hat{\Lambda}_{\rho\sigma}] = 0 \quad (42)$$

$$\begin{aligned} [M_{\mu\nu}, \hat{\Lambda}_{\rho\sigma}] &= \frac{u}{\kappa}\theta_\rho^\alpha \hat{\Lambda}_{\alpha\sigma}(a_\mu p_\nu - a_\nu p_\mu) \\ &\quad + \hat{\Lambda}_{\alpha\sigma}(e^{-u\mathcal{K}})_{[\mu}^\alpha (e^{-u\mathcal{K}})_{\rho\nu]} \end{aligned} \quad (43)$$

$$\begin{aligned} [M_{\mu\nu}, \hat{\xi}_\rho] &= \hat{\xi}_\gamma (e^{u\mathcal{K}})_{[\mu}^\gamma (e^{u\mathcal{K}})_{\nu]\rho} + \frac{u}{\kappa}\theta_\rho^\alpha \hat{\xi}_\alpha (a_\mu p_\nu - a_\nu p_\mu) \\ &\quad + \frac{i(u-1)}{2\kappa}\theta^{\alpha\beta} M_{\alpha\beta} a_{[\mu} (e^{-u\mathcal{K}})_{\rho\nu]} + \frac{i(u-1)}{\kappa} a_\rho M_{[\mu\alpha} \theta_{\nu]}^\alpha \end{aligned} \quad (44)$$

One can show that the generators of quantum Poincare group algebra, satisfying relations (30)-(32), (41)-(44) can be expressed in terms of classical group parameters  $(\xi_\mu, \Lambda_{\rho\sigma})$  as follows

$$\hat{\xi}_\mu = \xi_\alpha (e^{-u\mathcal{K}})_\mu^\alpha + (1-u) \frac{ia_\mu}{2\kappa} \theta M \quad (45)$$

$$\hat{\Lambda}_{\rho\sigma} = (e^{-u\mathcal{K}})_\rho^\alpha \Lambda_{\alpha\beta} (e^{-\frac{1}{\kappa}(p\Lambda a)^\theta})^\beta_\sigma. \quad (46)$$

The group manifold  $\widehat{\mathbb{G}}$  as  $\mathbb{H}$ -module algebra can be promoted to the base algebra of Hopf algebroid. We get in such a way the application of Lu construction [1], which provides for Heisenberg double  $\mathcal{HD} = \mathbb{H} \rtimes \widehat{\mathbb{G}}$ , with generators  $(\hat{\xi}_\mu, \hat{\Lambda}_{\mu\nu}; p_\mu, M_{\mu\nu})$  the Hopf algebroid structure<sup>8</sup>.

Let us introduce in place of relations (31) the following set of parameter-dependent algebraic relations ( $s$ -real parameter)

$$[\hat{\xi}_\mu^{(s)}, \hat{\Lambda}_{\rho\sigma}^{(s)}] = \frac{i}{\kappa} \left( s \hat{\Lambda}_{\rho\alpha}^{(s)} \theta^\alpha_\sigma a^\gamma \hat{\Lambda}_{\mu\gamma}^{(s)} - a_\mu \theta_\rho^\alpha \hat{\Lambda}_{\alpha\sigma}^{(s)} \right). \quad (47)$$

One can show that the relation (47) forms together with relations (30), (32) the consistent algebra of associative generalized quantum Poincare phase spaces, satisfying Jacobi relations. However, for  $\xi = 0$  and  $\xi = 1$  one can supplement the Hopf algebroid structure, namely

- i) if  $s = 1$  ( $\hat{\xi}_\mu^{(1)} \equiv \hat{\xi}_\mu, \hat{\Lambda}_{\rho\sigma}^{(1)} \equiv \hat{\Lambda}_{\rho\sigma}$ ) one can construct the Hopf algebroid  $\mathcal{H}^{(10,10)}$  with base algebra  $\widehat{\mathbb{G}} = (\hat{\xi}_\mu, \hat{\Lambda}_{\mu\nu})$  described by algebraic quantum Poincare group manifold.
- ii) if  $s = 0$  ( $\hat{\xi}_\mu^{(0)} \equiv \hat{x}_\mu, \hat{\Lambda}_{\rho\sigma}^{(0)} \equiv \tilde{\Lambda}_{\rho\sigma}$ ) one gets alternative Hopf algebroid, denoted by  $\tilde{\mathcal{H}}^{(10,10)}$ , with algebra  $\tilde{\mathbb{X}} = (\hat{x}_\mu, \tilde{\Lambda}_{\mu\nu})$  as base algebra, with the multiplication given by the star product formula (2). The formula (2) can be also written as follows

$$f(X) \star_{\mathcal{F}} k(X') = \widehat{f(\hat{X})} \triangleright k(X') \quad (48)$$

where  $\widehat{f(\hat{X})}$  denotes the noncommutative representation of  $f(\hat{X})$  defined as follows

$$f(\hat{X}) \simeq \widehat{f(\hat{X})} = m[\mathcal{F}^{-1}(\triangleright \otimes 1)(f(X) \otimes 1)] \quad (49)$$

where in formulae (48), (49) we use (see also (2))

$$h \triangleright X_A = [h, X_A], \quad h = \{p_\mu, M_{\mu\nu}\}, X_A = \{x_\mu, \Lambda_{\mu\nu}\} \quad (50)$$

---

<sup>8</sup>See [1], Sect. 5. The compact formula for the corresponding target map is under consideration. We add that in [1] it was proved explicitly that the Heisenberg doubles of finite-dimensional Hopf algebras carry a Hopf algebroid structure. Our Hopf algebra  $H$  is infinite-dimensional, however filtered by finite-dimensional subalgebras  $H_N$ , and the limit  $N \rightarrow \infty$  should be considered for delivering the rigorous proof.

and for Lorentz sector we obtain

$$[M_{\mu\nu}, \Lambda_{\rho\sigma}] = \eta_{\rho\nu}\Lambda_{\mu\sigma} - \eta_{\rho\mu}\Lambda_{\nu\sigma}. \quad (51)$$

Substituting in (49) the twist (16) we get the following explicit formulas describing base algebra  $\hat{X}_A = \{\hat{x}_\mu, \tilde{\Lambda}_{\mu\nu}\}$

$$\begin{aligned} \hat{x}_\mu &= m[\mathcal{F}^{-1}(\triangleright \otimes 1)(x_\mu \otimes 1)] \\ &= x_\alpha (e^{-u\mathcal{K}})_\mu^\alpha + (1-u) \frac{ia_\mu}{2\kappa} \theta M \end{aligned} \quad (52)$$

$$\begin{aligned} \tilde{\Lambda}_{\rho\sigma} &= m[\mathcal{F}^{-1}(\triangleright \otimes 1)(\Lambda_{\rho\sigma} \otimes 1)] \\ &= (e^{-u\mathcal{K}})_\rho^\alpha \Lambda_{\alpha\sigma} \end{aligned} \quad (53)$$

satisfying the following algebraic relations

$$[\hat{x}_\mu, \hat{x}_\nu] = \frac{i}{\kappa} (a_\nu \theta_\mu^\alpha - a_\mu \theta_\nu^\alpha) \hat{x}_\alpha \quad (54)$$

$$[\hat{x}_\mu, \tilde{\Lambda}_{\rho\sigma}] = -\frac{i}{\kappa} a_\mu \theta_\rho^\alpha \tilde{\Lambda}_{\alpha\sigma} \quad (55)$$

$$[\tilde{\Lambda}_{\mu\nu}, \tilde{\Lambda}_{\rho\sigma}] = 0. \quad (56)$$

The generators  $\Lambda_{\rho\sigma}$  and  $M_{\mu\nu}$  satisfying the relation (51) describe the pair of undeformed dual variables in Lorentz sector, defined by the limit  $\kappa \rightarrow \infty$  of the formula (43).

## 4 Hopf algebroid structures

The Hopf algebroid is described by the data  $\mathcal{B}(H, A; s, t; \tilde{\Delta}, S, \epsilon)$  where  $H$  is total algebra,  $A$  the base algebra, and bialgebroidal coproducts  $\tilde{\Delta}$  are described by the maps  $H \rightarrow H \otimes_A H$  into  $(A, A)$  bimodules  $H \otimes_A H$ , which do not inherit the  $H$  algebra structure. The bimodule  $(A, A)$  property follows from the assumed existence of two maps:

- i) source map  $s(a) : A \rightarrow H$  (homomorphism),
- ii) target maps  $t(a) : A \rightarrow H$  (antihomomorphism),

where

$$[s(a), t(b)] = 0 \quad a, b \in A \quad s(a), t(b) \in H. \quad (57)$$

The relation (57) permits to introduce the basic  $(A, A)$  bimodule formula, namely  $ahb = ht(a)s(b)$ . Further, the algebra  $H \otimes_A H$  can be defined as the quotient of  $H \otimes H$  by left ideal  $\mathcal{I}_L$  generated by the following elements<sup>9</sup>

$$\mathcal{I}_L = s(a) \otimes 1 - 1 \otimes t(a) \quad a \in A. \quad (58)$$

---

<sup>9</sup>The use of left ideal (58)-(59) describes right bialgebroid  $\tilde{\mathcal{H}}^R$ . The elements of left bialgebroid  $\tilde{\mathcal{H}}^L$  are intertwined with  $\tilde{\mathcal{H}}^R$  by antipode map  $S$ .

The canonical choice  $s(a) = a$  of the source map leads to canonical form of (58)

$$\mathcal{I}_L^c = a \otimes 1 - 1 \otimes t^c(a) \quad a \in A \quad (59)$$

with the target map  $t^c(a)$  determining the explicit form of coalgebra gauge freedom (see e.g. [7]). Further the canonical coproducts of the elements of base algebra  $A$  are chosen to be half-primitive (see e.g. [1]-[2])

$$\tilde{\Delta}(\hat{a}) = \hat{a} \otimes 1 \quad \hat{a} \in A \quad (60)$$

i.e. the coproducts (60) are homomorphic in trivial way to the algebra structure of  $A$ .

In this paper we have two choices of algebra  $A$ : given by quantum group algebra  $\widehat{\mathbb{G}}$ , with quantum space-times translations  $\hat{\xi}_\mu$ , and by generalized coordinates sector  $\widehat{\mathbb{X}}$  with quantum-deformed Minkowski space-time coordinates introduced as the module algebra of twisted Poincare algebra. Below we shall consider in some detail the second choice.

If the quantum deformation is generated by twist, the bialgebroid source and target map are described by the formulae analogous to relations (49) (see [2],[9]). For the choice (16) of twist factor one gets:

- source map

$$\begin{aligned} s_{\mathcal{F}}(\hat{x}_\mu) &= m[\mathcal{F}^{-1}(\triangleright \otimes 1)(s_0(x_\mu) \otimes 1)] \\ &= x_\alpha (e^{-u\mathcal{K}})_\mu^\alpha + (1-u) \frac{ia_\mu}{2\kappa} \theta M \equiv \hat{x}_\mu \end{aligned} \quad (61)$$

$$\begin{aligned} s_{\mathcal{F}}(\tilde{\Lambda}_{\rho\sigma}) &= m[\mathcal{F}^{-1}(\triangleright \otimes 1)(s_0(\Lambda_{\rho\sigma}) \otimes 1)] \\ &= (e^{-u\mathcal{K}})_\rho^\alpha \Lambda_{\alpha\sigma} \equiv \tilde{\Lambda}_{\rho\sigma} \end{aligned} \quad (62)$$

- target map

$$\begin{aligned} t(\hat{x}_\mu) &= m[(\mathcal{F}^{-1})^\tau(\triangleright \otimes 1)(t_0(x_\mu) \otimes 1)] \\ &= x_\alpha (e^{(1-u)\mathcal{K}})_\mu^\alpha - u \frac{ia_\mu}{2\kappa} \theta M = \hat{x}_\alpha (e^{\mathcal{K}})_\mu^\alpha - i \frac{a_\mu}{2\kappa} \theta M \end{aligned} \quad (63)$$

$$\begin{aligned} t(\tilde{\Lambda}_{\rho\sigma}) &= m[(\mathcal{F}^{-1})^\tau(\triangleright \otimes 1)(t_0(\Lambda_{\rho\sigma}) \otimes 1)] \\ &= (e^{(1-u)\mathcal{K}})_\rho^\alpha \Lambda_{\alpha\sigma} = (e^{\mathcal{K}})_\rho^\alpha \tilde{\Lambda}_{\alpha\sigma}. \end{aligned} \quad (64)$$

One can check that images of source and target maps commute as follows

$$[s(\hat{X}_A), s(\hat{X}_B)] = \mathcal{C}_{AB}^C s(\hat{X}_C) \quad (65)$$

$$[t(\hat{X}_A), t(\hat{X}_B)] = -\mathcal{C}_{AB}^C t(\hat{X}_C) \quad (66)$$

$$[s(\hat{X}_A), t(\hat{X}_B)] = 0 \quad (67)$$

where constant structures  $\mathcal{C}_{AB}^C$  describe the Lie algebraic structure of commutation relations (54)-(56). The coproducts of base algebra elements are given by

relations (60), and the coalgebra of generalized momentum sector  $H$  is described by Hopf-algebraic formulae (14) and (19)-(20).

In the case of our twisted Hopf algebroid the canonical ideal (59) is the following

$$\mathcal{I}_L^c(\hat{x}_\mu) = \hat{x}_\mu \otimes 1 - 1 \otimes (\hat{x}_\alpha (e^\mathcal{K})_\mu^\alpha - i \frac{a_\mu}{2\kappa} \theta M) \quad (68)$$

$$\mathcal{I}_L^c(\tilde{\Lambda}_{\rho\sigma}) = \tilde{\Lambda}_{\rho\sigma} \otimes 1 - 1 \otimes (e^\mathcal{K})_\rho^\alpha \tilde{\Lambda}_{\alpha\sigma} \quad (69)$$

with counits given by the canonical formula

$$\epsilon(\hat{X}_A) = m[\mathcal{F}^{-1}(\triangleright \otimes 1)(\epsilon_0(X_A) \otimes 1)] = \hat{X}_A \quad (70)$$

and  $\epsilon(h) = 0$  and  $\epsilon(1) = 1$ . The antipodes should satisfy the following relations

$$S(t(\hat{X}_A)) = s(\hat{X}_A) = \hat{X}_A \quad (71)$$

$$m[(1 \otimes S) \circ \tilde{\Delta}] = s\epsilon = \epsilon \quad (72)$$

$$m[(S \otimes 1) \circ \tilde{\Delta}] = t\epsilon S. \quad (73)$$

Using (71) one gets explicit formulae for the antipodes

$$S(\hat{x}_\mu) = (e^\mathcal{K})_\mu^\alpha \hat{x}_\alpha - i \frac{a_\mu}{2\kappa} \theta^{\alpha\beta} M_{\alpha\beta} = t(\hat{x}_\mu) \quad (74)$$

$$S(\tilde{\Lambda}_{\mu\nu}) = (e^\mathcal{K})_\mu^\alpha \tilde{\Lambda}_{\alpha\nu} = t(\tilde{\Lambda}_{\mu\nu}). \quad (75)$$

In our case we have  $S^2 = 1$  and the anchor map is not needed (see [1]-[3]).

## 5 Final remarks

The generalization of Hopf algebras, which we define over commutative ring, are generalized to Hopf-algebraic structures over noncommutative ring, called base algebra  $A$ , leads to the notion of Hopf algebroids. If the algebra sector of Hopf algebroid  $\mathcal{B}$  is endowed with (pre)symplectic structure, as it occurs in Heisenberg double construction presented here, we obtain various models of quantum-deformed phase space algebras. We presented in this paper two types of quantum-deformed phase spaces with Hopf algebroid structure: with coordinates described by quantum-deformed Poincare group algebra and with NC space-time coordinates generated by twist-dependent star product formula (48).

Our considerations have been illustrated by explicit example of Drinfeld twist, but our considerations can be generalized to arbitrary Drinfeld twist case. It has been shown (see [3], Sect. 4) that there exists a generic construction of bialgebroid associated with  $\mathcal{FRT}$  quantization method of quantum matrix groups, which also in general case provides the braid-commutative choice of base algebras defining the Yetter-Drinfeld module [3].

Hopf algebroids are defined in their coalgebraic sectors by the coproducts  $\tilde{\Delta} : H \rightarrow H \otimes H$  defined modulo the coproduct gauge freedom (see e.g. [7];

we stress that  $\otimes$  denotes standard tensor product). It is an interesting problem to characterize the algebraic manifold  $\mathcal{C}$  parametrizing the coproduct gauge transformations, and subsequently define the coproduct gauge-invariant quantum two-particle phase space as the factor algebra  $H \otimes H/\mathcal{C}$  which does not depend on the choice of coproduct gauge.

We add that  $\mathcal{HD}$  construction and corresponding Hopf algebroid structures can be introduced also for infinite-dimensional affine algebras, which are linked with integrable many-body systems of Cologero type (e.g. Cologero-Moser [38]-[39] and Ruijsenaars-Schneider [40]-[41] integrable hierarchies) as well as applied for dynamical quantum groups, described by parameter-dependent (dynamical) Yang-Baxter equations (see e.g. [42]). Other extension, which has as well the link with algebroid structures, is related with recent development in string models with applications of generalized geometries and T-duality covariance. Such generalized phase space dynamics formulated in the framework of string theories, related with double field theories, has been recently investigated under the name of metastring theories (see e.g. [43], [44]).

## Acknowledgements

One of the authors (J.L.) would like to thank Andrzej Borowiec and Zoran Skoda for valuable comments. J.L. and M.W. have been supported by Polish National Science Center, project 2017/27/B/ST2/01902.

## References

- [1] J-H Lu, *Hopf algebroids and quantum grupoids*, *Int. Journ. Math.* **7** (1996) **47**, arXiv:q-alg/9505024
- [2] Ping Xu, *Quantum grupoids*, *Commun. Math. Phys.* **216** (2001) 539, arXiv:math/9905192
- [3] T. Brzezinski, G. Militaru, *Bialgebroids,  $\times_A$ -bialgebras and duality*, *Journ. of Alg.* **251** (2002) 279, arXiv:math/0012164 [math.QA]
- [4] G. Bohm, *Hopf algebroids*, Handbook of Algebra vol. 6, Elsevier/North-Holland, Amsterdam, 2009, arXiv:0805.3806[math.QA]
- [5] T. Jurić, S. Meljanac and R. Štrajn, *Twists, realizations and Hopf algebroid structure of kappa-deformed phase space*, *Int. J. Mod. Phys. A* **29** (2014) 1450022, arXiv:1305.3088
- [6] T. Jurić, D. Kovačević, S. Meljanac,  *$\kappa$ -deformed phase space, Hopf algebroid and twisting*, *SIGMA* **10** (2014), 106, arXiv:1402.0397

- [7] J. Lukierski, Z. Skoda, M. Woronowicz,  $\kappa$ -deformed covariant quantum phase spaces as Hopf algebroids, *Phys. Lett.* **B750** (2015) 401, arXiv.1507.02612
- [8] S. Meljanac, Z. Skoda, M. Stojić, Lie algebra type noncommutative phase spaces as Hopf algebroid, *Lett. Math. Phys.* **107** (2017) 475, arXiv.1409.8188[math.QA]
- [9] A. Borowiec, A. Pachoł, Twisted bialgebroids versus bialgebroids from a Drinfeld twist, *J. Phys.* **A50** (2017) 055205, arXiv.1603.09280
- [10] J. Lukierski, D. Meljanac, S. Meljanac, D. Pikutić, M. Woronowicz, Lie-deformed quantum Minkowski spaces from twists: Hopf-algebraic versus Hopf-algebroid approach, *Phys. Lett.* **B777** (2018) 1, arXiv.1710.09772
- [11] J. M. Souriau, *Structure des systemes dynamiques*, ed. Dunod, Paris, 1970.
- [12] B. Kostant, *Quantization and representation theory*, Springer Lecture Notes in Math. **170** (1970) 87
- [13] H. Bacry, *Space-time and degrees of freedom of the elementary particle*, *Comm. Math. Phys.* **5** (1967) 97
- [14] F. Lurçat, *Strongly Decaying Particles and Relativistic Invariance*, *Phys. Rev.* **173** (1968) 1461
- [15] H. Bacry, A. Kihlberg, *Wavefunctions on Homogeneous Spaces*, *J. Math. Phys.* **10** (1969) 2132
- [16] A.P.Balachandran, G.Marmo, B.S.Skagerstam, A.Stern, *Spinning Particles in General Relativity*, *Phys. Lett.* **B89** (1980) 199
- [17] A.P.Balachandran, G.Marmo, B.S.Skagerstam, A.Stern, in *Gauge Theories and Fiber Bundles*, ed. J. Ehlers et al., Springer Lecture Notes in Physics 188, Chapt. 5 and 8
- [18] D. Finkelstein, *Internal Structure of Spinning Particles*, *Phys. Rev.* **100** (1955) 924
- [19] F. Lurçat, *Quantum field theory on the Poincare group*, in Cargese Lectures in Theor. Physics. Proc. of Summer School in Cargese (Sept. 1965) (1967) 83
- [20] E. P. Wigner, *Theoretical physics*, ed. A. Salam, Trieste, ICTP, (1963) p.59
- [21] L. C. Biedenharn, H. W. Braden, P. Truini, H. van Dam, *Relativistic wavefunctions on spinor spaces*, *J. Phys.* **A21** (1988) 3593
- [22] S.M. Kuzenko, S.L. Lyakhovich, A.Yu. Segal, *A Geometric Model of Arbitrary Spin Massive Particle*, *Int.J.Mod.Phys.* **A10** (1995), hep-th/9403196

- [23] A. A. Kirillov, *Elements of the theory of representations*, ed. Springer (1976)
- [24] V. I. Arnold, *Mathematical methods of classical mechanics*, ed. Springer (1980)
- [25] V. G. Drinfeld, *Proc. of Int. Math. Congress*, Berkeley (1985), p. 798
- [26] J. H. Lu, A. Weinstein, *Poisson-Lie groups, dressing transformations and Bruhat decompositions*, *J. Diff. Geom.* **31** (1990) 501
- [27] A. Yu. Alekseev, A. Z. Malkin, *Symplectic structures associated to lie-Poisson groups*, *Comm. Math. Phys.* **162** (1994) 147; hep-th/9303038
- [28] A. Ballesteros, F. J. Herranz, F. Musso, *On quantum deformations of (anti-)de Sitter algebras in (2+1) dimensions*, *J.Phys.Conf.Ser.* **532** (2014) 012002, arXiv:1302.0684 [hep-th]
- [29] A. Ballesteros, F. J. Herranz, C. Meusburger, P. Naranjo, *Twisted (2+1)  $\kappa$ -AdS Algebra, Drinfel'd Doubles and Non-Commutative Spacetimes*, *SIGMA* **10** (2014) 052, arXiv:1403.4773 [hep-th]
- [30] J. Lukierski, M. Woronowicz, *New Lie-Algebraic and Quadratic Deformations of Minkowski Space from Twisted Poincare Symmetries*, *Phys. Lett.* **B633** (2006) 116, arXiv:hep-th/0508083
- [31] S. Majid, *Foundations of Quantum Groups*, Cambridge University Press (1995)
- [32] C. Gonera, P. Kosinski, P. Maslanka, S. Giller, *Space-time symmetry of noncommutative field theory*, *Phys.Lett.* **B622** (2005) 192, arXiv:hep-th/0504132
- [33] A. Borowiec, A. Pachol, *Heisneberg Doubles of Quantized Poincare Algebras*, *Theor. and Math. Phys.* **169** (2011) 1620
- [34] A. Borowiec, J. Lukierski, V.N. Tolstoy, *Basic Quantizations of D=4 Euclidean, Lorentz, Kleinian and Quaternionic  $o^*(4)$  Symmetries*, *JHEP* **1711** (2017) 187, arXiv:1708.09848
- [35] D. Meljanac, S. Meljanac, D. Pikutić, K. S. Gupta, *Twisted statistics in Lie-deformed Minkowski spaces*, *Phys. Rev.* **D 96** (2017) 105008, arXiv:1703.09511
- [36] A. P. Balachandran, A. Joseph, P. Padmanabhan, *Non-Pauli Transitions From Spacetime Noncommutativity*, *Phys.Rev.Lett.* **105** (2010) 051601, arXiv:1003.2250; A. P. Balachandran, P. Padmanabhan, *Non-Pauli Effects from Noncommutative Spacetimes*, *JHEP* **1012** (2010) 001, arXiv:1006.1185

- [37] L. D. Faddeev, N. Yu. Reshetikhin, L. Takhtajan, *Quantization of Lie groups and Lie algebras*, *Algebra i Analiz*, **1** (1989) 178 (in Russian); English version in *Leningrad Math. Journ.* **1** (1990)
- [38] E. K. Sklyanin, *Dynamical  $r$ -matrices for the Elliptic Calogero-Moser Model*, *Alg.Anal.* **6** (1994) 227-237; *St.Petersburg Math.J.* **6** (1995) 397-406, arXiv:hep-th/9308060
- [39] A. Gorsky, N. Nekrasov, *Relativistic Calogero-Moser model as gauged WZW theory*, *Nucl.Phys.* **B436** (1995) 582, arXiv:hep-th/9401017
- [40] S. N. M. Ruijsenaars, *Complete integrability of relativistic Calogero-Moser systems and elliptic function identities*, *Comm. Math. Phys.* **110** (1987) 191
- [41] G. E. Arutyunov, S. A. Frolov, P. B. Medvedev, *Elliptic Ruijsenaars-Schneider model via the Poisson reduction of the Affine Heisenberg Double*, *J. Phys.* **A30** (1997) 5051, arXiv:hep-th/9607170
- [42] P. Etingof, A. Varchenko, *Geometry and classification of solutions of the Classical Dynamical Yang-Baxter Equation*, *Comm. Math. Phys.* **192** (1998) 77; *Solutions of the quantum dynamical Yang-Baxter equation and dynamical quantum groups*, *ibid.* **196** (1998) 591
- [43] L. Freidel, R. G. Leigh, D. Minic, *Metastring Theory and Modular Spacetime*, *JHEP* **1506** (2015) 006, arXiv:1502.08005 [hep-th]
- [44] L. Freidel, F. J. Rudolph, D. Svoboda, *Generalized Kinematics for Double Field Theory*, *JHEP* 1711, (2017) 175, arXiv:1706.07089 [hep-th]