



Review

Short Overview of Some Assays for the Measurement of Antioxidant Activity of Natural Products and Their Relevance in Dermatology

Morana Jaganjac ¹, Vesna Sredoja Tisma ² and Neven Zarkovic ^{1,*}

¹ Laboratory for Oxidative Stress, Division of Molecular Medicine Rudjer Boskovic Institute, 10000 Zagreb, Croatia; morana.jaganjac@irb.hr

² Polyclinic Department of Dermatology and Venereology, University Hospital Dubrava, 10000 Zagreb, Croatia; vesna.tisma@mail.inet.hr

* Correspondence: zarkovic@irb.hr

Abstract: Impaired systemic redox homeostasis is implicated in the onset and development of various diseases, including skin diseases. Therefore, continuous search for natural products with antioxidant bioactivities applicable in biomedicine is attractive topic of general interest. Research efforts aiming to validate antioxidant potentials of natural products has led to the development of several assays based on various test principles. Hence, understanding the advantages and limitations of various assays is important for selection of assays useful to study antioxidant and related bioactivities of natural products of biomedical interest. This review paper gives a short overview on some chemical and cellular bioassays used to estimate the antioxidant activity of chosen natural products together with a brief overview on the use of natural products with antioxidant activities as adjuvant medicinal remedies in dermatology.

Keywords: oxidative stress; antioxidants; assays; natural products; skin diseases



Citation: Jaganjac, M.; Sredoja Tisma, V.; Zarkovic, N. Short Overview of Some Assays for the Measurement of Antioxidant Activity of Natural Products and Their Relevance in Dermatology. *Molecules* **2021**, *26*, 5301. <https://doi.org/10.3390/molecules26175301>

Academic Editors: José Pinela, Lillian Barros and Maria Ines Dias

Received: 23 July 2021

Accepted: 25 August 2021

Published: 31 August 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Physical, psychological, chemical or environmental stress may provoke biological responses that can induce excessive production of reactive oxygen and nitrogen species (ROS and RNS) [1,2], which are otherwise continuously formed endogenously and contribute to the normal, oxidative energy metabolism of cells. ROS were first perceived as harmful byproducts of aerobic metabolism that may promote the onset and development of different diseases, however, their importance in redox signaling and normal cellular functioning is well recognized today. Under normal conditions generation and removal of ROS are in a fine balance. Antioxidant defense systems balance the level of prooxidants and antioxidants to maintain redox homeostasis. Disrupted redox equilibrium in favor of prooxidants will lead to oxidative/nitrosative stress. Depending on the concentration, reactivity and diffusion distance [3], ROS can react with surrounding molecules via different mechanisms, such as hydrogen abstraction and donation or acceptance of electrons [4]. The interaction of ROS/RNS with macromolecules, like proteins, carbohydrates, lipids and nucleic acids, can result in the loss or gain of function, or in the case of lipids can trigger a chain reaction of lipid peroxidation [5–7]. Peroxidation of lipids is of particular biological relevance as it can alter membrane fluidity, transmembrane transport and interaction of macromolecules (i.e., lipid–lipid and lipid–protein) thus impairing normal cell function or even leading to cell death [8]. Among the final products of lipid peroxidation are reactive aldehydes, such as 4-hydroxynonenal (4-HNE), which can, depending on the concentration, have a role either in the physiology or pathology of the cell [9]. In addition, reactive aldehydes are involved in various cellular processes such as regulation of cell growth, inflammation, signal transduction and apoptosis [10–14]. Eventually, 4-HNE can persist in the form of protein

adducts being able to induce and/or propagate oxidative stress even in the absence of ROS [6,9]. Therefore, it is crucial to maintain cellular antioxidant defenses in order to avoid a rise in 4-HNE concentration from physiological to pathological levels, especially for the cells exposed to potentially harmful effects of chemical (pro-oxidants) or physical inducers (α -rays, UV-irradiation) of ROS/RNS. This is particularly relevant for skin, the largest body organ that serves as biological barrier for physical and chemical environmental factors. These factors may act as oxidants or mediators in the process of generation of ROS and RNS, and together with endogenously formed reactive species make the skin a major target for oxidative stress contributing to the onset and development of various skin pathologies. In addition, skin inflammation is associated with a number of cutaneous diseases. The formation of peroxynitrite (ONOO^-), in the reaction between superoxide and nitric oxide (NO^\bullet), induces the level of nitrated proteins in the skin contributing to inflammation [15]. Furthermore, ONOO^- can induce lipid peroxidation yielding nitrogen-containing oxidized lipid derivatives [16] and may also induce DNA strand breakage further contributing to pathophysiology of inflammation [17].

Natural products with antioxidant bioactivity have long been recognized as a valuable tool in the management of oxidative/nitrosative stress-induced pathologies. This review discusses the natural product discovery workflow to identify products with antioxidant activity, with a focus on available chemical and cellular bioassays used to estimate the antioxidant activity. Finally, at the end of this review, special attention is given to dys-regulated redox homeostasis in cutaneous diseases with a brief overview of the potential use of natural products with antioxidant activity as an adjuvant therapeutic approach in dermatology.

2. Synthetic Antioxidants vs. Natural Antioxidants

Substances that can delay or inhibit oxidation of a substrate, or that can upregulate antioxidant defense systems are defined as antioxidants [18,19]. Antioxidant defense systems found in humans are divided into two types: enzymatic antioxidants and non-enzymatic antioxidants. Enzymatic antioxidants are almost exclusively endogenously formed, while the origin of non-enzymatic antioxidants can be either endogenous or exogenous. The nuclear factor erythroid 2-like 2 (Nrf2), thioredoxin and glutathione (GSH) systems are among the major endogenous antioxidant defenses and their mechanisms of action for cellular ROS detoxification have been recently reviewed [20]. Exogenous antioxidants may be of natural or synthetic origin. Synthetic phenolic antioxidants (SPA) are the most frequently used synthetic antioxidants due to their low cost, higher stability and availability. Butylated hydroxytoluene, butylated hydroxyanisole and tert-butyl hydroquinone are among the most frequently used SPAs. SPAs are increasingly used globally in the food industry and consumer products and are unintended contaminants of the environment [21–24]. The emerging evidence stresses that long-term exposure to SPAs has adverse effects on human health [24], and the use of some common SPAs in the European Union has been restricted by several directives. Hence, there is a need to replace them with antioxidants of natural origin. Plants are good sources of natural antioxidants that can in general be classified as phenolic compounds, vitamins and carotenoids [25]. However, although many naturally derived compounds have antioxidant properties, the majority are not suitable for human use. Among the properties to consider are stability, active concentration and safety, just as with medicines and nutraceuticals. Thus, the evaluation of toxicity in the *in vitro* cellular models should be part of the natural products discovery workflow (Figure 1).




Figure 1. Workflow for the natural products discovery. Extracts from a natural source, such as a plant, may be prepared by either solvent extraction or solvent-free extraction. Solvent extraction is the conventional technique and the polarity index of solvents used will determine the composition of crude extracts [26]. Crude extracts are then examined for potential antioxidant properties by chemical antioxidant and only those identified as bioactive ‘hit’ extracts are further evaluated in the cellular antioxidant assays. Cellular assays also include a toxicity assay to evaluate safety. Extracts acknowledged for their potential antioxidant activity are fractionated based on specific properties in order to reduce the complexity of the extract and preferably isolate pure compounds. High throughput screening (HTS) of each fraction for the antioxidant activity is followed by the purification of compounds from fractions with promising antioxidant activity, structural elucidation and compound identification.

Indeed, before antioxidants can be used as food additives and/or adjuvant medicinal remedies for the integrative biomedicine purposes or as ordinary pharmacological therapeutics, their safety should be assessed by different toxicity tests as requested by the regulatory bodies, such as the European Food Safety Authority and US Food and Drug Administration [25]. Moreover, reactivity and stoichiometric factor, liposolubility and secondary reactions of new natural products with antioxidant activity should be evaluated as well [27].

3. Assays for the Measurement of Antioxidant Activity

Traditionally antioxidant assays are divided into hydrogen atom transfer (HAT), or single electron transfer (SET) based methods (Figure 2) [28]. Although the final products of both mechanisms might be identical, their kinetics greatly differ.




Figure 2. Quenching mechanisms of target radicals by hydrogen atom transfer and/or single electron transfer. AH—antioxidant; A[•]—antioxidant radical; Phe-O[•]—aryloxy radical; Phe-OH—phenolic compound; ROO[•]—target radical.

HAT assays are based on the transfer of hydrogen atoms from antioxidant or phenolic compounds to target radicals, and the kinetics of reaction depends on the solvent used. The hydrogen donation is enhanced in aqueous solutions compared to alcohol ones [29]. However, HAT assays are not dependent on pH, while pH is important for SET assays as an increase in the pH will accelerate electron transfer [30]. Some of the popular HAT-based antioxidant assays are oxygen radical absorbance capacity (ORAC), total radical-trapping antioxidant parameter (TRAP) and β -carotene bleaching assay, while commonly used SET assays include total phenolic assay, 2,3-diphenyl-1-picrylhydrazyl (DPPH) free radical method, trolox equivalent antioxidant capacity (TEAC) and ferric reducing-antioxidant power (FRAP) assay.

3.1. Chemical-Based Antioxidant Assays

Low cost and high-throughput screening (HTS) assays are chemical-based methods preferred for the initial screening process of the natural products discovery workflow. Antioxidants may have different mechanisms of action, according to which antioxidant assays are divided into either (1) scavenging activity assays, (2) reducing antioxidant power assays, (3) lipid peroxidation inhibitory potential or (4) Metal ion chelation (Figure 3).




Figure 3. Antioxidant chemical-based assays divided based on the antioxidant activity to reducing power, lipid peroxidation inhibition, metal ion chelation and scavenging activity assay. Abbreviations: ABTS—2,2-azinobis-(3-ethylbenzothiazoline-6-sulphonic acid); CERAC—cerium reducing antioxidant capacity; CUPRAC—cupric ion reducing antioxidant capacity; DMPD—*N,N*-dimethyl-*p*-phenylenediamine; DPPH—2,3-diphenyl-1-picrylhydrazyl; FOX—ferrous oxidation-xylenol orange; FTC—ferric thiocyanate; FRAP—ferric reducing-antioxidant power; LDL—low density lipoprotein; LIPIC—lipid peroxidation inhibition capacity; ORAC—oxygen radical absorbance capacity; TEAC—trolox equivalent antioxidant capacity; TRAP—total radical-trapping antioxidant parameter; TBARS—thiobarbituric acid reactive substances.

- Scavenging activity assays are based on the ability of an antioxidant to scavenge stable free radicals. Some of the most common antioxidant assays based on scavenging activity include ORAC, DPPH, 2,2'-azinobis-(3-ethylbenzothiazole-6-sulphonate) (ABTS)/TEAC assay and *N,N*-dimethyl-*p*-phenylenediamine radical scavenging (DMPD) assay [31,32]. Although the basic principle is the same, their antioxidant solubility differs from those soluble in aqueous and alcoholic media (e.g., ABTS^{•+} and

DMPD^{•+} radical) to those soluble in organic solvents (e.g., DPPH[•] radical). Other scavenging assays include those specific to certain ROS/RNS, such as ONOO⁻ [33], hydroxyl radical [34], superoxide anion (O₂^{•-}) [35] or NO[•] scavenging assays [36].

- Reducing antioxidant power assays are based on the principle that antioxidants acts as reductants by accepting electrons, for example, from transition metals, notably iron and copper. The most frequently used reducing antioxidant power assays are FRAP, cupric ions reducing power assay (CUPRAC) and total phenolic content assay (Folin–Ciocalteu method) [32].
- The antioxidant assays widely used to evaluate antioxidant ability to inhibit lipid peroxidation include β-carotene bleaching assay, thiobarbituric acid reactive species assay (TBARS), lipid peroxidation inhibition capacity assay (LIPIC) and conjugated dienes [37,38].
- Metal ion chelation is an important property of an antioxidant as transition metals, via the Fenton reaction, promote oxidative stress and lipid peroxidation [2,39,40]. Ferrous ion is commonly used to assess the metal chelation capacity of antioxidants [41]. Curcumin, resveratrol, L-carnitine and ferrozine are among many well documented antioxidants with metal chelation ability [32].

Several comprehensive reviews on the strengths and limitations of the above chemical antioxidant assays including the detailed mechanisms behind each assay have been recently published [32,37,41–43].

3.2. Cellular-Based Antioxidant Assays

Natural products identified with potential antioxidant activity need to be further evaluated in the cellular model. Antioxidant activity of a large number of natural products will not extrapolate its performance in the biological system, either in vitro as cellular assays or in vivo as animal model studies. Thus, it is necessary to examine the bioavailability, metabolism and mechanism of action in a living system to prove potential antioxidant activities of new natural products [27]. Although in vivo studies would best reflect the effectiveness of the natural products with antioxidant activity, they are not preferred due to low throughput, occasionally bioethical uncertainties, very high costs and are time-consuming. On the other hand, in vitro cellular models, have high throughput, lower cost and are much faster. Nowadays, a plethora of redox-sensitive probes is available that enable live-cell monitoring of oxidative/nitrosative stress (Figure 4). According to the mechanism of antioxidant activity, cellular antioxidant assays could be divided into those screening for: 1. Direct reaction of antioxidants with ROS/RNS, 2. Organelle/membrane-specific antioxidant activity, 3. Inhibition of oxidant enzymes, and 4. Activators of transcription factors promoting antioxidant defense.

A variety of technologies have been developed to detect ROS/RNS, the selection of which to use will depend on several criteria, such as availability, applicability, specificity, selectivity, throughput and cost. Probes for detection of ROS/RNS in living cells are utilized to evaluate the direct effect of antioxidants on the level of reactive species. Decrease in the level of ROS/RNS can be either due to antioxidants scavenging of reactive species or due to inhibition of their generation. The majority of available probes lack specificity but can provide a good indication of potential antioxidant activity. Such probes are 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA), dihydrorhodamine 123 (DHR) and chemiluminescent probes. Although initially H₂DCFDA and DHR were perceived to detect hydrogen peroxide and ONOO⁻, respectively, it was later found that other ROS/RNS, as well as some other cellular molecules, can promote oxidation of the probes giving false-positive results [44]. Chemiluminescent assays, such as luminol or lucigenin-enhanced chemiluminescence have higher sensitivity but are susceptible to redox cycling inducing ROS formation and can, in long-term measurements, give a false increase in the signal. A number of sensitive and selective fluorescent probes for live cells/tissue detection and imaging of ONOO⁻, such as boronate-based polymeric fluorescent [45], *N*-phenylrhodol-based HKGreen-4 [46] and rhodamine based HKYellow probes [47] have been developed

and could be utilized to monitor the effect of antioxidants on ONOO^- . Dihydroethidium is another readily used probe that can react with $\text{O}_2^{\bullet-}$ yielding 2-hydroxyethidium that emits red fluorescence (2-OH-E^+). Unfortunately, DHE also nonspecifically reacts with other ROS forming ethidium, a fluorescent product whose spectra overlaps with the spectra of 2-OH-E^+ [48]. Thus, when selecting those probes, it is crucial to be aware of all factors affecting the signal and to take precautions to reduce false-positive results.




Figure 4. Probes used in HTS assays for the detection of ROS/RNS and lipid peroxidation. Abbreviations: 2-OH-E^+ —2-hydroxyethidium; AO—antioxidants; DCF—dichlorofluorescein; DHE—dihydroethidium; DHR—dihydrorhodamine 123; Et—ethidium; F—fluorescent; H_2DCFDA —2',7'-dichlorodihydrofluorescein diacetate; R^+ —rhodamine 123.

Spin traps and spin probes used for electron-spin resonance assays have high selectivity and specificity and are among the essential tools in oxidative stress research. Similarly, redox proteomics is rapidly emerging in oxidative stress research as it can identify ROS targets and modifications induced [49–51]. However, low throughput and expensive instruments requiring a high level of expertise reduce their applicability. Hence, these analytical approaches could hardly be the preferred choice for HTS of new natural products.

Organelle/membrane targeted redox sensitive probes allow selection of those antioxidants with organelle/membrane specific effects. The addition of a triphenylphosphonium group to DHE led to the novel probe mitoSOX, that has specificity and selectivity for mitochondria derived $\text{O}_2^{\bullet-}$ [52,53]. Afterwards, numerous organelle-targeting redox sensitive fluorescent probes were developed and validated including the redox-sensitive probes based on the chimeric fluorescent protein [54].

Screening for the antioxidant ability to protect cells from lipid peroxidation should preferably be monitored by the radiometric probe, such is $\text{C11-BODIPY}^{581/591}$. The $\text{C11-BODIPY}^{581/591}$ is a fatty acid analog that emits bright red fluorescence in the intact form. The probe is sensitive to ROS and when oxidized the fluorescence shifts to green [55,56].

Moreover, natural products might not act as scavengers of reactive species, but instead might inhibit activity of pro-oxidant enzymes. Among the pro-oxidant enzymes are NADPH oxidases (NOXs) and their excessive ROS production is implicated in various pathologies. A number of compounds, such as polyphenols and alkaloids from natural sources, were found to decrease or to inhibit NOX activity [57].

In addition, antioxidants could exhibit indirect activity via transcriptional regulation of Nrf2. Nrf2 is the primary redox sensor and the key regulator of endogenous antioxidant

defenses, such as members of the glutathione and thioredoxin systems and superoxide dismutase (SOD) [20].

Based on all the above, two or more cellular-based antioxidant assays should be performed and interpreted in the context of data obtained from both assays to confirm the antioxidant activity of natural products.

4. Natural Products with Antioxidant Activity as Adjuvant Therapeutic Approach in Dermatology

Skin diseases are multifactorial, and oxidative stress plays an important role in the pathophysiology of many autoimmune and inflammatory diseases, as well as in other stress- and age-dependent diseases. Accumulating evidence shows that impaired antioxidant defenses [39,40,58–60] and increased levels of oxidants [39,59–64] are important mediators in the pathology of a plethora of cutaneous diseases (Table 1).

Table 1. Imbalance in the redox system of cutaneous diseases.

Skin Disease	Imbalance in the Redox System	Reference
Psoriasis	Myeloperoxidase and GSH/GSSG ratio are increased SOD level is decreased	[65]
Alopecia areata	SOD, paraoxonase and glutathione peroxidase are decreased Total antioxidant capacity is decreased	[58]
Vitiligo	Advanced oxidation protein products, advanced glycation products and malondialdehyde levels are increased Catalase is decreased	[66–69]
Rosacea	Serum peroxide and cutaneous ferritin are increased. Total antioxidative potential is decreased	[39]
Acne vulgaris	Serum levels of malondialdehyde and xanthine oxidase activity are increased. SOD and catalase activity are decreased.	[61]
Oral lichen planus	Salivary uric acid is decreased. Serum gamma glutamyl transferase (GGT) and saliva total antioxidant capacity are increased.	[70]
Localized scleroderma (morphea)/systemic sclerosis	Total oxidant capacity, arylesterase and oxidative stress index are elevated. Nitric oxide, malondialdehyde, asymmetric dimethylarginine, and ROOH in the blood are elevated. Levels of SOD and vitamin C are decreased.	[64]
Chronic venous insufficiency	Malondialdehyde, serum iron and total antioxidant capacity are elevated. Uric acid level in the circulation is low.	[71]
Pemphigus vulgaris	Serum bilirubin, uric acid and albumin are decreased. Serum total oxidant capacity, lipid hydroperoxides and oxidative stress index are increased.	[63,72]
Eczema/dermatitis	SOD, catalase, GPX, GSH, and vitamins A, C, and E, total antioxidant status are decreased in the blood. Total oxidative status, total peroxides and oxidative stress index are increased.	[59,60,73]

The SOD enzyme that accelerates the conversion of the $O_2^{\bullet-}$ to hydrogen peroxide (H_2O_2), is decreased in psoriasis [65], alopecia areata [58], acne vulgaris [61], systemic sclerosis [62] and seborrheic dermatitis [59]. Alterations in the glutathione system, one of the major endogenous antioxidant defenses, were reported for alopecia areata [58], psoriasis [65], oral lichen planus [70] and atopic dermatitis [73]. In addition, vitiligo is also accompanied by decreased antioxidant defenses [67] and elevated oxidation products [68,69] that mediate the death of melanocytes [74]. Deregulated redox homeostasis and oxidation of macromolecules further accompany cutaneous disease pathology [40,61,68,71]. Nowadays, antioxidants from plants with potential antioxidant activity are emerging as potential adjuvant therapies for cutaneous diseases in order to prevent the development of various symptoms. In the Table 2 we provide a list of some natural products with antioxidant activity and with traditional, ethnomedical applications as adjuvant treatments for cutaneous diseases, or those experimentally shown in a disease model to have beneficial effects.

Table 2. Adjuvant therapeutic approach for the treatment of cutaneous diseases.

Skin Disease	Herbal Therapeutic Options	Mechanism of Action	References
Psoriasis	<i>Aloe greatheadii</i> var. <i>davyana</i>	Aqueous ethanol (95%) leaf gel extract has high polyphenol content and high antioxidant capacity.	[75]
	<i>Artemisia anomala</i> S.	Extract inactivates MAPK and caspase pathway, promotes viability of human keratinocytes and increases antioxidant capacity.	[76]
	<i>Astragalus sinicus</i> L.	Aqueous and methanol extracts possess anti-inflammatory activity, and antioxidant activity by regulating cellular redox homeostasis and NF- κ B, JAK/STAT and PI3/Akt signaling pathways.	[77]
	Berry extracts	Wild blueberry, bilberry, cranberry, elderberry, raspberry seed, and strawberry possess antioxidant activity and inhibit VEGF expression and impair angiogenesis.	[78]
	Canadian wood species	Yellow birch extract and black spruce extract had highest antioxidant capacity compared to other species. Black spruce extract demonstrated low toxicity and inhibited proliferation of normal human keratinocytes and non-lesional psoriatic keratinocytes but was not selective.	[79]
	<i>Centella asiatica</i> (L.)	Polar extract modulates cyclooxygenase and lipoxygenase activities suggesting its use for the treatment of psoriasis.	[80]
	<i>Citrus sudachi</i>	Peel extract demonstrated good radical scavenging activity and high ability in reducing power. It also inhibits EGFR-ERK signaling pathway, suppressing proliferation and inducing cell differentiation.	[81]
	<i>Copaifera langsdorffii</i> Desf.	Oleoresin reduces the release of pro-inflammatory cytokines by stimulated monocytes and its treatment improved typical clinical signs.	[82]
	<i>Datura metel</i> L.	Its application significantly reduced typical clinical signs of psoriasis. It also inhibited the inflammatory response which was suggested to be due to the TLR7/8-MyD88-NF- κ B-NLRP3 inflammasome pathway inhibition.	[83]
	French maritime pine bark	High antioxidant and anti-inflammatory properties by inhibiting expression of inducible intercellular adhesion molecule-1 and interferon-gamma mediated activation of Stat1.	[84]

Table 2. Cont.

Skin Disease	Herbal Therapeutic Options	Mechanism of Action	References
	Indian medicinal plants	Extracts from <i>Phyllanthus simplex</i> Retz., <i>Crotolaria juncea</i> Linn., <i>Leucas aspera</i> Linn., and <i>Vitex glabrata</i> R.Br. plants inhibit NO production and lipid peroxidation in keratinocytes and have promising antiproliferative activity.	[85]
	<i>Melissa officinalis</i> ssp. <i>Altissima</i>	The decoction showed high free radical scavenging activity and contributed to psoriasis treatment by decreasing inflammation and enhancing barrier function.	[86]
	<i>Oryza sativa</i> L.	Crude extract has an antioxidative property by enhancing Nrf2, induces expression of anti-inflammatory cytokines while reduces proinflammatory cytokines, impairs expression of psoriasis-associated genes and improves typical clinical signs of disease.	[87]
	<i>Plectranthus madagascariensis</i>	Contains abietane diterpenoids with excellent antioxidant activity.	[88]
	<i>Solanum xanthocarpum</i> Schrad. & Wendl.	Ethanollic stem extract has antioxidant properties and was found to inhibit the expression of proinflammatory cytokines and improves typical clinical signs of psoriasis.	[89]
Alopecia areata	Ginger (<i>Zingiber officinale</i> (L.) Rosc)	Orally administered ginger powder elevated GSH level and reduced malondialdehyde level of erythrocytes and lymphocytes, and improved total antioxidant status in alopecia areata patients	[90]
	Herbal extract	Extract prepared from <i>Urtica dioica</i> root, <i>Urtica urens</i> Leaf, <i>Equisetum arvense</i> leaf, <i>Achillea millefolium</i> aerial part, <i>Matricaria chamomilla</i> flower and <i>Ceratonia siliqua</i> fruit with known antioxidant and anti-inflammatory properties, downregulates expression of IL-1alpha a mediator for the hair loss.	[91]
Vitiligo	<i>Clusia minor</i> L.	Extract of this plant used to treat vitiligo exhibit antioxidant activity as determined by radical scavenging activity and ferro-reducing activity.	[92]
	Date seed	Date seed oil has radical scavenging activity, inhibits lipid peroxidation and protects against H ₂ O ₂ -induced cell death of melanocytes.	[93]
	Ginger	An active compound 6-shogaol has protective effects against H ₂ O ₂ -induced cell stress and activates Nrf2 pathway in epidermal melanocytes.	[94]
	<i>Ginko biloba</i>	Terpenoid bilobalide protects melanocytes from H ₂ O ₂ -induced apoptosis, promotes catalase and glutathione peroxidase 1. Bilobalide also exhibited immunoprotective effect by reducing the release of Hsp70.	[95]
	Green tea	Protects melanocytes from H ₂ O ₂ -induced cell death. Among the major constituents of green tea is Epigallocatechin-3-gallate with high antioxidant and anti-inflammatory potential that was also found to inhibit Janus kinase 2 thus suppressing trafficking of T lymphocytes to melanocytes.	[96,97]

Table 2. Cont.

Skin Disease	Herbal Therapeutic Options	Mechanism of Action	References
	<i>Pyrostegia venusta</i>	Topical and oral administration of leaves extract has antioxidative and anti-inflammatory properties and increases epidermal melanin level in and animal vitiligo model.	[98]
	<i>Scutellaria baicalensis</i>	Baicalein extracted from the plant protects melanocytes from H ₂ O ₂ -induced apoptosis and promotes activation of Nrf2 pathway.	[99]
	<i>Vernonia anthelmintica</i> (L.) Willd.	The extract contains compounds with antioxidant properties and promotes melanogenesis.	[100]
Rosacea	Turmeric (<i>Curcuma longa</i>)	Turmeric has antioxidant and anti-inflammatory properties and the administration of turmeric polyherbal formulation reduces facial redness intensity and distribution.	[101,102]
	<i>Artemisia vulgaris</i>	Essential oil has antioxidant properties with strong metal chelation activity and inhibits growth of <i>Streptococcus pyogenes</i> and <i>Propionibacterium acnes</i> .	[103]
	<i>Cephalaria uralensis</i>	Ethanol extract of aerial parts demonstrated radical scavenging activity, inhibits cyclooxygenase-1 and -2, and inhibits growth of <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , and <i>Propionibacterium acnes</i> .	[104]
	<i>Clausena anisata</i>	Extract inhibits growth of <i>Propionibacterium acnes</i> , has potent antioxidant activity, inhibits lipase and hyaluronidase activity and decreases IL-8 production.	[105]
	<i>Helichrysum kraussii</i>	Extract inhibits growth of <i>Propionibacterium acnes</i> , has potent antioxidant activity	[105]
	<i>Humulus lupulus</i> L.	Hop extracts demonstrated antibacterial activity against five acne causing bacteria, anticollagenase inhibitory activity and good antioxidant capacity.	[106]
	Keishibukuryogan-ka-yokuinin (KBGY)	Oral administration inhibits formation of lipid hydroperoxides and scavenges ROS in plasma.	[107]
Acne vulgaris	<i>Mangifera indica</i> L.	Kernel extract inhibits growth of <i>Propionibacterium acnes</i> , has strong radical scavenging properties, inhibits linoleic acid peroxidation and secretion of IL-8.	[108]
	<i>Neolitsea aciculata</i>	Essential oil inhibits growth of <i>Propionibacterium acnes</i> and <i>Staphylococcus epidermidis</i> , has antioxidant properties and reduces release of inflammatory cytokines.	[109]
	<i>Origanum vulgare</i>	Ethanol extract reduces generation of inflammatory cytokines and suppresses <i>Propionibacterium acnes</i> induced skin inflammation	[110]
	<i>Sargassum polycystum</i> C. Agardh	Methanolic fractions inhibit growth of <i>Propionibacterium acnes</i> , inhibits lipase and have high radical scavenging activities.	[111]
	<i>Selaginella involvens</i>	The extract inhibits production of NO, has NO scavenging effect and inhibits growth of <i>Propionibacterium acnes</i> .	[112]
	<i>Syzygium jambos</i> L.	The ethanol extract inhibits the growth of <i>Propionibacterium acnes</i> , exhibits strong antioxidant activity and inhibits the release of inflammatory cytokines.	[113]

Table 2. Cont.

Skin Disease	Herbal Therapeutic Options	Mechanism of Action	References
Oral lichen planus	Neem tree	Mouthwash with aqueous neem leaves extract improved symptoms of disease in patients.	[114]
	Purslane	Oral administration of antioxidant-rich purslane led to partial or complete clinical improvement in majority of patients.	[115]
Chronic venous insufficiency	Red-vine-leaf	Extract AS195 induces activation of endothelial and red blood cell nitric oxide synthase increasing NO bioavailability and ameliorates tert-butylhydroperoxide induced ROS.	[116]
	<i>Ruscus aculeatus</i>	<i>Ruscus</i> extract showed significant venular constriction, antioxidative and anti-inflammatory properties.	[117]
Eczema/dermatitis	<i>Erythrina stricta</i> Roxb.	<i>Erythrina</i> extracts are active against <i>Staphylococcus aureus</i> and <i>Candida albicans</i> , and extracted erynone demonstrated significant radical scavenging activity.	[118]
	<i>Sapium sebiferum</i> (L.) Roxb.	Phenolic extracts from leaves increase activities of catalase and SOD, increase GSH level and exhibit anti-inflammatory properties in an animal dermatitis model.	[119]
	<i>Sophora alopecuroides</i> L.	The root extract has strongest antioxidant activity and showed inhibitory activity for different enzymes.	[120]

It is obvious that numerous plant extracts possess the capacity for medicinal applications aiming at attenuating symptoms of skin diseases, preventing their occurrence and even to be used for adjuvant therapeutic remedies or nutraceuticals, However, must follow strict regulatory rules and should be proven for their efficiency as antioxidants before being used as such.

5. Conclusions and Future Perspectives

While stress- and age-associated disorders are usually considered to affect internal organs, they very often manifest as skin diseases, mostly associated with skin exposure to UV light, environmental pollutants, persistent oxidative stress and chronic inflammatory processes. However, even local manifestations of the many diseases are often associated with systemic disorders of oxidative homeostasis. Therefore, if intended to be used as adjuvant medicinal remedies or even cosmetics, natural products with antioxidant potential are obliged to meet the high safety and efficacy standards, as do other medicinal remedies. They should be tested for their bioactivities using complementary assays for antioxidants. Whenever possible, antioxidant effects should be complemented by analysis of the potential pro-oxidant effects of the chosen substance(s) in the presence of biofluids (preferably serum), thus resembling complex in vivo processes, as do enzymatic assays for antioxidant capacity and peroxides [121]. Complementary to that, options for in vitro bioassays should be developed to reveal at the same time pro- and anti-oxidant activities thus resembling living systems as in the case of the 4-HNE-Cell ELISA assay, which was recently used to prove cell-type specific bioactivities of the *Aloe vera* extracts [122].

Author Contributions: Conceptualization, M.J. and N.Z.; investigation, M.J. and V.S.T.; writing—original draft preparation, M.J., V.S.T. and N.Z.; writing—review and editing, M.J. and N.Z.; visualization, M.J. and V.S.T. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Spoljaric, D.; Cipak, A.; Horvatic, J.; Andrisic, L.; Waeg, G.; Zarkovic, N.; Jaganjac, M. Endogenous 4-hydroxy-2-nonenal in microalga *Chlorella kessleri* acts as a bioactive indicator of pollution with common herbicides and growth regulating factor of hormesis. *Aquat. Toxicol.* **2011**, *105*, 552–558. [[CrossRef](#)]
2. Poljak-Blazi, M.; Jaganjac, M.; Sabol, I.; Mihaljevic, B.; Matovina, M.; Grce, M. Effect of ferric ions on reactive oxygen species formation, cervical cancer cell lines growth and E6/E7 oncogene expression. *Toxicol. Vitro.* **2011**, *25*, 160–166. [[CrossRef](#)]
3. Jaganjac, M.; Cipak, A.; Schaur, R.J.; Zarkovic, N. Pathophysiology of neutrophil-mediated extracellular redox reactions. *Front. Biosci. Landmark* **2016**, *21*, 839–855. [[CrossRef](#)]
4. Slater, T.F. Free-radical mechanisms in tissue injury. *Biochem. J.* **1984**, *222*, 1–15. [[CrossRef](#)]
5. Jaganjac, M.; Čačev, T.; Čipak, A.; Kapitanović, S.; Trošelj, K.G.; Žarković, N. Even stressed cells are individuals: Second messengers of free radicals in pathophysiology of cancer. *Croat. Med. J.* **2012**, *53*, 304–309. [[CrossRef](#)]
6. Zarkovic, N.; Cipak, A.; Jaganjac, M.; Borovic, S.; Zarkovic, K. Pathophysiological relevance of aldehydic protein modifications. *J. Proteom.* **2013**, *92*, 239–247. [[CrossRef](#)]
7. Jaganjac, M.; Cindrić, M.; Jakovčević, A.; Žarković, K.; Žarković, N. Lipid peroxidation in brain tumors. *Neurochem. Int.* **2021**, *149*, 105118. [[CrossRef](#)]
8. Catalá, A.; Díaz, M. Editorial: Impact of lipid peroxidation on the physiology and pathophysiology of cell membranes. *Front. Physiol.* **2016**, *7*, 423. [[CrossRef](#)]
9. Jaganjac, M.; Milkovic, L.; Gegotek, A.; Cindric, M.; Zarkovic, K.; Skrzydlewska, E.; Zarkovic, N. The relevance of pathophysiological alterations in redox signaling of 4-hydroxynonenal for pharmacological therapies of major stress-associated diseases. *Free Radic. Biol. Med.* **2020**, *157*, 128–153. [[CrossRef](#)]
10. Awasthi, Y.C.; Sharma, R.; Cheng, J.Z.; Yang, Y.; Sharma, A.; Singhal, S.S.; Awasthi, S. Role of 4-hydroxynonenal in stress-mediated apoptosis signaling. *Mol. Asp. Med.* **2003**, *24*, 219–230. [[CrossRef](#)]
11. Zarkovic, N.; Ilic, Z.; Jurin, M.; Schaur, R.J.; Puhl, H.; Esterbauer, H. Stimulation of HeLa cell growth by physiological concentrations of 4-hydroxynonenal. *Cell Biochem. Funct.* **1993**, *11*, 279–286. [[CrossRef](#)]
12. Živković, M.; Žarković, K.; Škrinjar, L.; Waeg, G.; Poljak-Blaži, M.; Šunjić, S.B.; Schaur, R.J.; Žarković, N. A new method for detection of HNE-histidine conjugates in rat inflammatory cells. *Croat. Chem. Acta* **2005**, *78*, 91–98.
13. Jaganjac, M.; Matijević, T.; Cindric, M.; Cipak, A.; Mrakovcic, L.; Gubisch, W.; Zarkovic, N. Induction of CMV-1 promoter by 4-hydroxy-2-nonenal in human embryonic kidney cells. *Acta Biochim. Pol.* **2010**, *57*, 179–183. [[CrossRef](#)]
14. Elrayess, M.A.; Almuraikhy, S.; Kafienah, W.; Al-Menhali, A.; Al-Khelaifi, F.; Bashah, M.; Zarkovic, K.; Zarkovic, N.; Waeg, G.; Alsayrafi, M.; et al. 4-hydroxynonenal causes impairment of human subcutaneous adipogenesis and induction of adipocyte insulin resistance. *Free Radic. Biol. Med.* **2017**, *104*, 129–137. [[CrossRef](#)]
15. Greenacre, S.A.B.; Evans, P.; Halliwell, B.; Brain, S.D. Formation and loss of nitrated proteins in peroxynitrite-treated rat skin in vivo. *Biochem. Biophys. Res. Commun.* **1999**, *262*, 781–786. [[CrossRef](#)]
16. Rubbo, H.; Radi, R.; Trujillo, M.; Telleri, R.; Kalyanaraman, B.; Barnes, S.; Kirk, M.; Freeman, B.A. Nitric oxide regulation of superoxide and peroxynitrite-dependent lipid peroxidation. Formation of novel nitrogen-containing oxidized lipid derivatives. *J. Biol. Chem.* **1994**, *269*, 26066–26075. [[CrossRef](#)]
17. Szabó, C. DNA strand breakage and activation of poly-ADP ribosyltransferase: A cytotoxic pathway triggered by peroxynitrite. *Free Radic. Biol. Med.* **1996**, *21*, 855–869. [[CrossRef](#)]
18. Møller, P.; Loft, S. Dietary antioxidants and beneficial effect on oxidatively damaged DNA. *Free Radic. Biol. Med.* **2006**, *41*, 388–415. [[CrossRef](#)]
19. Harwell, B. Biochemistry of oxidative stress. *Biochem. Soc. Trans.* **2007**, *35*, 1147–1150. [[CrossRef](#)]
20. Jaganjac, M.; Milkovic, L.; Sunjic, S.B.; Zarkovic, N. The NRF2, Thioredoxin, and Glutathione System in Tumorigenesis and Anticancer Therapies. *Antioxidants* **2020**, *9*, 1151. [[CrossRef](#)]
21. Wu, Y.; Venier, M.; Hites, R.A. Broad Exposure of the North American Environment to Phenolic and Amino Antioxidants and to Ultraviolet Filters. *Environ. Sci. Technol.* **2020**, *54*, 9345–9355. [[CrossRef](#)]
22. Wu, Y.; Venier, M.; Hites, R.A. Identification of Unusual Antioxidants in the Natural and Built Environments. *Environ. Sci. Technol. Lett.* **2019**, *6*, 443–447. [[CrossRef](#)]
23. Wang, W.; Xiong, P.; Zhang, H.; Zhu, Q.; Liao, C.; Jiang, G. Analysis, occurrence, toxicity and environmental health risks of synthetic phenolic antioxidants: A review. *Environ. Res.* **2021**, *201*, 111531. [[CrossRef](#)]
24. Liu, R.; Mabury, S.A. Synthetic Phenolic Antioxidants: A Review of Environmental Occurrence, Fate, Human Exposure, and Toxicity. *Environ. Sci. Technol.* **2020**, *54*, 11706–11719. [[CrossRef](#)]
25. Lourenço, S.C.; Moldão-Martins, M.; Alves, V.D. Antioxidants of natural plant origins: From sources to food industry applications. *Molecules* **2019**, *24*, 4132. [[CrossRef](#)] [[PubMed](#)]
26. Wakeel, A.; Jan, S.A.; Ullah, I.; Shinwari, Z.K.; Xu, M. Solvent polarity mediates phytochemical yield and antioxidant capacity of *Isatis tinctoria*. *PeerJ* **2019**, *2019*, e7857. [[CrossRef](#)]
27. López-Alarcón, C.; Denicola, A. Evaluating the antioxidant capacity of natural products: A review on chemical and cellular-based assays. *Anal. Chim. Acta* **2013**, *763*, 1–10. [[CrossRef](#)]
28. Apak, R.; Gorinstein, S.; Böhm, V.; Schaich, K.M.; Özyürek, M.; Güçlü, K. Methods of measurement and evaluation of natural antioxidant capacity/activity (IUPAC technical report). *Pure Appl. Chem.* **2013**, *85*, 957–998. [[CrossRef](#)]

29. Litwinienko, G.; Ingold, K.U. Solvent effects on the rates and mechanisms of reaction of phenols with free radicals. *Acc. Chem. Res.* **2007**, *40*, 222–230. [[CrossRef](#)]
30. Prior, R.L.; Wu, X.; Schaich, K. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J. Agric. Food Chem.* **2005**, *53*, 4290–4302. [[CrossRef](#)]
31. Shahidi, F.; Zhong, Y. Measurement of antioxidant activity. *J. Funct. Foods* **2015**, *18*, 757–781. [[CrossRef](#)]
32. Gulcin, İ. Antioxidants and antioxidant methods: An updated overview. *Arch. Toxicol.* **2020**, *94*, 651–715. [[CrossRef](#)]
33. Krzyżanowska-Kowalczyk, J.; Kowalczyk, M.; Ponczek, M.B.; Pecio, Ł.; Nowak, P.; Kolodziejczyk-Czepas, J. Pulmonaria obscura and pulmonaria officinalis extracts as mitigators of peroxynitrite-induced oxidative stress and cyclooxygenase-2 inhibitors—in vitro and in silico studies. *Molecules* **2021**, *26*, 631. [[CrossRef](#)]
34. Kunchandy, E.; Rao, M.N.A. Oxygen radical scavenging activity of curcumin. *Int. J. Pharm.* **1990**, *58*, 237–240. [[CrossRef](#)]
35. Fazilatun, N.; Nornisah, M.; Zhari, I. Superoxide radical scavenging properties of extracts and flavonoids isolated from the leaves of *Blumea balsamifera*. *Pharm. Biol.* **2004**, *42*, 404–408. [[CrossRef](#)]
36. Asokkumar, K.; Umamaheswari, M.; Sivashanmugam, A.T.; Subhadradevi, V.; Subhashini, N.; Ravi, T.K. Free radical scavenging and antioxidant activities of *Glinus oppositifolius* (carpet weed) using different in vitro assay systems. *Pharm. Biol.* **2009**, *47*, 474–482. [[CrossRef](#)]
37. Zhang, J.; Stanley, R.A.; Melton, L.D. Lipid peroxidation inhibition capacity assay for antioxidants based on liposomal membranes. *Mol. Nutr. Food Res.* **2006**, *50*, 714–724. [[CrossRef](#)]
38. Carocho, M.; Ferreira, I.C.F.R. A review on antioxidants, prooxidants and related controversy: Natural and synthetic compounds, screening and analysis methodologies and future perspectives. *Food Chem. Toxicol.* **2013**, *51*, 15–25. [[CrossRef](#)] [[PubMed](#)]
39. Tisma, V.S.; Basta-Juzbasic, A.; Jaganjac, M.; Brcic, L.; Dobric, I.; Lipozencic, J.; Tatzber, F.; Zarkovic, N.; Poljak-Blazi, M. Oxidative stress and ferritin expression in the skin of patients with rosacea. *J. Am. Acad. Dermatol.* **2009**, *60*, 270–276. [[CrossRef](#)] [[PubMed](#)]
40. Sredoja Tisma, V.; Bulimbasic, S.; Galesic Ljubanovic, D.; Galesic, K.; Morovic-Vergles, J.; Mitrovic, J.; Uchida, K.; Tatzber, F.; Zarkovic, N.; Jaganjac, M. The Onset of Systemic Oxidative Stress Associated with the Accumulation of Lipid Peroxidation Product Acrolein in the Skin of Patients with Small-Vessel Vasculitis. *Molecules* **2021**, *26*, 2344. [[CrossRef](#)]
41. Sadeer, N.B.; Montesano, D.; Albrizio, S.; Zengin, G.; Mahomoodally, M.F. The versatility of antioxidant assays in food science and safety—Chemistry, applications, strengths, and limitations. *Antioxidants* **2020**, *9*, 709. [[CrossRef](#)]
42. Munteanu, I.G.; Apetrei, C. Analytical methods used in determining antioxidant activity: A review. *Int. J. Mol. Sci.* **2021**, *22*, 3380. [[CrossRef](#)]
43. Félix, R.; Valentão, P.; Andrade, P.B.; Félix, C.; Novais, S.C.; Lemos, M.F.L. Evaluating the in vitro potential of natural extracts to protect lipids from oxidative damage. *Antioxidants* **2020**, *9*, 231. [[CrossRef](#)]
44. Dikalov, S.I.; Harrison, D.G. Methods for detection of mitochondrial and cellular reactive oxygen species. *Antioxid. Redox Signal.* **2014**, *20*, 372–382. [[CrossRef](#)]
45. Lao, H.K.; Tan, J.; Wang, C.; Zhang, X. Ratiometric polymer probe for detection of peroxynitrite and the application for live-cell imaging. *Molecules* **2019**, *24*, 465. [[CrossRef](#)] [[PubMed](#)]
46. Peng, T.; Wong, N.-K.; Chen, X.; Chan, Y.-K.; Ho, D.H.-H.; Sun, Z.; Hu, J.J.; Shen, J.; El-Nezami, H.; Yang, D. Molecular imaging of peroxynitrite with HKGreen-4 in live cells and tissues. *J. Am. Chem. Soc.* **2014**, *136*, 11728–11734. [[CrossRef](#)]
47. Peng, T.; Chen, X.; Gao, L.; Zhang, T.; Wang, W.; Shen, J.; Yang, D. A rationally designed rhodamine-based fluorescent probe for molecular imaging of peroxynitrite in live cells and tissues. *Chem. Sci.* **2016**, *7*, 5407–5413. [[CrossRef](#)]
48. Zielonka, J.; Kalyanaraman, B. Hydroethidine- and MitoSOX-derived red fluorescence is not a reliable indicator of intracellular superoxide formation: Another inconvenient truth. *Free Radic. Biol. Med.* **2010**, *48*, 983–1001. [[CrossRef](#)] [[PubMed](#)]
49. Al-Thani, A.M.; Voss, S.C.; Al-Menhali, A.S.; Barcaru, A.; Horvatovich, P.; Al Jaber, H.; Nikolovski, Z.; Latiff, A.; Georgakopoulos, C.; Merenkov, Z.; et al. Whole blood storage in CPDA1 blood bags alters erythrocyte membrane proteome. *Oxid. Med. Cell. Longev.* **2018**, *2018*. [[CrossRef](#)] [[PubMed](#)]
50. Ludtmann, M.H.R.; Angelova, P.R.; Horrocks, M.H.; Choi, M.L.; Rodrigues, M.; Baev, A.Y.; Berezhnov, A.V.; Yao, Z.; Little, D.; Banushi, B.; et al. α -synuclein oligomers interact with ATP synthase and open the permeability transition pore in Parkinson’s disease. *Nat. Commun.* **2018**, *9*, 2293. [[CrossRef](#)]
51. Esteras, N.; Kopach, O.; Maiolino, M.; Lariccia, V.; Amoroso, S.; Qamar, S.; Wray, S.; Rusakov, D.A.; Jaganjac, M.; Abramov, A.Y. Mitochondrial ROS control neuronal excitability and cell fate in frontotemporal dementia. *Alzheimer’s Dement.* **2021**. [[CrossRef](#)]
52. Robinson, K.M.; Janes, M.S.; Pehar, M.; Monette, J.S.; Ross, M.F.; Hagen, T.M.; Murphy, M.P.; Beckman, J.S. Selective fluorescent imaging of superoxide in vivo using ethidium-based probes. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 15038–15043. [[CrossRef](#)]
53. Al-Menhali, A.S.; Jameela, S.A.; Latiff, A.A.; Elrayess, M.A.; Alsayrafi, M.; Jaganjac, M. Cistanche tubulosa induces reactive oxygen species-mediated apoptosis of primary and metastatic human colon cancer cells. *J. Appl. Pharm. Sci.* **2017**, *7*, 39–45. [[CrossRef](#)]
54. Belousov, V.V.; Fradkov, A.F.; Lukyanov, K.A.; Staroverov, D.B.; Shakhbazov, K.S.; Terskikh, A.V.; Lukyanov, S. Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat. Methods* **2006**, *3*, 281–286. [[CrossRef](#)]
55. Drummen, G.P.; Van Liebergen, L.C.; Op den Kamp, J.A.; Post, J.A. C11-BODIPY^{581/591}, an oxidation-sensitive fluorescent lipid peroxidation probe: (Micro)spectroscopic characterization and validation of methodology. *Free Radic. Biol. Med.* **2002**, *33*, 473–490. [[CrossRef](#)]

56. Al-Menhali, A.S.; Banu, S.; Angelova, P.R.; Barcaru, A.; Horvatovich, P.; Abramov, A.Y.; Jaganjac, M. Lipid peroxidation is involved in calcium dependent upregulation of mitochondrial metabolism in skeletal muscle. *Biochim. Biophys. Acta Gen. Subj.* **2020**, *1864*, 129487. [[CrossRef](#)]
57. Maraldi, T. Natural compounds as modulators of NADPH oxidases. *Oxid. Med. Cell. Longev.* **2013**. [[CrossRef](#)] [[PubMed](#)]
58. Acharya, P.; Mathur, M.C. Oxidative stress in alopecia areata: A systematic review and meta-analysis. *Int. J. Dermatol.* **2020**, *59*, 434–440. [[CrossRef](#)] [[PubMed](#)]
59. Emre, S.; Metin, A.; Demirseren, D.D.; Akoglu, G.; Oztekin, A.; Neselioglu, S.; Erel, O. The association of oxidative stress and disease activity in seborrheic dermatitis. *Arch. Dermatol. Res.* **2012**, *304*, 683–687. [[CrossRef](#)] [[PubMed](#)]
60. Kaur, S.; Zilmer, K.; Leping, V.; Zilmer, M. Allergic contact dermatitis is associated with significant oxidative stress. *Dermatol. Res. Pract.* **2014**, *2014*. [[CrossRef](#)]
61. Sarici, G.; Cinar, S.; Armutcu, F.; Altinyazar, C.; Koca, R.; Tekin, N.S. Oxidative stress in acne vulgaris. *J. Eur. Acad. Dermatol. Venereol.* **2010**, *24*, 763–767. [[CrossRef](#)]
62. Luo, J.-Y.; Liu, X.; Jiang, M.; Zhao, H.-P.; Zhao, J.-J. Oxidative stress markers in blood in systemic sclerosis: A meta-analysis. *Mod. Rheumatol.* **2017**, *27*, 306–314. [[CrossRef](#)] [[PubMed](#)]
63. Yesilova, Y.; Ucmak, D.; Selek, S.; Dertlioglu, S.B.; Sula, B.; Bozkus, F.; Turan, E. Oxidative stress index may play a key role in patients with pemphigus vulgaris. *J. Eur. Acad. Dermatol. Venereol.* **2013**, *27*, 465–467. [[CrossRef](#)] [[PubMed](#)]
64. Kilinc, F.; Sener, S.; Akbaş, A.; Metin, A.; Kirbaş, S.; Neselioglu, S.; Erel, O. Oxidative stress parameters in localized scleroderma patients. *Arch. Dermatol. Res.* **2016**, *308*, 625–629. [[CrossRef](#)]
65. Baek, J.-O.; Byamba, D.; Wu, W.H.; Kim, T.-G.; Lee, M.-G. Assessment of an imiquimod-induced psoriatic mouse model in relation to oxidative stress. *Arch. Dermatol. Res.* **2012**, *304*, 699–706. [[CrossRef](#)]
66. Richmond, J.M.; Frisoli, M.L.; Harris, J.E. Innate immune mechanisms in vitiligo: Danger from within. *Curr. Opin. Immunol.* **2013**, *25*, 676–682. [[CrossRef](#)]
67. Schallreuter, K.U.; Wood, J.M.; Berger, J. Low catalase levels in the epidermis of patients with vitiligo. *J. Investig. Dermatol.* **1991**, *97*, 1081–1085. [[CrossRef](#)]
68. Vaccaro, M.; Bagnato, G.; Cristani, M.; Borgia, F.; Spatari, G.; Tigano, V.; Saja, A.; Guarneri, F.; Cannavò, S.P.; Gangemi, S. Oxidation products are increased in patients affected by non-segmental generalized vitiligo. *Arch. Dermatol. Res.* **2017**, *309*, 485–490. [[CrossRef](#)]
69. Mehaney, D.A.; Darwish, H.A.; Hegazy, R.A.; Nooh, M.M.; Tawdy, A.M.; Gawdat, H.I.; El-Sawalhi, M.M. Analysis of oxidative stress status, catalase and catechol-O-Methyltransferase polymorphisms in Egyptian vitiligo patients. *PLoS ONE* **2014**, *9*, e99286. [[CrossRef](#)] [[PubMed](#)]
70. Battino, M.; Greabu, M.; Totan, A.; Bullon, P.; Bucur, A.; Tovar, S.; Mohora, M.; Didilescu, A.; Parlatescu, I.; Spinu, T.; et al. Oxidative stress markers in oral lichen planus. *BioFactors* **2008**, *33*, 301–310. [[CrossRef](#)]
71. Budzyń, M.; Iskra, M.; Krasiński, Z.; Dzieciuchowicz, L.; Kasprzak, M.; Gryszczyńska, B. Serum iron concentration and plasma oxidant/antioxidant balance in patients with chronic venous insufficiency. *Med. Sci. Monit.* **2011**, *17*, CR719. [[CrossRef](#)]
72. Li, W.C.; Mo, L.J.; Shi, X.; Lin, Z.Y.; Li, Y.Y.; Yang, Z.; Wu, C.L.; Li, X.H.; Luo, Y.Z.; Qin, L.Q.; et al. Antioxidant status of serum bilirubin, uric acid and albumin in pemphigus vulgaris. *Clin. Exp. Dermatol.* **2018**, *43*, 158–163. [[CrossRef](#)] [[PubMed](#)]
73. Sivaranjani, N.; Venkata Rao, S.; Rajeev, G. Role of reactive oxygen species and antioxidants in atopic dermatitis. *J. Clin. Diagn. Res.* **2013**, *7*, 2683–2685. [[CrossRef](#)]
74. Chen, J.; Li, S.; Li, C. Mechanisms of melanocyte death in vitiligo. *Med. Res. Rev.* **2021**, *41*, 1138–1166. [[CrossRef](#)]
75. Botes, L.; Van Der Westhuizen, F.H.; Loots, D.T. Phytochemical contents and antioxidant capacities of two *Aloe gratifolia* var. *davyana* extracts. *Molecules* **2008**, *13*, 2169–2180. [[CrossRef](#)]
76. Gao, Y.; Yuan, J.; Liang, X.; He, Y.; Li, P.; Yang, M. *Artemisia anomala* extracts enhance the viability and anti-oxidation capacity of human keratinocytes. *Trop. J. Pharm. Res.* **2019**, *18*, 61–67. [[CrossRef](#)]
77. Kim, B.-H.; Oh, I.; Kim, J.-H.; Jeon, J.-E.; Jeon, B.; Shin, J.; Kim, T.-Y. Anti-inflammatory activity of compounds isolated from *Astragalus sinicus* L. in cytokine-induced keratinocytes and skin. *Exp. Mol. Med.* **2014**, *46*, e87. [[CrossRef](#)]
78. Roy, S.; Khanna, S.; Alessio, H.M.; Vider, J.; Bagchi, D.; Bagchi, M.; Sen, C.K. Anti-angiogenic property of edible berries. *Free Radic. Res.* **2002**, *36*, 1023–1032. [[CrossRef](#)]
79. García-Pérez, M.-E.; Royer, M.; Duque-Fernandez, A.; Diouf, P.N.; Stevanovic, T.; Pouliot, R. Antioxidant, toxicological and antiproliferative properties of Canadian polyphenolic extracts on normal and psoriatic keratinocytes. *J. Ethnopharmacol.* **2010**, *132*, 251–258. [[CrossRef](#)]
80. Bader, A.; Martini, F.; Schinella, G.R.; Rios, J.L.; Prieto, J.M. Modulation of Cox-1, 5-, 12- And 15-Lox by popular herbal remedies used in Southern Italy against psoriasis and other skin diseases. *Phyther. Res.* **2015**, *29*, 108–113. [[CrossRef](#)]
81. Abe, S.; Ueno, M.; Nishitani, M.; Akamatsu, T.; Sato, T.; Shimoda, M.; Kanaoka, H.; Nii, Y.; Yamasaki, H.; Yuasa, K. Citrus sudachi peel extract suppresses cell proliferation and promotes the differentiation of keratinocytes through inhibition of the EGFR-ERK signaling pathway. *Biomolecules* **2020**, *10*, 1468. [[CrossRef](#)]
82. Gelmini, F.; Beretta, G.; Anselmi, C.; Centini, M.; Magni, P.; Ruscica, M.; Cavalchini, A.; Maffei Facino, R. GC-MS profiling of the phytochemical constituents of the oleoresin from *Copaifera langsdorffii* Desf. and a preliminary in vivo evaluation of its antipsoriatic effect. *Int. J. Pharm.* **2013**, *440*, 170–178. [[CrossRef](#)]

83. Yang, B.-Y.; Cheng, Y.-G.; Liu, Y.; Liu, Y.; Tan, J.-Y.; Guan, W.; Guo, S.; Kuang, H.-X. Datura Metel L. Ameliorates imiquimod-induced psoriasis-like dermatitis and inhibits inflammatory cytokines production through TLR7/8-MyD88-NF- κ B-NLRP3 inflammasome pathway. *Molecules* **2019**, *24*, 2157. [[CrossRef](#)] [[PubMed](#)]
84. Bito, T.; Roy, S.; Sen, C.K.; Packer, L. Pine bark extract pycnogenol downregulates IFN- γ -induced adhesion of T cells to human keratinocytes by inhibiting inducible ICAM-1 expression. *Free Radic. Biol. Med.* **2000**, *28*, 219–227. [[CrossRef](#)]
85. Singh, S.K.; Chouhan, H.S.; Sahu, A.N.; Narayan, G. Assessment of in vitro antipsoriatic activity of selected Indian medicinal plants. *Pharm. Biol.* **2015**, *53*, 1295–1301. [[CrossRef](#)]
86. Dimitris, D.; Ekaterina-Michaela, T.; Christina, K.; Ioannis, S.; Ioanna, S.K.; Aggeliki, L.; Sophia, H.; Michael, R.; Helen, S. Melissa officinalis ssp. altissima extracts: A therapeutic approach targeting psoriasis in mice. *J. Ethnopharmacol.* **2020**, *246*. [[CrossRef](#)] [[PubMed](#)]
87. Ampawong, S.; Kengkoom, K.; Sukphopetch, P.; Aramwit, P.; Muangkaew, W.; Kanjanapruthipong, T.; Buaban, T. Evaluating the effect of rice (*Oryza sativa* L.: SRNC05053-6-2) crude extract on psoriasis using in vitro and in vivo models. *Sci. Rep.* **2020**, *10*, 17618. [[CrossRef](#)] [[PubMed](#)]
88. Ndjoubi, K.O.; Sharma, R.; Badmus, J.A.; Jacobs, A.; Jordaan, A.; Marnewick, J.; Warner, D.F.; Hussein, A.A. Antimycobacterial, cytotoxic, and antioxidant activities of abietane diterpenoids isolated from *Plectranthus madagascariensis*. *Plants* **2021**, *10*, 175. [[CrossRef](#)]
89. Parmar, K.M.; Itankar, P.R.; Joshi, A.; Prasad, S.K. Anti-psoriatic potential of *Solanum xanthocarpum* stem in Imiquimod-induced psoriatic mice model. *J. Ethnopharmacol.* **2017**, *198*, 158–166. [[CrossRef](#)]
90. Abbas, A.N. Ginger (*Zingiber officinale* (L.) Rosc) improves oxidative stress and trace elements status in patients with alopecia areata. *Niger. J. Clin. Pract.* **2020**, *23*, 1555–1560. [[CrossRef](#)]
91. Pekmezci, E.; Dundar, C.; Turkoglu, M. Proprietary Herbal Extract Downregulates the Gene Expression of IL-1 α in HaCaT Cells: Possible Implications Against Nonscarring Alopecia. *Med. Arch. (Sarajevobosnia Herzeg.)* **2018**, *72*, 136–140. [[CrossRef](#)]
92. León, L.M.; García, J.C.L.; Duarte, C.C.; García, K.G.; Guerra, I.R.; Marín, R.M. Antioxidant capacity of *clusia minor* l. Leaves | Capacidad antioxidante de las hojas de la especie *clusia minor* l. *Rev. Cuba. Plantas Med.* **2020**, *25*.
93. Dammak, I.; Boudaya, S.; Abdallah, F.B.; Hamida, T.; Attia, H. Date seed Oil inhibits hydrogen peroxide-induced oxidative stress in normal human epidermal melanocytes. *Connect. Tissue Res.* **2009**, *50*, 330–335. [[CrossRef](#)] [[PubMed](#)]
94. Yang, L.; Yang, F.; Teng, L.; Katayama, I. 6-shogaol protects human melanocytes against oxidative stress through activation of the nrf2-antioxidant response element signaling pathway. *Int. J. Mol. Sci.* **2020**, *21*, 3537. [[CrossRef](#)]
95. Lu, L.; Wang, S.; Fu, L.; Liu, D.; Zhu, Y.; Xu, A. Bilobalide protection of normal human melanocytes from hydrogen peroxide-induced oxidative damage via promotion of antioxidant expression and inhibition of endoplasmic reticulum stress. *Clin. Exp. Dermatol.* **2016**, *41*, 64–73. [[CrossRef](#)]
96. Jeong, Y.-M.; Choi, Y.-G.; Kim, D.-S.; Park, S.-H.; Yoon, J.-A.; Kwon, S.-B.; Park, E.-S.; Park, K.-C. Cytoprotective effect of green tea extract and quercetin against hydrogen peroxide-induced oxidative stress. *Arch. Pharm. Res.* **2005**, *28*, 1251–1256. [[CrossRef](#)]
97. Ning, W.; Wang, S.; Dong, X.; Liu, D.; Fu, L.; Jin, R.; Xu, A. Epigallocatechin-3-gallate (EGCG) suppresses the trafficking of lymphocytes to epidermal melanocytes via inhibition of JAK2: Its implication for vitiligo treatment. *Biol. Pharm. Bull.* **2015**, *38*, 1700–1706. [[CrossRef](#)]
98. Moreira, C.G.; Carrenho, L.Z.B.; Pawloski, P.L.; Soley, B.S.; Cabrini, D.A.; Otuki, M.F. Pre-clinical evidences of *Pyrostegia venusta* in the treatment of vitiligo. *J. Ethnopharmacol.* **2015**, *168*, 315–325. [[CrossRef](#)]
99. Ma, J.; Li, S.; Zhu, L.; Guo, S.; Yi, X.; Cui, T.; He, Y.; Chang, Y.; Liu, B.; Li, C.; et al. Baicalein protects human vitiligo melanocytes from oxidative stress through activation of NF-E2-related factor2 (Nrf2) signaling pathway. *Free Radic. Biol. Med.* **2018**, *129*, 492–503. [[CrossRef](#)]
100. Lai, Y.; Feng, Q.; Zhang, R.; Shang, J.; Zhong, H. The great capacity on promoting melanogenesis of three compatible components in *vernonia anthelmintica* (L.) willd. *Int. J. Mol. Sci.* **2021**, *22*, 4073. [[CrossRef](#)]
101. Kim, H.; Ban, I.; Choi, Y.; Yu, S.; Youn, S.J.; Baik, M.-Y.; Lee, H.; Kim, W. Puffing of turmeric (*Curcuma longa* L.) enhances its anti-inflammatory effects by upregulating macrophage oxidative phosphorylation. *Antioxidants* **2020**, *9*, 931. [[CrossRef](#)]
102. Vaughn, A.R.; Pourang, A.; Clark, A.K.; Burney, W.; Sivamani, R.K. Dietary supplementation with turmeric polyherbal formulation decreases facial redness: A randomized double-blind controlled pilot study. *J. Integr. Med.* **2019**, *17*, 20–23. [[CrossRef](#)]
103. Bhatt, L.R.; Lim, J.A.; Chai, K.Y.; Kang, J.I.; Oh, H.K.; Baek, S.H. Antioxidative and antimicrobial activities of essential oil from *Artemisia vulgaris*. *Nat. Prod. Sci.* **2006**, *12*, 226–231.
104. Chrzaszcz, M.; Miazga-Karska, M.; Klimek, K.; Granica, S.; Tchórzewska, D.; Ginalska, G.; Szewczyk, K. Extracts from *cephalaria uralensis* (Murray) roem. & schult. and *cephalaria gigantea* (Ledeb.) bobrov as potential agents for treatment of acne vulgaris: Chemical characterization and in vitro biological evaluation. *Antioxidants* **2020**, *9*, 796. [[CrossRef](#)]
105. De Canha, M.N.; Kishore, N.; Kumar, V.; Meyer, D.; Nehar, S.; Singh, B.; Lall, N. The potential of *Clausena anisata* (Willd.) Hook.f. ex Benth against *Propionibacterium acnes*. *S. Afr. J. Bot.* **2018**, *119*, 410–419. [[CrossRef](#)]
106. Yamaguchi, N.; Satoh-Yamaguchi, K.; Ono, M. In vitro evaluation of antibacterial, anticollagenase, and antioxidant activities of hop components (*Humulus lupulus*) addressing acne vulgaris. *Phytomedicine* **2009**, *16*, 369–376. [[CrossRef](#)]
107. Matsubara, Y.; Matsumoto, T.; Sekiguchi, K.; Koseki, J.; Kaneko, A.; Yamaguchi, T.; Kurihara, Y.; Kobayashi, H.; Iriti, M. Oral administration of the Japanese traditional medicine Keishibukuryogan-ka-yokuinin decreases reactive oxygen metabolites in rat plasma: Identification of chemical constituents contributing to antioxidant activity. *Molecules* **2017**, *22*, 256. [[CrossRef](#)] [[PubMed](#)]

108. Poomanee, W.; Chaiyana, W.; Mueller, M.; Viernstein, H.; Khunkitti, W.; Leelapornpisid, P. In-vitro investigation of anti-acne properties of *Mangifera indica* L. kernel extract and its mechanism of action against *Propionibacterium acnes*. *Anaerobe* **2018**, *52*, 64–74. [[CrossRef](#)]
109. Kim, S.S.; Kim, J.E.; Hyun, C.-G.; Lee, N.H. Neolitea aciculata essential oil inhibits drug-resistant skin pathogen growth and *Propionibacterium acnes*-induced inflammatory effects of human monocyte leukemia. *Nat. Prod. Commun.* **2011**, *6*, 1193–1198. [[CrossRef](#)]
110. Chuang, L.-T.; Tsai, T.-H.; Lien, T.-J.; Huang, W.-C.; Liu, J.-J.; Chang, H.; Chang, M.-L.; Tsai, P.-J. Ethanolic extract of *origanum vulgare* suppresses *propionibacterium acnes*-induced inflammatory responses in human monocyte and mouse ear edema models. *Molecules* **2018**, *23*, 1987. [[CrossRef](#)]
111. Kok, J.M.-L.; Jee, J.-M.; Chew, L.-Y.; Wong, C.-L. The potential of the brown seaweed *Sargassum polycystum* against *acne vulgaris*. *J. Appl. Phycol.* **2016**, *28*, 3127–3133. [[CrossRef](#)]
112. Seong, S.J.; Su, K.J.; Sung, G.K.; Choi, J.-S.; Kwang, W.H.; Do, I.L. Anti-acne activity of *Selaginella involvens* extract and its non-antibiotic antimicrobial potential on *Propionibacterium acnes*. *Phyther. Res.* **2008**, *22*, 335–339. [[CrossRef](#)]
113. Sharma, R.; Kishore, N.; Hussein, A.; Lall, N. Antibacterial and anti-inflammatory effects of *Syzygium jambos* L. (Alston) and isolated compounds on *acne vulgaris*. *BMC Complement. Altern. Med.* **2013**, *13*, 1–10. [[CrossRef](#)]
114. Kalaskar, A.R.; Bhowate, R.R.; Kalaskar, R.R.; Ghonmode, S. Novel neem leaves extract mouthwash therapy for oral lichen planus. *J. Herb. Med.* **2021**, *26*, 100408. [[CrossRef](#)]
115. Agha-Hosseini, F.; Borhan-Mojabi, K.; Monsef-Esfahani, H.-R.; Mirzaii-Dizgah, I.; Etemad-Moghadam, S.; Karagah, A. Efficacy of purslane in the treatment of oral lichen planus. *Phyther. Res.* **2010**, *24*, 240–244. [[CrossRef](#)]
116. Grau, M.; Bölck, B.; Bizjak, D.A.; Stabenow, C.J.A.; Bloch, W. The red-vine-leaf extract AS195 increases nitric oxide synthase-dependent nitric oxide generation and decreases oxidative stress in endothelial and red blood cells. *Pharmacol. Res. Perspect.* **2016**, *4*, e00213. [[CrossRef](#)] [[PubMed](#)]
117. De Almeida Cyrino, F.Z.G.; Balthazar, D.S.; Sicuro, F.L.; Bouskela, E. Effects of venotonic drugs on the microcirculation: Comparison between *Ruscus* extract and micronized diosmine. *Clin. Hemorheol. Microcirc.* **2018**, *68*, 361–370. [[CrossRef](#)]
118. Akter, K.; Barnes, E.C.; Loa-Kum-Cheung, W.L.; Yin, P.; Kichu, M.; Brophy, J.J.; Barrow, R.A.; Imchen, I.; Vemulpad, S.R.; Jamie, J.F. Antimicrobial and antioxidant activity and chemical characterisation of *Erythrina stricta* Roxb. (Fabaceae). *J. Ethnopharmacol.* **2016**, *185*, 171–181. [[CrossRef](#)]
119. Fu, R.; Zhang, Y.-T.; Guo, Y.-R.; Huang, Q.-L.; Peng, T.; Xu, Y.; Tang, L.; Chen, F. Antioxidant and anti-inflammatory activities of the phenolic extracts of *Sapium sebiferum* (L.) Roxb. leaves. *J. Ethnopharmacol.* **2013**, *147*, 517–524. [[CrossRef](#)] [[PubMed](#)]
120. Arumugam, R.; Sarikurcu, C.; Mutlu, M.; Tepe, B. *Sophora alopecuroides* var. *alopecuroides*: Phytochemical composition, antioxidant and enzyme inhibitory activity of the methanolic extract of aerial parts, flowers, leaves, roots, and stems. *S. Afr. J. Bot.* **2020**. [[CrossRef](#)]
121. Gveric-Ahmetasevic, S.; Sunjic, S.B.; Skala, H.; Andrisic, L.; Stroser, M.; Zarkovic, K.; Skrablin, S.; Tatzber, F.; Cipak, A.; Jaganjac, M.; et al. Oxidative stress in small-for-gestational age (SGA) term newborns and their mothers. *Free Radic. Res.* **2009**, *43*, 376–384. [[CrossRef](#)] [[PubMed](#)]
122. Cesar, V.; Jozić, I.; Begović, L.; Vuković, T.; Mlinarić, S.; Lepeduš, H.; Šunjić, S.B.; Žarković, N. Cell-type-specific modulation of hydrogen peroxide cytotoxicity and 4-hydroxynonenal binding to human cellular proteins in vitro by antioxidant aloe vera extract. *Antioxidant* **2018**, *7*, 125. [[CrossRef](#)] [[PubMed](#)]