

# The $\theta$ -angle and axion physics of two-color QCD at fixed baryon charge

Jahmall Bersini,<sup>a,f</sup> Alessandra D'Alise,<sup>b</sup> Francesco Sannino<sup>b,c,d,e</sup> and Matías Torres<sup>b</sup>

<sup>a</sup>*Rudjer Boskovic Institute, Division of Theoretical Physics,  
Bijenička 54, Zagreb 10000, Croatia*

<sup>b</sup>*Dipartimento di Fisica “E. Pancini”, Università di Napoli Federico II and  
INFN sezione di Napoli, Complesso Universitario di Monte S. Angelo,  
Edificio 6, via Cintia, Napoli 80126, Italy*

<sup>c</sup>*Scuola Superiore Meridionale,  
Largo S. Marcellino 10, Napoli 80138, Italy*

<sup>d</sup>*CP<sup>3</sup>-Origins and D-IAS, University of Southern Denmark,  
Campusvej 55, Odense M 5230, Denmark*

<sup>e</sup>*CERN, Theoretical Physics Department,  
Geneva 23 1211, Switzerland*

<sup>f</sup>*Kavli IPMU (WPI), UTIAS, The University of Tokyo,  
Kashiwa, Chiba 277-8583, Japan*

*E-mail:* [jbersini@irb.hr](mailto:jbersini@irb.hr), [alessandra.dalise@unina.it](mailto:alessandra.dalise@unina.it),  
[sannino@cp3.sdu.dk](mailto:sannino@cp3.sdu.dk), [matiasignacio.torressandoval@unina.it](mailto:matiasignacio.torressandoval@unina.it)

**ABSTRACT:** We analyze the impact of the  $\theta$ -angle and axion dynamics for two-color (in fact any  $\text{Sp}(2N)$ ) QCD at nonzero baryon charge and as a function of the number of matter fields on the vacuum properties, the pattern of chiral symmetry breaking as well as the spectrum of the theory. We show that the vacuum acquires a rich structure when the underlying  $CP$  violating topological operator is added to the theory. We discover novel phases and analyse the order of their transitions characterizing the dynamics of the odd and even number of flavours. We further determine the critical chemical potential as function of the  $\theta$  angle separating the normal from the superfluid phase of the theory. Our results will guide numerical simulations and novel tests of the model's dynamics. The results are also expected to better inform phenomenological applications of the model ranging from composite Higgs physics to strongly interacting massive dark matter models featuring number changing interactions. In the companion work [1] we repurpose and upgrade the approach to determine the impact of the  $\theta$ -angle and axion physics on non-perturbative near conformal dynamics related to the fixed baryon charge sector.

**KEYWORDS:** Chiral Lagrangian, Finite Temperature or Finite Density, Global Symmetries, Spontaneous Symmetry Breaking

ARXIV EPRINT: [2208.09226](https://arxiv.org/abs/2208.09226)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Two-color chiral Lagrangian</b>	<b>3</b>
2.1	Baryon charge	4
2.2	The $\theta$ -angle and the U(1) problem	4
2.3	The axion and the baryon charge	5
<b>3</b>	<b>Determining the vacuum</b>	<b>6</b>
3.1	$N_f = 2$	9
3.2	$N_f = 3$	11
3.3	$N_f = 6$	12
3.4	$N_f = 7$	14
3.5	$N_f = 8$	15
3.6	Considerations for general $N_f$	17
3.7	Ground state energy with the axion	18
<b>4</b>	<b>Symmetry-breaking pattern and spectrum</b>	<b>19</b>
4.1	Symmetry-breaking pattern without the axion	19
4.2	Symmetry-breaking pattern with the axion	20
4.3	Fluctuations spectrum	21
<b>5</b>	<b>Conclusions</b>	<b>22</b>

---

## 1 Introduction

Understanding the dynamics of strongly coupled theories remains an open challenge making it one of the most fascinating topics in theoretical physics. Partial progress has been made both analytically and numerically via a number of tools developed in the past decades. Quantum Chromo Dynamics (QCD), featuring three colors and several light flavors, has been chosen by Nature as one of the pillars of the Standard Model of particle interactions. QCD therefore stands out within the landscape of strongly coupled theories. However, several relevant questions remain unanswered about its dynamics from the QCD physical spectrum to the interplay between chiral symmetry breaking and confinement, to its phase structure as a function of light matter fields (conformal window), temperature and matter density. For example, knowledge about the phase diagram informs a number of physical applications ranging from the inner structure of Neutron stars [2] to the thermal history of the early Universe [3] and last but not least to applications beyond standard model physics and cosmology (see [4, 5] and references therein). Another unsolved puzzle is the absence or unexplained strong suppression [6, 7] of the otherwise legitimate presence in the theory of

the topological term responsible for strong  $CP$  violation. Its possible resolutions in terms of either the axion physics [8–15] or its alternatives such as the ones in which  $CP$  is broken spontaneously [16–18] or models in which the emergent Goldstone boson is not an axion (which would have a non-zero mass) [19] (see also [20]) are under active experimental and theoretical investigation.

Reducing the number of colors from three to two provides a number of theoretical advantages and increases potential phenomenological applications beyond the QCD template. This is due, in a nutshell, to the symplectic nature of the matter representation that enhances the quantum global symmetries for the light flavors from the  $SU(N_f) \times SU(N_f) \times U(1)_B$  to  $SU(2N_f)$ . The enhancement of the quantum global symmetries (demonstrated on the lattice in [21, 22]) has far-reaching consequences [23, 24], such as the possibility of the minimal construction of composite Goldstone Higgs theories [25] to the natural introduction of number changing operators (stemming from the Wess-Zumino term) crucial to models in which dark matter genesis occurs within the dark sector itself [26–29]. When discussing non-zero baryon charge the model allows for well defined lattice simulations because the action remains real, differently from ordinary QCD. For a review of the various applications of the theory beyond the ones mentioned here we refer to [4, 5]. Last but not least this model together with ordinary QCD has been one of the most studied theories on the lattice as function of light flavors in the hunt for the predicted lower edge of the conformal window [30–33] and its dynamical properties [21, 22, 34, 34–44], finite baryon density [45–65] and last but not least the investigation of gravity-free asymptotic safety at large number of matter fields (proven rigorously in [66, 67] for gauge-Yukawa theories) and suggested for gauge-fermion theories in [68] investigated on the lattice in [69, 70].

The goal of our work is to go beyond the state-of-the-art by providing an in depth analysis of the  $\theta$ -angle and axion physics at non-zero baryon chemical potential of two-color QCD. Earlier studies appeared in [71]. We organise our work by introducing in the next section the two-color effective pion Lagrangian at non-zero baryon charge, including both the  $\theta$ -angle term as well as the axion field. In section 3 we will determine the vacuum structure of the theory both in the normal and superfluid phase as a function of the different number of matter fields. Here we will show how new phases emerge due to the interplay of the  $\theta$ -angle term, quark masses, and baryon chemical potential. We will also characterise the type of phase transitions. In particular, we will study the dynamics of  $N_f = 2, 3, 6, 7, 8$  flavors and unveil general properties depending on the even versus odd number of flavors. We stop at eight flavours since around and above this number one expects the dynamics to either become near conformal or conformal [33]. The analysis for near conformal dynamics is performed in [1]. Here we discover that in the case of even flavors we observe a first order  $CP$  phase transition for  $\theta = \pi$  both in the normal and superfluid phase, except for the case of  $N_f = 2$  for which in the superfluid case the transition disappears as also discussed in [71]. For the odd case, the normal phase still supports  $CP$  breaking at  $\theta = \pi$  while in the superfluid phase the  $\theta = \pi$  is replaced by novel first order phase transitions for  $\theta = \frac{\pi}{2}$  and  $\theta = \frac{3\pi}{2}$ . We summarize the patterns of chiral symmetry breaking in the normal and superfluid phases and determine the spectrum and associated dispersion relations in section 4. We offer our conclusions in section 5.

## 2 Two-color chiral Lagrangian

The Lagrangian of  $N_f$  Dirac fermions transforming according to the fundamental representation of two-color QCD reads:

$$\mathcal{L} = -\frac{1}{4g^2} \vec{G}_{\mu\nu} \cdot \vec{G}^{\mu\nu} + i\bar{Q}\bar{\sigma}^\nu \left[ \partial_\nu - i\vec{G}_\nu \cdot \frac{\vec{\tau}}{2} \right] Q - \frac{1}{2} m_q Q^T \tau_2 E Q + \text{h.c.} \quad (2.1)$$

Here  $G_{\mu\nu}^a$  and  $G_\mu^a$  with  $a = 1, 2, 3$  are respectively the gluon field strength and the field itself,  $\tau^a$  are the Pauli matrices for the SU(2) color group, and  $Q_\alpha^{c,i}$  is a two-spinor fermion field transforming according to the fundamental representation of color with  $c = 1, 2$  and  $i = 1, \dots, 2N_f$ . In terms of  $q_{L,R}$ , which are the original left and right handed quarks stemming from the Dirac notation it reads

$$Q = \begin{pmatrix} q_L \\ i\sigma_2 \tau_2 q_R^* \end{pmatrix}. \quad (2.2)$$

At zero fermion mass (i.e.  $m_q = 0$ ) the theory exhibits the classical  $U(2N_f)$  symmetry broken at the quantum level by the Adler-Bell-Jackiw anomaly to  $SU(2N_f)$ . The Dirac mass term breaks explicitly the symmetry to  $Sp(2N_f)$  and the  $2N_f \times 2N_f$  matrix  $E$  reads:

$$E = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \otimes \mathbb{1}_{N_f}. \quad (2.3)$$

At small number of flavors the dynamics is expected to be strong enough for a fermion condensate to form breaking the  $SU(2N_f)$  global symmetry spontaneously to a smaller subgroup expected to be the maximal diagonal one. However, only recently, first-principle lattice simulations have been able to demonstrate this pattern of chiral symmetry breaking in [21] which has been further confirmed in [22] for two Dirac flavors. Increasing the number of flavors the dynamics can change and one expects the existence of a critical number of flavors  $N_f^*$  above which an IR interacting conformal fixed point emerges. This dynamics has been investigated theoretically [33] and via first principle lattice simulations [72]. A recent up-to-date review can be found in [5]. In fact, even the dynamics at very large number of flavours is extremely interesting both theoretically and phenomenologically. It is also being investigated on the lattice [69] searching for the existence of an interacting non-perturbative UV fixed point [68] similar to the one shown to exist in [66, 67] for gauge-Yukawa theories. For any  $2 < N_f < N_f^*$  it is therefore natural to expect the pattern of spontaneous symmetry breaking to be  $SU(2N_f) \rightarrow Sp(2N_f)$ . Generalizing this theory to arbitrary  $N$  while keeping the same global symmetries requires considering  $Sp(2N)$  gauge groups and the associated conformal window has been discussed in [33]. The associated chiral Lagrangian embodying the previous pattern of symmetry breaking reads [4, 23]:

$$\mathcal{L}_{\text{eff}} = \nu^2 \text{Tr}\{\partial_\mu \Sigma \partial^\mu \Sigma^\dagger\} + m_\pi^2 \nu^2 \text{Tr}\{M\Sigma + M^\dagger \Sigma^\dagger\}, \quad (2.4)$$

with  $\Sigma$  transforming linearly under a chiral rotation as

$$\Sigma \rightarrow u \Sigma u^T, \quad u \in SU(2N_f), \quad (2.5)$$

and the democratic mass matrix  $M$  being

$$M = -i\sigma_2 \otimes \mathbb{1}_{N_f} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \otimes \mathbb{1}_{N_f}. \quad (2.6)$$

## 2.1 Baryon charge

We can take into account a non-vanishing baryon charge by coupling it to a chemical potential  $\mu$ . The latter can be introduced as the zero component of a background gauge field via the covariant derivative

$$\partial_\mu \mapsto D_\mu = \partial_\mu - i\mu\delta_\mu^0 B, \quad B \equiv \begin{pmatrix} 1/2 & 0 \\ 0 & -1/2 \end{pmatrix} \otimes \mathbb{1}_{N_f} \quad (2.7)$$

which reproduces the usual coupling of the chemical potential to the Noether charge in the Dirac notation. In fact

$$\bar{q}\gamma^0 q = \begin{pmatrix} q_L^* \\ q_R^* \end{pmatrix}^T \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \mathbb{1}_{N_f} \begin{pmatrix} q_L \\ q_R \end{pmatrix} = \underbrace{Q^\dagger \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes \mathbb{1}_{N_f}}_{\equiv 2B} Q = 2Q^\dagger B Q. \quad (2.8)$$

Notice that the baryon charge generator belongs to the  $su(2N_f)$  algebra being proportional to the third Pauli matrix.

From the form of the  $B$  matrix, we see that for non-zero  $\mu$  the Lagrangian is no longer invariant under  $SU(2N_f)$  transformations. To fix the ideas, when the mass term is zero the  $SU(2N_f)$  symmetry is explicitly broken as

$$SU(2N_f) \xrightarrow{m=0, \mu \neq 0} SU(N_f)_L \times SU(N_f)_R \times U(1)_B, \quad (2.9)$$

conversely, for  $m \neq 0$  one has

$$SU(2N_f) \xrightarrow{m \neq 0, \mu \neq 0} SU(N_f)_V \times U(1)_B. \quad (2.10)$$

By using the covariant derivative (2.7) into eq. (2.4), we obtain the effective Lagrangian describing the theory at non-zero baryon charge

$$\begin{aligned} \mathcal{L}_B = & \nu^2 \text{Tr}\{\partial_\mu \Sigma \partial^\mu \Sigma^\dagger\} + 4\mu\nu^2 \text{Tr}\{B \Sigma^\dagger \partial_0 \Sigma\} + m_\pi^2 \nu^2 \text{Tr}\{M \Sigma + M^\dagger \Sigma^\dagger\} \\ & + 2\mu^2 \nu^2 \left[ \text{Tr}\{\Sigma B^T \Sigma^\dagger B\} + \text{Tr}\{B B\} \right]. \end{aligned} \quad (2.11)$$

## 2.2 The $\theta$ -angle and the $U(1)$ problem

In order to discuss the physics of the  $\theta$ -angle and of the axial anomaly [73–75], we introduce the topological charge density

$$q(x) = \frac{g^2}{64\pi^2} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu}^a F_{\rho\sigma}^a, \quad (2.12)$$

which we incorporate in the effective Lagrangian as

$$\mathcal{L}_{q(x)} = \frac{i}{4} q(x) \text{Tr}\{\log \Sigma - \log \Sigma^\dagger\} - \theta q(x) + \frac{q(x)^2}{4a\nu^2}. \quad (2.13)$$

	[SU(2)]	SU( $N_f$ ) <sub>L</sub> ×	SU( $N_f$ ) <sub>R</sub> ×	U(1) <sub>V</sub> ×	U(1) <sub>A</sub>
$q_L$	□	□	<b>1</b>	+1	+1
$i\sigma_2\tau_2q_R^*$	□	<b>1</b>	$\bar{\square}$	-1	+1
	[SU(2)]	SU(2 $N_f$ ) ×	U(1) <sub>A</sub>		
$\mathcal{Q}$	□	□	+1		

**Table 1.** Transformation properties of  $q_L$ ,  $i\sigma_2\tau_2q_R^*$  and  $\mathcal{Q}$  under the action of the symmetry groups.

Here the new  $\Sigma$  is related to the old one that transforms only under SU(2 $N_f$ ) via [76]

$$\Sigma \rightarrow \Sigma e^{i\frac{S}{\sqrt{2N_f}}\mathbb{1}_{2N_f}}, \tag{2.14}$$

with the  $S$ -field a singlet of SU(2 $N_f$ ) transforming under the anomalous U(1)<sub>A</sub>, and it is parent to the  $\eta'$  particle in ordinary QCD.

The minimal choice to neglect orders higher than  $q^2(x)$  is justified at large number of colors, see discussion in [76, 77]. Here, to keep the same pattern of chiral symmetry breaking the SU(2) of color generalizes to Sp(2 $N$ ) and not SU( $N$ ). The linear term allows accommodating for the U(1)<sub>A</sub> anomaly. The coefficient of the quadratic term in  $q(x)$  is known as the topological susceptibility of the Yang-Mills theory. The coefficients of these terms are coherently chosen such that we reproduce the axial anomaly  $\partial_\mu J_5^\mu = 4N_f q(x)$  in two-color QCD with quarks in the fundamental representation [76]. Notice that  $q(x)$  is a background auxiliary field introduced to implement the anomalous transformation at the action level in linear fashion. It can, therefore, be integrated out via its equations of motion, yielding the following effective Lagrangian

$$\begin{aligned} \mathcal{L}_\theta = & \nu^2 \text{Tr}\{\partial_\mu \Sigma \partial^\mu \Sigma^\dagger\} + 4\mu\nu^2 \text{Tr}\{B\Sigma^\dagger \partial_0 \Sigma\} + m_\pi^2 \nu^2 \text{Tr}\{M\Sigma + M^\dagger \Sigma^\dagger\} \\ & + 2\mu^2 \nu^2 \left[ \text{Tr}\{\Sigma B^T \Sigma^\dagger B\} + \text{Tr}\{BB\} \right] - a\nu^2 \left( \theta - \frac{i}{4} \text{Tr}\{\log \Sigma - \log \Sigma^\dagger\} \right)^2. \end{aligned} \tag{2.15}$$

The action of the symmetry groups is summarised in table 1 where the axial generator  $A$  in our notation is

$$A = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \mathbb{1}_{N_f}. \tag{2.16}$$

### 2.3 The axion and the baryon charge

For the Standard Model three colors QCD there is no experimental evidence of strong  $CP$  violation. This constrains the associated value of the effective theta angle  $\bar{\theta} = \theta + \arg \det(M)$ , with  $M$  the physical quark mass matrix, to be  $\bar{\theta} < 10^{-10}$ . The limit comes from the bound on the neutron electric dipole moment  $|d_n| = C_{\text{EDM}} e \bar{\theta} < 1.8 \times 10^{-26} e \text{ cm}$  [6, 7], where  $C_{\text{EDM}} = 2.4(1.0) \times 10^{-16} \text{ cm}$  [78] is related to the effective nucleon interactions with the axion explained in the appendix of [76]. From a theoretical viewpoint, a tiny value of a physical parameter unprotected by any symmetry requires an explanation. One solution for the so-called strong  $CP$  problem was proposed in the 70s by R. Peccei and H. Quinn [8, 9].

The proposal makes use of an additional  $U(1)_{PQ}$  symmetry which is quantum mechanically anomalous and spontaneously broken leading to the axion as an extra (pseudo-)Goldstone boson. Axion physics has led to a great deal of both theoretical and phenomenological work including active experimental searches beyond the original QCD axion [79]. An alternative solution to the strong  $CP$  problem featuring new composite dynamics and heavier (if not completely absent) axions was proposed in [19] and reconsidered in [20] and applied extensively in model building in [80–82].

Of course, when discussing the  $\theta$ -angle physics here, depending on the physical application of the model, we can allow for new sources of  $CP$  breaking, perhaps relevant for some models of dark matter or Standard Model secluded sectors [79]. Nevertheless, it is interesting to entertain the possibility that an axion is present in the theory and therefore explore its finite chemical potential dynamics and impact. We denote by  $\nu_{PQ}$  the scale of  $U(1)_{PQ}$  spontaneous symmetry breaking and by  $a_{PQ}$  the coefficient of the  $U(1)_{PQ}$  anomalous term. We include the axion in our theory by extending (2.15) to the following effective Lagrangian

$$\begin{aligned} \mathcal{L}_{\hat{a}} = & \nu^2 \text{Tr}\{\partial_\mu \Sigma \partial^\mu \Sigma^\dagger\} + \nu_{PQ}^2 \partial_\mu N \partial^\mu N^\dagger + 4\mu\nu^2 \text{Tr}\{B \Sigma^\dagger \partial_0 \Sigma\} + m_\pi^2 \nu^2 \text{Tr}\{M \Sigma + M^\dagger \Sigma^\dagger\} \\ & + 2\mu^2 \nu^2 \left[ \text{Tr}\{\Sigma B^T \Sigma^\dagger B\} + \text{Tr}\{BB\} \right] - a\nu^2 \left( \theta - \frac{i}{4} \text{Tr}\{\log \Sigma - \log \Sigma^\dagger\} - \frac{i}{4} a_{PQ} (\log N - \log N^\dagger) \right)^2. \end{aligned} \quad (2.17)$$

The details on how to construct the extended effective theory can be found in [76].

### 3 Determining the vacuum

In this section, we focus on the vacuum structure of theory (2.15) in the presence of the  $\theta$ -angle and finite baryon charge. In particular, we will first study the general conditions determining the classical vacuum solution as a function of  $\theta$  and  $\mu$ . Armed with these results, we then carefully analyze the properties of the vacuum in the concrete cases  $N_f = 2, 3, 6, 7, 8$  and  $N_f$  arbitrary. To begin with, we notice that for vanishing  $\theta$  the vacuum is determined by the competition of mass and baryon chemical potential. In other words, the ground state can be written as [83]

$$\Sigma_c = \begin{pmatrix} 0 & \mathbf{1}_{N_f} \\ -\mathbf{1}_{N_f} & 0 \end{pmatrix} \cos \varphi + i \begin{pmatrix} \mathcal{I} & 0 \\ 0 & \mathcal{I} \end{pmatrix} \sin \varphi \quad \text{where} \quad \mathcal{I} = \begin{pmatrix} 0 & -\mathbf{1}_{N_f/2} \\ \mathbf{1}_{N_f/2} & 0 \end{pmatrix}, \quad (3.1)$$

where the angle  $\varphi$  is determined by the equations of motion.

To study the effect of  $\theta$  on the vacuum solutions it is convenient to introduce the Witten variables  $\alpha_i$  as [84]

$$\Sigma_0 = U(\alpha_i) \Sigma_c, \quad U(\alpha_i) \equiv \text{diag}\{e^{-i\alpha_1}, \dots, e^{-i\alpha_{N_f}}, e^{-i\alpha_1}, \dots, e^{-i\alpha_{N_f}}\}. \quad (3.2)$$

These variables reflect the possibility to generalize the mass matrix in (2.6) by replacing the  $\mathbf{1}_{N_f}$  with  $\text{diag}\{e^{-i\alpha_1}, \dots, e^{-i\alpha_{N_f}}\}$ . Each phase is the overall axial transformation for each left-right quark pair. The reason why we cannot consider a single (odd number of) Weyl fermion(s) at the time (as is the case for real gauge representations) is that one

cannot write a mass term in this case and the theory overall suffers from a topological anomaly [85].

We take the above to be the general ansatz for the vacuum. The Lagrangian evaluated on this ansatz reads

$$\mathcal{L}_\theta[\Sigma_0] = \nu^2 \left[ 4m_\pi^2 X \cos \varphi + 2\mu^2 N_f \sin^2 \varphi - a\bar{\theta}^2 \right] \quad (3.3)$$

where for later convenience we introduced

$$\bar{\theta} = \theta - \sum_i^{N_f} \alpha_i, \quad X = \sum_i^{N_f} \cos \alpha_i \quad (3.4)$$

where  $\bar{\theta}$  is the effective theta angle that enters physical observables. The equations of motion read

$$\sin \varphi \left( N_f \cos \varphi - \frac{m_\pi^2}{\mu^2} X \right) = 0 \quad (3.5)$$

$$2m_\pi^2 \sin \alpha_i \cos \varphi = a\bar{\theta}, \quad i = 1, \dots, N_f \quad (3.6)$$

and the energy of the system is

$$E = -\nu^2 \left[ 4m_\pi^2 X - a\bar{\theta}^2 \right], \quad \text{normal phase } (\varphi = 0) \quad (3.7)$$

$$E = -\nu^2 \left[ 2 \frac{N_f^2 \mu^4 + m_\pi^4 X^2}{N_f \mu^2} - a\bar{\theta}^2 \right], \quad \text{superfluid phase } \left( \cos \varphi = \frac{m_\pi^2}{N_f \mu^2} X \right). \quad (3.8)$$

Note that when  $a \gg m_\pi$  all the  $\theta$ -dependence is contained in an effective pion mass  $m_\pi^2(\theta) \equiv \frac{m_\pi^2 X}{N_f}$ . For  $\theta = 0$  (i.e.  $U(\alpha_i) = \mathbb{1}$ ) one has  $X = N_f$  and  $\bar{\theta} = 0$ , and we recover the results in [83]. In this case, a normal to superfluid phase transition occurs when  $\mu = m_\pi$ , i.e. when  $\cos \varphi = 1$ . For  $\theta \neq 0$  the  $\theta$ -dependence of the energy may be different in the two phases. Therefore, to find the conditions for the onset of the superfluid phase we first need to determine the  $\theta$  vacuum in both phases. To this end, we observe that in the normal (superfluid) phase the energy is minimized when  $X$  ( $X^2$ ) is maximized. In the former case, the Witten variables are related to  $\theta$  by the well-known equation

$$2m_\pi^2 \sin \alpha_i = a\bar{\theta} = a \left( \theta - \sum_i^{N_f} \alpha_i \right). \quad (3.9)$$

For the general solution we must have for any  $\bar{\theta}$  fixed  $\sin \alpha_i = \sin \alpha_j$ . To solve for the  $\alpha_i$  we consider the expansion in the parameter  $\frac{m_\pi^2}{a}$  that we take to be small. Concretely, at the leading order one needs to solve for  $\bar{\theta} = 0$  and the angles  $\alpha_i$  satisfy

$$\alpha_i = \begin{cases} \pi - \alpha, & i = 1, \dots, n \\ \alpha, & i = n + 1, \dots, N_f \end{cases} \quad (3.10)$$

where  $\alpha$  is the solution of the following modular equation

$$n(\pi - \alpha) + (N_f - n)\alpha = \theta \text{ Mod } 2\pi. \quad (3.11)$$

The modulo comes from the fact that if a solution  $\{\alpha_i\}$  of eq. (3.9) is found, then it is possible to build another solution as follows

$$\alpha_1(\theta + 2\pi) = \alpha_1(\theta) + 2\pi, \quad \alpha_i(\theta + 2\pi) = \alpha_i(\theta), \quad i = 2, \dots, N_f. \quad (3.12)$$

However, since the physics depends only on  $e^{-i\alpha_i}$ , the dynamics is invariant under  $\theta \rightarrow \theta + 2\pi$ . The solution of eq. (3.11) can be written as

$$\alpha = \frac{\theta + (2k - n)\pi}{(N_f - 2n)}, \quad k = 0, \dots, N_f - 2n - 1, \quad n = 0, \dots, \left\lfloor \frac{N_f - 1}{2} \right\rfloor. \quad (3.13)$$

The range for  $k$  above emerges because for  $k \geq N_f - 2n$  we repeat the solution for a given  $n$ . Note that when  $n \neq 0$ , the vacuum spontaneously breaks  $\text{Sp}(2N_f)$  because of the different phases for each quark flavour.

It is well-known that  $CP$  is preserved when  $\bar{\theta} = 0$ . For equal mass quarks as considered here, this happens when  $m_\pi = 0$  or  $\theta = 0$ . On the other hand, for  $\theta = \pi$  the Lagrangian (2.15) possess  $CP$  symmetry but in the normal phase the latter is spontaneously broken by the vacuum [76, 86–88],<sup>1</sup> leading to a strong  $\theta$ -dependence near  $\theta = \pi$ . In fact, assuming that the ground state does not break  $\text{Sp}(2N_f)$  spontaneously (i.e.  $n = 0$ ), the vacua lie at [87]

$$U(\alpha_i) = e^{i\frac{\theta+2\pi k}{N_f}} \mathbb{1}_{2N_f}. \quad (3.14)$$

For  $\theta = \pi$  one has  $X = \cos\left(\frac{(2k+1)\pi}{N_f}\right)$ , which is maximized when  $k = 0$  and  $k = N_f - 1$ , that is

$$U(\alpha_i) = e^{i\frac{\pi}{N_f}} \mathbb{1}_{2N_f}, \quad U(\alpha_i) = e^{-i\frac{\pi}{N_f}} \mathbb{1}_{2N_f}. \quad (3.15)$$

The two solutions are related by a  $CP$  transformation  $U \rightarrow U^\dagger$  and thus  $CP$  is spontaneously broken. For  $N_f > 2$  the minima are separated by an energy barrier while for  $N_f = 2$  the leading order quark-mass induced potential vanishes,<sup>2</sup> apparently leading to a paradoxical situation according to which one has massless pions and no explicit breaking of chiral symmetry. The paradox is simply resolved by going to higher orders in the mass for the chiral Lagrangian. Once these corrections are taken into account they lift, as expected, the vacuum degeneracy yielding two minima separated by a barrier [89, 90]. The spontaneous breaking of  $CP$  at  $\theta = \pi$  is known as Dashen’s phenomenon [86] and has been thoroughly studied in the literature [84, 87–92]. As we shall see, in the superfluid phase  $CP$  may be violated or not at  $\theta = \pi$  depending on the value of  $N_f$ . Once determined the  $\alpha_i$  to the leading order in  $\frac{a}{m_\pi^2}$ , subleading corrections can be more easily computed. Note that in the superfluid phase the equation of motion (3.6) becomes

$$\frac{2m_\pi^4}{N_f \mu^2} X \sin \alpha_i = a\bar{\theta}, \quad i = 1, \dots, N_f. \quad (3.16)$$

Hence, in this case the natural expansion parameter is  $\frac{m_\pi^4}{a\mu^2}$  leading to an enhanced suppression compared to the normal phase.

<sup>1</sup>See also the discussion about the impact on the  $\theta = \pi$  solution due to quark masses ordering provided in the appendix of [76].

<sup>2</sup>As well explained in the work by Smilga [89] this phenomenon occurs because the trace of  $\text{SU}(2)$  matrices are real.

We now consider specific values of  $N_f$  to cover most of the phase diagram for which chiral symmetry is expected to break spontaneously. In the companion paper [1] we will re-examine the values of  $N_f$  that are expected to cover the near-conformal dynamics.

### 3.1 $N_f = 2$

We now focus on the case  $N_f = 2$  which has been previously considered in [71] at the leading order in  $\frac{m_\pi^2}{a}$ . In this limit, the two angles  $\{\alpha_1, \alpha_2\}$  satisfy  $\alpha_1 + \alpha_2 = \theta + 2k\pi$ . Additionally, because  $\sin \alpha_1 = \sin \alpha_2$  we have

$$\sin \alpha_1 = \sin (\theta + 2k\pi - \alpha_1) \quad (3.17)$$

yielding the solutions


$$\sin \alpha_1 = \sin \alpha_2 = \sin \left( \frac{\theta}{2} + k\pi \right). \quad (3.18)$$

Only  $k = 0$  and  $k = 1$  are independent solutions of the equations of motion and agree with the general result in equation (3.13) for the  $\alpha_i$  for  $N_f = 2$  that has the two solutions corresponding to  $n = 0$  and  $k = 0, 1$  (i.e.  $\{\alpha_1, \alpha_2\} = \left\{ \frac{\theta}{2}, \frac{\theta}{2} \right\}$  and  $\{\alpha_1, \alpha_2\} = \left\{ \frac{\theta+2\pi}{2}, \frac{\theta+2\pi}{2} \right\}$ ). To determine the solution corresponding to the ground state for any value of  $\theta$  we need to consider the ground state energy. In the normal phase, the energy is linear in  $X$  and therefore the original equation of motion solutions cross at  $\theta = \pi$  where Dashen's phenomenon occurs stemming from spontaneous  $CP$  symmetry breaking. Because in the superfluid phase we have that the ground state energy is proportional to  $X^2$  the two solutions are identical yielding a degenerate ground state. Interestingly this degeneracy is not lifted by higher order corrections in  $\frac{m_\pi^2}{a}$ . The  $\theta$ -dependence in the two phases is shown in figure 1.

Additionally, when  $\theta = \pi$  the effective mass  $m_\pi^2(\theta) \sim m_\pi^2 \left| \cos \left( \frac{\theta}{2} \right) \right|$  vanishes up to correction of order  $\mathcal{O} \left( \frac{m_\pi^2}{a} \right)$ . Therefore, the mass term disappears from the Lagrangian and the global flavor symmetry is again  $SU(4)$  consequently leading to massless Goldstones when it spontaneously breaks to  $Sp(4)$  [71]. Nevertheless, there is no chiral symmetry restoration in the fundamental Lagrangian. As mentioned, this apparent paradox is solved by realising that  $SU(4)$  is still broken by higher order mass terms in the effective Lagrangian also for  $a \rightarrow \infty$  [71, 87, 89]. By including the first two subleading corrections in  $\frac{m_\pi^2}{a}$ , the energy in the two phases reads

$$E(\theta) = -8m_\pi^2 \nu^2 \left( \left| \cos \frac{\theta}{2} \right| + \frac{1}{2} \frac{m_\pi^2}{a} \sin^2 \frac{\theta}{2} - \frac{1}{4} \frac{m_\pi^4}{a^2} \left| \sin \frac{\theta}{2} \sin \theta \right| \right), \quad \text{normal phase} \quad (3.19)$$

$$E(\theta) = -\nu^2 \left( \frac{4 \left( m_\pi^4 \cos^2 \frac{\theta}{2} + \mu^4 \right)}{\mu^2} + \frac{m_\pi^8 \sin^2 \theta}{a\mu^4} - \frac{m_\pi^{12} \sin^2 \theta \cos \theta}{a^2 \mu^6} \right), \quad \text{superfluid phase.} \quad (3.20)$$



**Figure 1.**  $\theta$ -dependence of the energy for  $N_f = 2$ .

At fixed quark masses, the superfluid phase transition occurs at a critical value of the chemical potential given by

$$\mu_c = m_\pi(\theta) = m_\pi \left[ \sqrt{\left| \cos \frac{\theta}{2} \right|} + \mathcal{O} \left( \frac{m_\pi^2}{a} \right) \right], \quad (3.21)$$

implying that it can be realized for tiny values of the chemical potential when  $\theta \sim \pi$ . To estimate  $\mu_c$  in the region  $\theta \sim \pi$ , we introduce  $\phi \equiv \theta - \pi$  and take into account the leading correction in  $\frac{m_\pi^2}{a}$ . As a result, we have that for small  $|\phi|$  the critical chemical potential in the region  $\theta \sim \pi$  reads

$$\mu_c \sim m_\pi \sqrt{\frac{m_\pi^2}{a} + \frac{|\phi|}{2}}, \quad (3.22)$$

and vanishes for  $a \rightarrow \infty$  and  $\phi \rightarrow 0$ .


Still, in the superfluid phase the energy is an analytic function of  $\theta$ . This can be better appreciated by analysing the  $CP$  order parameter  $\langle F\tilde{F} \rangle \propto -\frac{\partial E}{\partial \theta}$  shown in figure 2 for the two phases. The normal phase is characterised by a discontinuous  $CP$  order parameter at  $\theta = \pi$  while it is clear from the right hand panel of the figure that the superfluid phase displays a smooth behaviour for any value of the  $\theta$ -angle and vanishes for  $\theta = \pi$ . The discontinuity of the  $CP$  order parameter in the normal phase translates into a divergent topological susceptibility  $\chi = \frac{\partial^2 E}{\partial \theta^2}$  at  $\theta = \pi$ . Note that the presence of the baryon chemical potential suppresses  $CP$  violation for every value of  $\theta$  as can be seen from the right panel of figure 2. Finally, we study the effective  $\bar{\theta}$  angle for  $\theta = \pi$  at higher orders in  $\frac{m_\pi^2}{a}$  for the solutions of the equations of motion. We find

$$\bar{\theta} = \frac{2m_\pi^2}{a} \sin \frac{\theta}{2} \Big|_{\theta=\pi} = \frac{2m_\pi^2}{a} + \mathcal{O} \left( \frac{m_\pi^6}{a^3} \right), \quad \text{normal phase} \quad (3.23)$$

$$\bar{\theta} = \frac{m_\pi^4}{a\mu^2} \sin \theta \Big|_{\theta=\pi} = 0, \quad \text{superfluid phase.} \quad (3.24)$$

Note that eq. (3.24) is exact to all orders in  $\frac{m_\pi^4}{a\mu^2}$  since the equation of motion (3.16) becomes

$$\frac{m_\pi^4}{a\mu^2} \sin(2\alpha) = \pi - 2\alpha, \quad (3.25)$$



(a)  $CP$  order parameter for  $N_f = 2$  in the normal phase.

(b)  $CP$  order parameter for  $N_f = 2$  in the superfluid phase.

**Figure 2.**  $CP$  order parameter as a function of  $\theta$  both in the normal and superfluid phase for  $N_f = 2$ .

and has the solution  $\alpha = \pi/2$  (equivalent to  $\alpha = 3\pi/2$  under a shift  $\theta \rightarrow \theta + 2\pi$ ). Therefore to the present order in the chiral expansion, there is no spontaneous  $CP$  symmetry breaking in the superfluid phase for  $\theta = \pi$ .

### 3.2 $N_f = 3$


Compared to the previous subsection, the  $N_f = 3$  case leads to a richer vacuum structure as has been first pointed out in [89], who studied the physics in the normal phase. In fact, we find the following four solutions for the set  $\{\alpha_i\} = \{\alpha_1, \alpha_2, \alpha_3\}$

$$\text{i. } \left\{ \frac{\theta}{3}, \frac{\theta}{3}, \frac{\theta}{3} \right\}, \quad \text{ii. } \left\{ \frac{\theta + 2\pi}{3}, \frac{\theta + 2\pi}{3}, \frac{\theta + 2\pi}{3} \right\}, \quad \text{iii. } \left\{ \frac{\theta + 4\pi}{3}, \frac{\theta + 4\pi}{3}, \frac{\theta + 4\pi}{3} \right\}, \quad \text{iv. } \{ \theta - \pi, \theta - \pi, 2\pi - \theta \}.$$

(3.26)

In figure 3(a) we show the value of the variable  $-X$  as a function of  $\theta$ . Recalling that in the normal phase, the energy is minimized when  $X$  is maximized, we observe that the physical vacuum is ruled by the solutions **i.** and **iii.** which cross at  $\theta = \pi$  where Dashen's phenomenon occurs [89]. In the superfluid phase the story changes since now the minimum of the energy is achieved when  $X^2$  is maximized. The situation is depicted in figure 3(b): here the relevant solutions are **i.**, **ii.**, **iii.**. Solutions **i.** and **ii.** cross at  $\theta = \frac{\pi}{2}$  whereas **ii.** and **iii.** cross at  $\theta = \frac{3\pi}{2}$ . Therefore, at  $\theta = \pi$  we have a unique minimum and  $CP$  is conserved. In other words, Dashen's phenomenon at  $\theta = \pi$  is again absent in the superfluid region. However, two new non-analytic points occur, one at  $\theta = \frac{\pi}{2}$  and the other at  $\theta = \frac{3\pi}{2}$ . These are indications of two novel first-order phase transitions because the derivative of the free energy (corresponding to the  $CP$  order parameter  $\langle F\tilde{F} \rangle$ ) jumps at these two values of  $\theta$ .

Having determined the  $\theta$ -vacuum in the two regions, we proceed by studying the transition between the normal and superfluid phases. Since the  $U(\alpha_i)$  that minimizes the energy depends on  $\theta$  and differs in the two phases, one should determine the critical  $\mu$  by comparing the energies (3.7) and (3.8) in the given intervals of  $\theta$ . We find that when the superfluid solution exists (i.e.  $\cos \varphi \leq 1$ ) it is always realized. In turn, the critical chemical



**Figure 3.**  $\theta$ -dependence of the energy for  $N_f = 3$ .

potential is  $\mu_c = m_\pi(\theta)$ , which is displayed in figure 4. Differently from the  $N_f = 2$  case,  $\mu_c$  exhibits a mild dependence on  $\theta$  and oscillates near the value  $\mu_c = m_\pi$ .

The  $CP$  order parameter is plotted for the normal and superfluid phases in figure 5(a) and figure 5(b), respectively. Finally, we show the value of  $\bar{\theta}$  at  $\theta = \pi$  for the ground state energy. The expression in the  $\frac{m_\pi^2}{a}$  expansion is

$$\bar{\theta} = \frac{\sqrt{3}m_\pi^2}{a} - \frac{m_\pi^4}{\sqrt{3}a^2} - \frac{m_\pi^6}{6\sqrt{3}a^3} + \mathcal{O}\left(\frac{m_\pi^8}{a^4}\right), \quad \text{normal phase} \quad (3.27)$$


$$\bar{\theta} = 0, \quad \text{superfluid phase.} \quad (3.28)$$

### 3.3 $N_f = 6$


For the case of  $N_f = 6$  we, at first, assume that chiral symmetry breaking occurs away from a potential nearby (as function of flavors) conformal phase is discussed at length in [1]. The full set of solutions is given by

$$\begin{aligned} \text{Solutions i-vi : } \alpha_1 = \alpha_2, \dots = \alpha_6 &= \frac{\theta + 2\pi k}{6}, & k = 0, \dots, 5 \\ \text{Solutions vii-ix : } \alpha_1 = \alpha_2 = \dots = \alpha_5 &= \frac{\theta - \pi + 2\pi k}{4}, \quad \alpha_6 = \pi - \alpha_1, & k = 0, \dots, 3 \\ \text{Solutions x-xii : } \alpha_1 = \alpha_2 = \dots = \alpha_4 &= \frac{\theta - 2\pi + 2\pi k}{2}, \quad \alpha_5 = \alpha_6 = \pi - \alpha_1, & k = 0, 1. \end{aligned} \quad (3.29)$$

Here the relevant solutions for  $U(\alpha_i)$  have  $\alpha_1 = \alpha_2 = \dots = \alpha_6 = \alpha$ . Moreover, all the solutions appear in pairs of degenerate energy in the superfluid phase as already observed in the  $N_f = 2$  case. This can be understood by noting that when  $N_f$  is even, then from a given solution  $\alpha$  of eq. (3.13) it is possible to build another solution as  $\alpha + \pi$  that will lead to the same  $-X^2$ . As displayed in figure 6(a), in the normal phase the minimum of the energy is achieved for  $\alpha = \frac{\theta}{6}$  (defined as solution i) and  $\alpha = \frac{\theta + 10\pi}{6}$  (solution iii) with the two solutions crossing at  $\theta = \pi$ . This occurs also in the superfluid phase but now solution i and iii are degenerate for every value of  $\theta$  with  $\alpha = \frac{\theta}{6} + \pi$  and  $\alpha = \frac{\theta + 4\pi}{6}$ , respectively. In figure 6(b) we show the phase diagram of the theory as a function of  $\theta$  and  $\mu$ . Differently




**Figure 4.** Normalized critical chemical potential ( $m_\pi = \nu = 1$ ) above which the superfluid phase is realized as a function of  $\theta$  for  $N_f = 3$ .



(a) The  $CP$  order parameter  $\langle F\tilde{F} \rangle$  as a function of  $\theta$  in the normal phase. We consider  $m_\pi = \nu = 1$ .

(b) The  $CP$  order parameter  $\langle F\tilde{F} \rangle$  as a function of  $\theta$  in the superfluid phase for different values of the chemical potential:  $\mu = 5, 10, 15$ . We consider  $m_\pi = \nu = 1$ .


**Figure 5.**  $\theta$ -dependence of the  $CP$  order parameter for  $N_f = 3$ .



(a)  $\theta$ -dependence of the energy for  $N_f = 6$  both in the normal and superfluid phase.

(b) Normalized critical chemical potential ( $m_\pi = \nu = 1$ ) above which the superfluid phase is realized as a function of  $\theta$ .

**Figure 6.**  $\theta$ -dependence of the energy and of the critical chemical potential for  $N_f = 6$ .



(a) The  $CP$  order parameter  $\langle F\tilde{F} \rangle$  as a function of  $\theta$  in the normal phase. We consider  $m_\pi = \nu = 1$ .

(b) The  $CP$  order parameter  $\langle F\tilde{F} \rangle$  as a function of  $\theta$  in the superfluid phase for different values of the chemical potential:  $\mu = 5, 10, 15$ . We consider  $m_\pi = \nu = 1$ .

**Figure 7.**  $\theta$ -dependence of the  $CP$  order parameter for  $N_f = 6$ .

from  $N_f = 2$  we still observe a Dashen phenomenon at  $\theta = \pi$  and therefore  $CP$  breaks spontaneously for this value.

In the  $N_f = 6$  case we therefore have spontaneous breaking of  $CP$  at  $\theta = \pi$  in both phases as can be seen considering the  $CP$  order parameter represented in figure 7(a) and figure 7(b).

Finally, we report the value of  $\bar{\theta}$  at  $\theta = \pi$  expanded in  $\frac{m_\pi^2}{a}$

$$\bar{\theta} = \frac{m_\pi^2}{a} - \frac{m_\pi^4}{2\sqrt{3}a^2} + \frac{5m_\pi^6}{72a^3} + \mathcal{O}\left(\frac{m_\pi^8}{a^4}\right), \quad \text{normal phase} \quad (3.30)$$


$$\bar{\theta} = -\frac{\sqrt{3}m_\pi^4}{2a\mu^2} + \frac{m_\pi^8}{4\sqrt{3}a^2\mu^4} + \frac{m_\pi^{12}}{48\sqrt{3}a^3\mu^6} + \mathcal{O}\left(\frac{m_\pi^{16}}{a^4\mu^8}\right), \quad \text{superfluid phase.} \quad (3.31)$$

### 3.4 $N_f = 7$

For the  $N_f = 7$  case we have thus 16 solutions of eq. (3.13) which are given by

$$\begin{aligned} \text{Solutions i-vii: } \alpha_1 = \alpha_2, \dots = \alpha_7 &= \frac{\theta + 2\pi k}{7}, & k = 0, \dots, 6 \\ \text{Solutions viii-xii: } \alpha_1 = \alpha_2 = \dots = \alpha_5 &= \frac{\theta - \pi + 2\pi k}{5}, \quad \alpha_6 = \alpha_7 = \pi - \alpha_1, & k = 0, \dots, 4 \\ \text{Solutions xiii-xv: } \alpha_1 = \alpha_2 = \alpha_3 &= \frac{\theta - 2\pi + 2\pi k}{3}, \quad \alpha_4 = \alpha_5 = \alpha_6 = \alpha_7 = \pi - \alpha_1, & k = 0, \dots, 2 \\ \text{Solution xvi: } \alpha_1 = \theta - 3\pi, & \alpha_2 = \alpha_3 = \dots = \alpha_7 = \pi - \alpha_1. \end{aligned} \quad (3.32)$$

The solution that minimizes the energy and their corresponding  $\theta$ -dependence in the two phases is shown in figure 8. As in the previous cases, the solutions that minimize the energy have all equal angles, corresponding here to the first set of solutions in (3.32). In particular, in the normal phase the minimum of the energy is achieved for  $\alpha = \frac{\theta}{7}$  and  $\alpha = \frac{\theta + 12\pi}{7}$  with the two solutions crossing at  $\theta = \pi$  (in figure 8(a)). The  $\theta$ -dependence in the superfluid phase is analogous to the  $N_f = 3$  case. In fact, the relevant solutions are



(a)  $\theta$ -dependence of the energy for  $N_f = 6$  both in the normal and superfluid phase.

(b) Normalized critical chemical potential ( $m_\pi = \nu = 1$ ) above which the superfluid phase is realized as a function of  $\theta$ .

**Figure 8.**  $\theta$ -dependence of the energy and of the critical chemical potential for  $N_f = 7$ .

$\alpha = \frac{\theta}{7}$ ,  $\alpha = \frac{\theta+6\pi}{7}$ , and  $\alpha = \frac{\theta+12\pi}{7}$  with the first two crossing at  $\theta = \frac{\pi}{2}$  while the second and the third at  $\theta = \frac{3\pi}{2}$ . As already described in the  $N_f = 3$  section, the energy is characterised by a unique minimum and  $CP$  intact symmetry at  $\theta = \pi$ .

The critical chemical potential is displayed in figure 8(b). It oscillates with period  $2\pi$  between the values  $\mu_c = m_\pi$  and  $\mu_c = m_\pi(\theta)$ . Figure 8 leads us to further investigate the physical meaning of the  $\theta = \frac{\pi}{2}$  and  $\theta = \frac{3\pi}{2}$  points. As a consequence, we analysed the  $CP$  order parameter which we show in figure 9 as a function of  $\theta$ . As can be seen from the figure, the aforementioned points are actually points of discontinuity of the  $CP$  order parameter therefore it signals the occurrence of a first-order phase transition.

Finally, we provide the values of  $\bar{\theta}$  at  $\theta = \pi$  in the  $\frac{m_\pi^2}{a}$  expansion


$$\bar{\theta} = \frac{2m_\pi^2 \sin \frac{\pi}{7}}{a} - \frac{2m_\pi^4 \cos \frac{3\pi}{14}}{7a^2} + \frac{2m_\pi^6 \sin \frac{\pi}{7} (1 + 3 \sin \frac{3\pi}{14})}{49a^3} + \mathcal{O}\left(\frac{m_\pi^8}{a^4}\right), \quad \text{normal phase} \quad (3.33)$$

$$\bar{\theta} = 0, \quad \text{superfluid phase.} \quad (3.34)$$

### 3.5 $N_f = 8$

We proceed with the general expression for the solutions of (3.13) for the  $N_f = 8$  case


$$\begin{aligned} \text{Solutions i-viii : } \alpha_1 = \dots = \alpha_8 &= \frac{\theta + 2\pi k}{8}, & k = 0, \dots, 7 \\ \text{Solutions ix-xiv : } \alpha_1 = \dots = \alpha_6 &= \frac{\theta - \pi + 2\pi k}{6}, & \alpha_7 = \alpha_8 = \pi - \alpha_1, & k = 0, \dots, 5 \\ \text{Solutions xv-xviii : } \alpha_1 = \dots = \alpha_4 &= \frac{\theta - 2\pi + 2\pi k}{4}, & \alpha_5 = \dots = \alpha_8 = \pi - \alpha_1, & k = 0, \dots, 3 \\ \text{Solutions xix-xx : } \alpha_1 = \alpha_2 &= \frac{\theta - 3\pi + 2\pi k}{2}, & \alpha_3 = \dots = \alpha_8 = \pi - \alpha_1, & k = 0, 1. \end{aligned} \quad (3.35)$$



(a) The  $CP$  order parameter  $\langle F\tilde{F} \rangle$  as a function of  $\theta$  in the normal phase. We consider  $m_\pi = \nu = 1$ .

(b) The  $CP$  order parameter  $\langle F\tilde{F} \rangle$  as a function of  $\theta$  in the superfluid phase for different values of the chemical potential:  $\mu = 5, 10, 15$ . We consider  $m_\pi = \nu = 1$ .

**Figure 9.**  $\theta$ -dependence of the  $CP$  order parameter for  $N_f = 7$ .



(a)  $\theta$ -dependence of the energy for  $N_f = 8$  both in the normal and superfluid phase.

(b) Normalized critical chemical potential ( $m_\pi = \nu = 1$ ) above which the superfluid phase is realized as a function of  $\theta$ .


**Figure 10.**  $\theta$ -dependence of the energy and of the critical chemical potential for  $N_f = 8$ .

In line with our previous analyses, the  $\theta$ -dependence of the energy is minimised by the solutions with all equal angles. Figure 10 displays the behaviour of this minimum in the range  $\theta \in [0, 2\pi]$  as well as the phase diagram of the theory. The  $\theta$ -dependence is analogous to  $N_f = 6$  case as it can be further deduced by studying the  $CP$  order parameter with its behaviour shown in figure 11.

The value of  $\bar{\theta}$  at  $\theta = \pi$  in an  $\frac{m_\pi^2}{a}$  expansion reads

$$\bar{\theta} = \frac{2m_\pi^2 \sin \frac{\pi}{8}}{a} - \frac{m_\pi^4}{4\sqrt{2}a^2} + \frac{\left(1 + \frac{3}{\sqrt{2}}\right)m_\pi^6 \sin \frac{\pi}{8}}{32a^3} + \mathcal{O}\left(\frac{m_\pi^8}{a^4}\right), \quad \text{normal phase} \quad (3.36)$$

$$\bar{\theta} = \frac{m_\pi^4}{\sqrt{2}a\mu^2} - \frac{m_\pi^8}{8a^2\mu^4} + \frac{m_\pi^{12}}{64\sqrt{2}a^3\mu^6} + \mathcal{O}\left(\frac{m_\pi^{16}}{a^4\mu^8}\right), \quad \text{superfluid phase.} \quad (3.37)$$



(a) The  $CP$  order parameter  $\langle F\tilde{F} \rangle$  as a function of  $\theta$  in the normal phase. We consider  $m_\pi = \nu = 1$ .

(b) The  $CP$  order parameter  $\langle F\tilde{F} \rangle$  as a function of  $\theta$  in the superfluid phase for different values of the chemical potential:  $\mu = 5, 10, 15$ . We consider  $m_\pi = \nu = 1$ .

**Figure 11.**  $\theta$ -dependence of the  $CP$  order parameter for  $N_f = 8$ .

### 3.6 Considerations for general $N_f$

Having analyzed in detail the physics for different values of  $N_f$  away from the conformal window, we take a step back and consider the emerging structure hinting at structural differences between the even/odd  $N_f$  cases in the superfluid phase. We start by noting that solutions of the EOM are generally not periodic of  $2\pi$  for  $\theta$ . In fact, the periodicity condition can be satisfied only if at least two solutions cross in the interval  $\theta \in [0, 2\pi]$ . Taking  $U = e^{-i\alpha} \mathbb{1}_{2N_f}$ , one can ask when two different solutions of the equation of motion can have the same energy. This corresponds to requiring

$$\cos\left(\frac{\theta + 2\pi k_1}{N_f}\right) = \cos\left(\frac{\theta + 2\pi k_2}{N_f}\right), \quad \text{normal phase} \quad (3.38)$$

$$\cos^2\left(\frac{\theta + 2\pi k_1}{N_f}\right) = \cos^2\left(\frac{\theta + 2\pi k_2}{N_f}\right), \quad \text{superfluid phase.} \quad (3.39)$$

Both conditions are satisfied when  $k_1 = -\frac{\theta}{\pi} - k_2 + N_f$ . Since  $k_1, k_2$  are integers, it follows that the energy can only be non-trivially equal for  $\theta = \pi$ . Therefore to find the two different solutions, in the normal phase, covering the full  $[0, 2\pi]$  interval for  $\theta$  it is sufficient to consider the case  $k_1 = 0$  that for  $[0, \pi]$  interval corresponds to the ground state energy, furthermore at  $\theta = \pi$  it forces the second solution to be  $k_2 = N_f - 1$ . This result is in agreement with the findings for the specific cases above for the normal phase.

In the superfluid phase, we have two additional solutions to the second condition (3.39):

- When  $N_f$  is even we have the solution  $k_1 = k_2 + \frac{N_f}{2}$  which does not depend on  $\theta$ . This corresponds to the observation that the solutions for even  $N_f$  organize themselves in pairs ( $\alpha$  and  $\alpha + \pi$ ) with the same energy for every  $\theta$ . Additionally, of course, we still have the normal phase solution  $k_1 = 0$  and  $k_2 = N_f - 1$  together with their superfluid degenerate partners. These two classes of solutions cross at  $\theta = \pi$  except for the special  $N_f = 2$  case for which the crossing disappears. Note that the pairs of completely degenerate solutions remain such to all orders in  $\frac{m_\pi^2}{a}$ . In fact, given the

EOM for a certain  $\alpha$

$$\frac{m_\pi^4}{a\mu^2} \sin(2\alpha) = \theta - N_f \alpha, \quad (3.40)$$

we have the same EOM for  $\alpha + \pi$ , upon shifting the  $\theta$ -angle as  $\theta \rightarrow \theta + N_f \pi$ . Being  $N_f$  even, this corresponds to a shift by an integer multiple of  $2\pi$ .

- When  $N_f$  is odd we have the solution  $k_1 = -k_2 + \frac{N_f}{2} - \frac{\theta}{\pi}$  which can be realized for  $\theta = \frac{\pi}{2}$  and  $\theta = \frac{3\pi}{2}$ . Since the solution that minimizes the energy near  $\theta = 0$  is  $\alpha = \frac{\theta}{N_f}$ , then it will cross at  $\theta = \frac{\pi}{2}$  with the solution for  $\theta > \frac{\pi}{2}$  that becomes  $\alpha = \frac{\theta + 2\pi k_2}{N_f} = \frac{\theta - \pi}{N_f} + \pi$ . The latter solution represents the minimum till  $\theta = \frac{3\pi}{2}$  where its energy crosses with  $\alpha = \frac{\theta - 2\pi}{N_f}$ . As a consequence, the vacuum is always non degenerate at  $\theta = \pi$  where  $CP$  is conserved. In particular, for  $\theta = \pi$ ,  $\bar{\theta}$  vanishes to all orders in  $\frac{m_\pi^4}{a\mu^2}$  since the equation of motion (3.16) admits the minimum energy solution  $\alpha = \theta = \pi$ . On the other hand, we have two novel first-order phase transitions at  $\theta = \frac{\pi}{2}, \frac{3\pi}{2}$  due to a jump of the physical vacuum between the two minima and characterized by a discontinuous  $CP$  order parameter.

Finally, we estimate the tension of the domain wall between the two degenerate vacua at  $\theta = \pi$  for even  $N_f$  in the superfluid phase. To this end, we consider the Lagrangian (2.15) and look for solutions  $\alpha_i = \alpha_i(x)$  that interpolate between the two degenerate ground states (3.15) at  $\theta = \pi$  with  $x$  the coordinate orthogonal to the domain wall. By considering the ansatz (3.2), we obtain the tension of the wall as

$$T = 2\nu^2 \int_{-\infty}^{\infty} dx \left[ \sum_{i=1}^{N_f} \alpha_i'(x)^2 - \frac{m_\pi^4}{\mu^2 N_f} \left( \sum_{i=1}^{N_f} \cos \left( \frac{\pi}{N_f} + \alpha_i(x) \right) \right)^2 \right] \quad (3.41)$$

where we have shifted  $\alpha_i$  as  $\alpha_i \rightarrow \alpha_i + \frac{\pi}{N_f}$  such that the boundary conditions are  $\alpha(-\infty) = 0$  and  $\alpha(\infty) = \pi$ . As discussed in [87], the boundary condition fixes  $\alpha_i = \alpha$  for  $i = 1, \dots, N_f - 1$  and  $\alpha_{N_f} = -(N_f - 1)\alpha$ , leading to

$$T = 2\nu^2 \int_{-\infty}^{\infty} dx \left[ (N_f - 1) N_f \alpha'(x)^2 - \frac{m_\pi^4}{\mu^2 N_f} \left( (N_f - 1) \cos \left( \alpha(x) + \frac{\pi}{N_f} \right) + \cos \left( \frac{\pi}{N_f} - (N_f - 1) \alpha(x) \right) \right)^2 \right]. \quad (3.42)$$

By introducing dimensionless coordinates as  $\bar{x} = x \frac{m_\pi^2}{\mu}$ , we obtain that regardless of the exact form of the wall's profile, its tension scales as

$$T \approx \frac{\nu^2 m_\pi^2}{\mu} \quad (3.43)$$

and shows a suppression by a factor of  $\frac{m_\pi}{\mu}$  compared to the normal phase result  $T \approx \nu^2 m_\pi$  [87].

### 3.7 Ground state energy with the axion

The effective action for two-color QCD at finite baryon number in the presence of the Peccei-Quinn axion field  $N$  is

$$L_{\text{eff}} = -a\nu^2 \left( \theta - \sum_{i=1}^{N_f} \alpha_i - \delta \right)^2 - \Lambda^4 + 4\nu^2 m_\pi^2 X \cos \varphi + 2\mu^2 \nu^2 N_f \sin^2 \varphi, \quad \langle N \rangle = e^{-i\delta/a_{PQ}}, \quad (3.44)$$

whose equation of motion  $\theta - \sum_{i=1}^{N_f} \alpha_i - \delta = 0$  is solved for  $\delta = \theta$  and  $\alpha_i = 0$ . This solution minimizes the energy and leads to  $X = N_f$ . Hence, the superfluid ground state energy in presence of the axion corresponds to the known result at  $\theta = 0$  [83]

$$E = -\frac{2\nu^2 N_f}{\mu^2} (m_\pi^4 + \mu^4) . \tag{3.45}$$

## 4 Symmetry-breaking pattern and spectrum

In the section above we concentrated on determining the vacuum structure of the theory. We now move to establish the associated symmetry-breaking pattern starting with the theory without an axion. The pattern will allow us to disentangle the light degrees of freedom of the theory. We will keep the analysis as general as possible so that it is applicable also for the continuation of our work related to near-conformal field theories in the large-charge expansion [1].

### 4.1 Symmetry-breaking pattern without the axion

The dynamics of the theory at hand depends on three parameters: the mass of the quarks  $m$ , the chemical potential  $\mu$ , and the scale of the axial anomaly  $a$ . The latter corresponds to the energy at which the pseudo-Goldstone (the  $S$ -particle) emerging from the spontaneous symmetry breaking of the axial symmetry acquires an anomalous mass. We are interested in the low energy theory where  $S$  is kept into the spectrum. We, therefore, consider the following hierarchy<sup>3</sup>

$$m \ll \sqrt{a} \leq \mu \ll 4\pi\nu , \tag{4.1}$$

where the last inequality implies the validity of the chiral EFT. For  $m = \mu = 0$  and in absence of the singlet  $S$ , the infrared spectrum consists of massless Goldstone bosons of the spontaneously broken chiral symmetry as summarized below.

$$\begin{array}{ll}
 m = 0 & \\
 \mu = 0 & \\
 \sqrt{a} \gg \nu & 
 \end{array}
 \qquad
 \begin{array}{l}
 2N_f^2 - N_f - 1 \\
 \text{Goldstones transforming under the} \\
 \text{antisymm. representation of } \text{Sp}(2N_f) \\
 \nearrow \\
 \text{SU}(2N_f) \xrightarrow{2N_f^2 - N_f - 1} \text{Sp}(2N_f)
 \end{array}$$

When setting to zero the anomaly, and in the absence of the quark mass, the  $S$ -field becomes the Goldstone boson of the  $U(1)_A$ .

We now provide the full spectrum of light particles in the following limits:

1.  $\mathbf{m} \neq \mathbf{0}, \mu = 0$  and  $\sqrt{a} \ll 4\pi\nu$ .

Working in steps, one first considers the would-be pattern of spontaneous symmetry breaking according to which  $\text{SU}(2N_f) \times U(1)_A$  breaks spontaneously to  $\text{Sp}(2N_f)$ . Then the associated  $2N_f^2 - N_f$  Goldstone bosons acquire masses via the non-zero quark mass and the anomaly.

---


<sup>3</sup>This hierarchy of scales is typical when modelling neutron stars in 2-color QCD at finite isospin chemical potential [2].

2.  $\mathbf{m} = \mathbf{0}, \mu \neq \mathbf{0}$  and  $\sqrt{a} \ll 4\pi\nu$ .

The introduction of the chemical potential breaks explicitly the final  $\text{Sp}(2N_f)$  to  $\text{SU}(N_f)_L \times \text{SU}(N_f)_R \times \text{U}(1)_B$  while the anomaly breaks  $\text{U}(1)_A$ . The relativistic Bose-Einstein condensation leads to the further spontaneous symmetry breaking  $\text{SU}(N_f)_L \times \text{SU}(N_f)_R \times \text{U}(1)_B \rightsquigarrow \text{Sp}(N_f)_L \times \text{Sp}(N_f)_R$ . Therefore, the final spectrum is composed of  $N_f^2 - N_f - 1$  massless Goldstone bosons the massive  $S$  state due to the anomaly, and  $N_f^2$  modes with mass proportional to the chemical potential. The latter belong to the  $(N_f, N_f)$  irreducible representation of  $\text{Sp}(N_f)_L \times \text{Sp}(N_f)_R$  whereas the true Goldstones arrange themselves into three different irreducible representations: a singlet  $(1, 1)$ , the  $(\frac{N_f(N_f-1)}{2} - 1, 1)$  representation and the  $(1, \frac{N_f(N_f-1)}{2} - 1)$ . The sum of the degrees of freedom adds to  $2N_f^2 - N_f$ .

3.  $\mathbf{m} \neq \mathbf{0}, \mu \neq \mathbf{0}$  and  $\sqrt{a} \ll 4\pi\nu$ .

The situation here is involved due to the competition between the quark mass term and the baryon chemical potential. The chemical potential breaks  $\text{Sp}(2N_f)$  down to  $\text{SU}(N_f)_L \times \text{SU}(N_f)_R \times \text{U}(1)_B$ , which, in turn, is explicitly broken by the mass term to its vectorial subgroup  $\text{SU}(N_f)_V \times \text{U}(1)_B$ . Finally, the superfluid vacuum breaks the latter symmetry group to  $\text{Sp}(N_f)_V$  yielding to  $\frac{N_f^2 - N_f}{2}$  massless Goldstone bosons as summarized below



The case of our interest is the last one, where we have  $\frac{N_f^2 - N_f}{2}$  Goldstones transforming according to the antisymmetric representation of  $\text{Sp}(N_f)_V$  plus a singlet. Additionally, there's the  $\eta'$  like state  $S$  with mass of order  $\sqrt{a}$ .

The naive counting above of the number of Goldstone bosons holds correct here since there are no Goldstone bosons of type II [93] in this case. The reason is that the dimension of the representation of the Goldstone bosons is identical to the number of broken generators (see discussion in [94]).

### 4.2 Symmetry-breaking pattern with the axion

In this section, we analyse the symmetry-breaking pattern for  $m \neq 0, \mu \neq 0, a \ll 4\pi\nu$  but including the presence of the additional  $\text{U}(1)_{PQ}$  symmetry discussed at the end of section 2. The latter is classically exact but quantum mechanically anomalous and spontaneously broken, leading to the existence of a new pseudo-Goldstone boson (the axion) with the mass

of the order of the scale of the anomaly. We summarize below the pattern of symmetry breaking

$$\begin{array}{c}
 \text{SU}(2N_f) \times \text{U}(1)_A \times \text{U}(1)_{PQ} \xrightarrow{2N_f^2 - N_f + 1} \text{Sp}(2N_f) \\
 \begin{array}{c}
 \swarrow m, a, a_{PQ} \\
 \downarrow \mu \\
 \text{SU}(N_f)_V \times \text{U}(1)_B \\
 \text{superfluid phase} \\
 \downarrow \frac{N_f^2 - N_f}{2} \\
 \text{Sp}(N_f)_V
 \end{array}
 \end{array}$$

From the picture is clear that the spectrum of Goldstones remains the same as in the case without the axion with an additional singlet massive state.

In the next section, we will study the dispersion relations of the light modes describing the infrared dynamics by explicitly expanding the Lagrangian at the quadratic order in the fluctuations around the ground state.

### 4.3 Fluctuations spectrum

To analytically determine the dispersion relations of the different relevant states around the vacuum we consider the large  $a/m_\pi^2$  expansion and stop at the leading order. We recover eqs. (69) and (84) of [83] for  $\theta = 0$ , and generalize them to include the  $\theta$ -angle by taking it into account via an effective quark mass matrix. We obtain

$$\omega_1^2 = k^2 + \mu^2, \quad \square \square \quad \frac{1}{2} N_f (N_f + 1) \quad (4.2)$$

$$\omega_2^2 = k^2 + \frac{m_\pi^4 X^2}{\mu^2 N_f^2}, \quad \square \quad \frac{1}{2} N_f (N_f - 1) - 1 \quad (4.3)$$

$$\omega_3^2 = k^2 + \frac{2(\mu^4 N_f^2 + 3m_\pi^4 X^2)}{N_f^2 \mu^2} + A, \quad \bullet + \square \quad \frac{1}{2} N_f (N_f - 1) \quad (4.4)$$

$$\omega_4^2 = k^2 + \frac{2(\mu^4 N_f^2 + 3m_\pi^4 X^2)}{N_f^2 \mu^2} - A, \quad \bullet + \square \quad \frac{1}{2} N_f (N_f - 1) \quad (4.5)$$

$$\omega_5^2 = k^2 + M_S^2, \quad \bullet \quad 1 \quad (4.6)$$

where

$$A = \frac{2}{N_f^2 \mu^2} \sqrt{(N_f^2 \mu^4 + 3m_\pi^4 X^2)^2 + 4N_f^2 \mu^2 m_\pi^4 k^2 X^2}, \quad (4.7)$$

$$M_S^2 = \frac{a\mu^4 N_f^3 + 2\mu^2 m_\pi^4 X^2}{2\mu^4 N_f^2 - 2m_\pi^4 X^2} \left( 1 - \frac{m_\pi^4 X^2}{\mu^2 N_f^2} \right). \quad (4.8)$$

The Young tableaux describe the irreducible representations of  $\text{Sp}(N_f)$  according to which the states transform with  $\bullet$  denoting the singlet representation. The number of degrees of freedom sum to  $\dim \left( \frac{\text{U}(2N_f)}{\text{Sp}(2N_f)} \right) = N_f(2N_f - 1)$ , i.e. the number of pseudo-Goldstones

of the chiral symmetry breaking plus the  $\eta'$  like  $S$ -particle. For  $\theta = 0$  the first four dispersion relations reduce to the ones found in [83].  $\omega_4$  describes true Goldstone modes with speed  $v_G = 1$  which parametrize the coset  $\frac{SU(N_f)}{Sp(N_f)}$  and correspond to the  $\pi$  modes considered in [94]. These are the modes in which we will be mainly interested in the companion paper of this work since they dominate the large-charge dynamics. In addition, we have modes with a mass of order  $\mu$  ( $\omega_1$  and  $\omega_3$ ) and modes with mass  $\frac{m_\pi^2 X}{\mu N_f}$ . If we include the axion in our theory, the dispersion relations of the modes described by eqs. (4.2)–(4.5) remain the same with  $X = N_f$ , i.e. we have the spectrum at  $\theta = 0$  as expected from the Peccei-Quinn mechanism. On top of that, we have the modes stemming from the mixing between the singlet  $S$  and the axion, which is described by the following inverse propagator

$$D^{-1} = \begin{pmatrix} \frac{\omega^2 - k^2}{\sin^2 \varphi} - M_S^2 & -\frac{a\sqrt{N_f}a_{PQ}}{4\sqrt{2}\nu_{PQ}\sin^2 \varphi} \\ -\frac{a\sqrt{N_f}a_{PQ}}{4\sqrt{2}\nu_{PQ}\sin^2 \varphi} & \frac{\omega^2 - k^2}{4\nu^2 \sin^2 \varphi} - M_{\hat{a}}^2 \end{pmatrix}, \quad (4.9)$$

where

$$M_S^2 = \frac{(a\mu^4 N_f + 2\mu^2 m_\pi^4)}{2\mu^4 - 2m_\pi^4} \quad (4.10)$$

$$M_{\hat{a}}^2 = \frac{a\mu^4 a_{PQ}^2}{16\nu_{PQ}^2 (\mu^4 - m_\pi^4)}. \quad (4.11)$$

The dispersion relations read

$$\omega_{6,7} = \frac{1}{2} \sqrt{4k^2 + 8\nu^2 M_{\hat{a}}^2 \sin^2 \varphi + 2M_S^2 \sin^2 \varphi \pm \frac{1}{\nu_{PQ}} \sqrt{2a^2 \nu^2 N_f a_{PQ}^2 + 4\nu_{PQ}^2 \sin^4 \varphi (M_S^2 - 4\nu^2 M_{\hat{a}}^2)^2}}. \quad (4.12)$$

As can be seen from the inverse propagator, the  $S$  particle and the axion decouple in the anomaly-free limits  $a \rightarrow 0$  and  $a_{PQ} \rightarrow 0$ . In the former case, the  $S$ -particle describes a pseudo-Goldstone mode with mass of order  $M_S^2 = \frac{\mu^2 m_\pi^4}{\mu^4 - m_\pi^4}$ , while for  $a_{PQ} \rightarrow 0$  we have  $M_S^2 = \frac{aN_f}{2}$  for  $\mu \gg m_\pi$ , in agreement with the Witten-Veneziano relation [73, 74]. On the other hand, in the limit of vanishing anomalies, the axion is massless being the  $U(1)_{PQ}$  symmetry exact.

## 5 Conclusions

We investigated the spectrum, the pattern of chiral symmetry breaking and the dispersion relations at low energy of two-color QCD as a function of the baryon chemical potential, the topological term responsible for strong  $CP$  breaking as well as the quark masses. The analysis is applicable to  $Sp(2N)$  gauge groups with fermions in the fundamental representation. We explicitly considered the dynamics stemming from two, three, six, seven and eight light matter flavors and determined the normal and superfluid ground state. We showed that the vacuum acquires a rich structure stemming from the presence of the  $CP$  violating topological operator and unveiled novel phases. We analysed these phases, studied the dependence of the critical chemical potential on the  $\theta$  angle, delineating the boundary between the normal phase and the superfluid phase. By investigating the behaviour

of the  $CP$  order parameter we characterised the order of the phase transitions which are shown to be first order. The results are readily applicable, in the normal phase, to the  $\theta$  and axion physics of composite Goldstone Higgs models. In particular, a new composite strong source of  $CP$  violation can be relevant to investigating composite electroweak baryogenesis [24, 95, 96] while the nonzero chemical potential analysis is useful for studying asymmetric dark matter [24, 28, 29] and the interplay with  $CP$  violation. Our predictions can guide and be tested by first principle lattice simulations at nonzero baryon chemical potential but including also the effects of the topological susceptibility [97, 98].

We stress that the results are valid for a number of flavours below the lower boundary of the conformal window whose value increases with the underlying number of colors of the  $Sp(2N_c)$  gauge theory investigated here. In the large  $N_c$  limit the eta-prime becomes a quasi-Goldstone boson and therefore validates the effective approach used here. The analysis is further performed respecting the hierarchies of scales of (4.1) allowing to consider the impact of the chemical potential as a modification of the underlying pattern of chiral symmetry breaking in the vacuum. In the future, higher derivative terms can be consistently included in the framework such as the Wess-Zumino-Witten term and its possible gauging [99] following [100].

## Acknowledgments

M.T. was supported by Agencia Nacional de Investigación y Desarrollo (ANID) grant 72210390.

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP<sup>3</sup> supports the goals of the International Year of Basic Sciences for Sustainable Development.

## References

- [1] J. Bersini, A. D’Alise, F. Sannino and M. Torres, *The  $\theta$ -angle and axion physics of two-color QCD at fixed baryon charge. Part II. Approaching the conformal window*, [arXiv:2208.09227](https://arxiv.org/abs/2208.09227) [[INSPIRE](https://inspirehep.net/literature/2208092)].
- [2] S. Gandolfi, J. Lippuner, A.W. Steiner, I. Tews, X. Du and M. Al-Mamun, *From the microscopic to the macroscopic world: from nucleons to neutron stars*, *J. Phys. G* **46** (2019) 103001 [[arXiv:1903.06730](https://arxiv.org/abs/1903.06730)] [[INSPIRE](https://inspirehep.net/literature/1903067)].
- [3] T. Boeckel, S. Schettler and J. Schaffner-Bielich, *The Cosmological QCD Phase Transition Revisited*, *Prog. Part. Nucl. Phys.* **66** (2011) 266 [[arXiv:1012.3342](https://arxiv.org/abs/1012.3342)] [[INSPIRE](https://inspirehep.net/literature/1012334)].
- [4] F. Sannino, *Conformal Dynamics for TeV Physics and Cosmology*, *Acta Phys. Pol. B* **40** (2009) 3533 [[arXiv:0911.0931](https://arxiv.org/abs/0911.0931)] [[INSPIRE](https://inspirehep.net/literature/0911093)].
- [5] G. Cacciapaglia, C. Pica and F. Sannino, *Fundamental Composite Dynamics: A Review*, *Phys. Rept.* **877** (2020) 1 [[arXiv:2002.04914](https://arxiv.org/abs/2002.04914)] [[INSPIRE](https://inspirehep.net/literature/2002049)].
- [6] C.A. Baker et al., *An Improved experimental limit on the electric dipole moment of the neutron*, *Phys. Rev. Lett.* **97** (2006) 131801 [[hep-ex/0602020](https://arxiv.org/abs/hep-ex/0602020)] [[INSPIRE](https://inspirehep.net/literature/131801)].

- [7] C. Abel et al., *Measurement of the Permanent Electric Dipole Moment of the Neutron*, *Phys. Rev. Lett.* **124** (2020) 081803 [[arXiv:2001.11966](#)] [[INSPIRE](#)].
- [8] R.D. Peccei and H.R. Quinn, *CP Conservation in the Presence of Instantons*, *Phys. Rev. Lett.* **38** (1977) 1440 [[INSPIRE](#)].
- [9] R.D. Peccei and H.R. Quinn, *Constraints Imposed by CP Conservation in the Presence of Instantons*, *Phys. Rev. D* **16** (1977) 1791 [[INSPIRE](#)].
- [10] S. Weinberg, *A New Light Boson?*, *Phys. Rev. Lett.* **40** (1978) 223 [[INSPIRE](#)].
- [11] F. Wilczek, *Problem of Strong P and T Invariance in the Presence of Instantons*, *Phys. Rev. Lett.* **40** (1978) 279 [[INSPIRE](#)].
- [12] J.E. Kim, *Weak Interaction Singlet and Strong CP Invariance*, *Phys. Rev. Lett.* **43** (1979) 103 [[INSPIRE](#)].
- [13] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Can Confinement Ensure Natural CP Invariance of Strong Interactions?*, *Nucl. Phys. B* **166** (1980) 493 [[INSPIRE](#)].
- [14] M. Dine, W. Fischler and M. Srednicki, *A Simple Solution to the Strong CP Problem with a Harmless Axion*, *Phys. Lett. B* **104** (1981) 199 [[INSPIRE](#)].
- [15] R.N. Mohapatra and G. Senjanović, *Natural Suppression of Strong p and t Noninvariance*, *Phys. Lett. B* **79** (1978) 283 [[INSPIRE](#)].
- [16] A.E. Nelson, *Naturally Weak CP-violation*, *Phys. Lett. B* **136** (1984) 387 [[INSPIRE](#)].
- [17] S.M. Barr, *A Natural Class of Nonpeccei-quinn Models*, *Phys. Rev. D* **30** (1984) 1805 [[INSPIRE](#)].
- [18] A.E. Nelson, *Calculation of  $\theta$  Barr*, *Phys. Lett. B* **143** (1984) 165 [[INSPIRE](#)].
- [19] S.D.H. Hsu and F. Sannino, *New solutions to the strong CP problem*, *Phys. Lett. B* **605** (2005) 369 [[hep-ph/0408319](#)] [[INSPIRE](#)].
- [20] F. Wilczek and G. Moore, *Superheavy Light Quarks and the Strong P, T Problem*, [arXiv:1601.02937](#) [[INSPIRE](#)].
- [21] R. Lewis, C. Pica and F. Sannino, *Light Asymmetric Dark Matter on the Lattice: SU(2) Technicolor with Two Fundamental Flavors*, *Phys. Rev. D* **85** (2012) 014504 [[arXiv:1109.3513](#)] [[INSPIRE](#)].
- [22] A. Hietanen, R. Lewis, C. Pica and F. Sannino, *Fundamental Composite Higgs Dynamics on the Lattice: SU(2) with Two Flavors*, *JHEP* **07** (2014) 116 [[arXiv:1404.2794](#)] [[INSPIRE](#)].
- [23] T. Appelquist, P.S. Rodrigues da Silva and F. Sannino, *Enhanced global symmetries and the chiral phase transition*, *Phys. Rev. D* **60** (1999) 116007 [[hep-ph/9906555](#)] [[INSPIRE](#)].
- [24] T.A. Ryttov and F. Sannino, *Ultra Minimal Technicolor and its Dark Matter TIMP*, *Phys. Rev. D* **78** (2008) 115010 [[arXiv:0809.0713](#)] [[INSPIRE](#)].
- [25] D.B. Kaplan and H. Georgi, *SU(2)  $\times$  U(1) Breaking by Vacuum Misalignment*, *Phys. Lett. B* **136** (1984) 183 [[INSPIRE](#)].
- [26] E.D. Carlson, M.E. Machacek and L.J. Hall, *Self-interacting dark matter*, *Astrophys. J.* **398** (1992) 43 [[INSPIRE](#)].
- [27] Y. Hochberg, E. Kuflik, T. Volansky and J.G. Wacker, *Mechanism for Thermal Relic Dark Matter of Strongly Interacting Massive Particles*, *Phys. Rev. Lett.* **113** (2014) 171301 [[arXiv:1402.5143](#)] [[INSPIRE](#)].

- [28] Y. Hochberg, E. Kuflik, H. Murayama, T. Volansky and J.G. Wacker, *Model for Thermal Relic Dark Matter of Strongly Interacting Massive Particles*, *Phys. Rev. Lett.* **115** (2015) 021301 [[arXiv:1411.3727](#)] [[INSPIRE](#)].
- [29] M. Hansen, K. Langæble and F. Sannino, *SIMP model at NNLO in chiral perturbation theory*, *Phys. Rev. D* **92** (2015) 075036 [[arXiv:1507.01590](#)] [[INSPIRE](#)].
- [30] F. Sannino and K. Tuominen, *Orientifold theory dynamics and symmetry breaking*, *Phys. Rev. D* **71** (2005) 051901 [[hep-ph/0405209](#)] [[INSPIRE](#)].
- [31] D.D. Dietrich, F. Sannino and K. Tuominen, *Light composite Higgs from higher representations versus electroweak precision measurements: Predictions for CERN LHC*, *Phys. Rev. D* **72** (2005) 055001 [[hep-ph/0505059](#)] [[INSPIRE](#)].
- [32] D.D. Dietrich and F. Sannino, *Conformal window of SU(N) gauge theories with fermions in higher dimensional representations*, *Phys. Rev. D* **75** (2007) 085018 [[hep-ph/0611341](#)] [[INSPIRE](#)].
- [33] F. Sannino, *Conformal Windows of Sp(2N) and SO(N) Gauge Theories*, *Phys. Rev. D* **79** (2009) 096007 [[arXiv:0902.3494](#)] [[INSPIRE](#)].
- [34] T. Karavirta, J. Rantaharju, K. Rummukainen and K. Tuominen, *Determining the conformal window: SU(2) gauge theory with  $N_f = 4, 6$  and 10 fermion flavours*, *JHEP* **05** (2012) 003 [[arXiv:1111.4104](#)] [[INSPIRE](#)].
- [35] H. Ohki et al., *Study of the scaling properties in SU(2) gauge theory with eight flavors*, *PoS LATTICE2010* (2010) 066 [[arXiv:1011.0373](#)] [[INSPIRE](#)].
- [36] J. Rantaharju, T. Karavirta, V. Leino, T. Rantalaiho, K. Rummukainen and K. Tuominen, *The gradient flow running coupling in SU(2) with 8 flavors*, *PoS LATTICE2014* (2014) 258 [[arXiv:1411.4879](#)] [[INSPIRE](#)].
- [37] A. Hietanen, R. Lewis, C. Pica and F. Sannino, *Composite Goldstone Dark Matter: Experimental Predictions from the Lattice*, *JHEP* **12** (2014) 130 [[arXiv:1308.4130](#)] [[INSPIRE](#)].
- [38] R. Arthur, V. Drach, M. Hansen, A. Hietanen, C. Pica and F. Sannino, *SU(2) gauge theory with two fundamental flavors: A minimal template for model building*, *Phys. Rev. D* **94** (2016) 094507 [[arXiv:1602.06559](#)] [[INSPIRE](#)].
- [39] V. Leino, K. Rummukainen, J.M. Suorsa, K. Tuominen and S. Tähtinen, *Infrared Behaviour of SU(2) Gauge Theory with  $N_f$  fundamental flavours*, *PoS Confinement2018* (2019) 225 [[arXiv:1811.12438](#)] [[INSPIRE](#)].
- [40] F. Bursa, L. Del Debbio, L. Keegan, C. Pica and T. Pickup, *Mass anomalous dimension in SU(2) with six fundamental fermions*, *Phys. Lett. B* **696** (2011) 374 [[arXiv:1007.3067](#)] [[INSPIRE](#)].
- [41] M. Hayakawa, K.I. Ishikawa, S. Takeda and N. Yamada, *Running coupling constant and mass anomalous dimension of six-flavor SU(2) gauge theory*, *Phys. Rev. D* **88** (2013) 094504 [[arXiv:1307.6997](#)] [[INSPIRE](#)].
- [42] T. Appelquist et al., *Two-Color Gauge Theory with Novel Infrared Behavior*, *Phys. Rev. Lett.* **112** (2014) 111601 [[arXiv:1311.4889](#)] [[INSPIRE](#)].
- [43] V. Leino, T. Rantalaiho, K. Rummukainen, J.M. Suorsa, K. Tuominen and S. Tähtinen, *Gradient flow running coupling in SU(2)  $N_f = 6$  flavors*, *PoS LATTICE2016* (2016) 218 [[arXiv:1610.09989](#)] [[INSPIRE](#)].

- [44] J.M. Suorsa et al., *Mass anomalous dimension of SU(2) using the spectral density method*, *PoS LATTICE2016* (2016) 389 [[arXiv:1611.02022](#)] [[INSPIRE](#)].
- [45] A. Nakamura, *Quarks and Gluons at Finite Temperature and Density*, *Phys. Lett. B* **149** (1984) 391 [[INSPIRE](#)].
- [46] S. Hands, J.B. Kogut, M.-P. Lombardo and S.E. Morrison, *Symmetries and spectrum of SU(2) lattice gauge theory at finite chemical potential*, *Nucl. Phys. B* **558** (1999) 327 [[hep-lat/9902034](#)] [[INSPIRE](#)].
- [47] J.B. Kogut, D. Toublan and D.K. Sinclair, *Diquark condensation at nonzero chemical potential and temperature*, *Phys. Lett. B* **514** (2001) 77 [[hep-lat/0104010](#)] [[INSPIRE](#)].
- [48] J.B. Kogut, D. Toublan and D.K. Sinclair, *The Phase diagram of four flavor SU(2) lattice gauge theory at nonzero chemical potential and temperature*, *Nucl. Phys. B* **642** (2002) 181 [[hep-lat/0205019](#)] [[INSPIRE](#)].
- [49] S. Muroya, A. Nakamura and C. Nonaka, *Behavior of hadrons at finite density: Lattice study of color SU(2) QCD*, *Phys. Lett. B* **551** (2003) 305 [[hep-lat/0211010](#)] [[INSPIRE](#)].
- [50] S. Hands, S. Kim and J.-I. Skullerud, *Deconfinement in dense 2-color QCD*, *Eur. Phys. J. C* **48** (2006) 193 [[hep-lat/0604004](#)] [[INSPIRE](#)].
- [51] S. Cotter, P. Giudice, S. Hands and J.-I. Skullerud, *Towards the phase diagram of dense two-color matter*, *Phys. Rev. D* **87** (2013) 034507 [[arXiv:1210.4496](#)] [[INSPIRE](#)].
- [52] T. Boz, S. Cotter, L. Fister, D. Mehta and J.-I. Skullerud, *Phase transitions and gluodynamics in 2-colour matter at high density*, *Eur. Phys. J. A* **49** (2013) 87 [[arXiv:1303.3223](#)] [[INSPIRE](#)].
- [53] V.V. Braguta, E.M. Ilgenfritz, A.Y. Kotov, A.V. Molochkov and A.A. Nikolaev, *Study of the phase diagram of dense two-color QCD within lattice simulation*, *Phys. Rev. D* **94** (2016) 114510 [[arXiv:1605.04090](#)] [[INSPIRE](#)].
- [54] L. Holicki, J. Wilhelm, D. Smith, B. Wellegehausen and L. von Smekal, *Two-colour QCD at finite density with two flavours of staggered quarks*, *PoS LATTICE2016* (2017) 052 [[arXiv:1701.04664](#)] [[INSPIRE](#)].
- [55] V.G. Bornyakov, V.V. Braguta, E.M. Ilgenfritz, A.Y. Kotov, A.V. Molochkov and A.A. Nikolaev, *Observation of deconfinement in a cold dense quark medium*, *JHEP* **03** (2018) 161 [[arXiv:1711.01869](#)] [[INSPIRE](#)].
- [56] T. Boz, O. Hajizadeh, A. Maas and J.-I. Skullerud, *Finite-density gauge correlation functions in  $QC_2D$* , *Phys. Rev. D* **99** (2019) 074514 [[arXiv:1812.08517](#)] [[INSPIRE](#)].
- [57] N.Y. Astrakhantsev et al., *Lattice study of static quark-antiquark interactions in dense quark matter*, *JHEP* **05** (2019) 171 [[arXiv:1808.06466](#)] [[INSPIRE](#)].
- [58] T. Boz, P. Giudice, S. Hands and J.-I. Skullerud, *Dense two-color QCD towards continuum and chiral limits*, *Phys. Rev. D* **101** (2020) 074506 [[arXiv:1912.10975](#)] [[INSPIRE](#)].
- [59] K. Iida, E. Itou and T.-G. Lee, *Two-colour QCD phases and the topology at low temperature and high density*, *JHEP* **01** (2020) 181 [[arXiv:1910.07872](#)] [[INSPIRE](#)].
- [60] J. Wilhelm, L. Holicki, D. Smith, B. Wellegehausen and L. von Smekal, *Continuum Goldstone spectrum of two-color QCD at finite density with staggered quarks*, *Phys. Rev. D* **100** (2019) 114507 [[arXiv:1910.04495](#)] [[INSPIRE](#)].
- [61] V.G. Bornyakov, V.V. Braguta, A.A. Nikolaev and R.N. Rogalyov, *Effects of Dense Quark Matter on Gluon Propagators in Lattice  $QC_2D$* , *Phys. Rev. D* **102** (2020) 114511 [[arXiv:2003.00232](#)] [[INSPIRE](#)].

- [62] K. Iida, E. Itou and T.-G. Lee, *Relative scale setting for two-color QCD with  $N_f = 2$  Wilson fermions*, *Prog. Theor. Exp. Phys.* **2021** (2021) 013B05 [[arXiv:2008.06322](#)] [[INSPIRE](#)].
- [63] N. Astrakhantsev, V.V. Braguta, E.M. Ilgenfritz, A.Y. Kotov and A.A. Nikolaev, *Lattice study of thermodynamic properties of dense  $QC_2D$* , *Phys. Rev. D* **102** (2020) 074507 [[arXiv:2007.07640](#)] [[INSPIRE](#)].
- [64] T.G. Khunjua, K.G. Klimenko and R.N. Zhokhov, *The dual properties of chiral and isospin asymmetric dense quark matter formed of two-color quarks*, *JHEP* **06** (2020) 148 [[arXiv:2003.10562](#)] [[INSPIRE](#)].
- [65] T. Kojo and D. Suenaga, *Thermal quarks and gluon propagators in two-color dense QCD*, *Phys. Rev. D* **103** (2021) 094008 [[arXiv:2102.07231](#)] [[INSPIRE](#)].
- [66] D.F. Litim and F. Sannino, *Asymptotic safety guaranteed*, *JHEP* **12** (2014) 178 [[arXiv:1406.2337](#)] [[INSPIRE](#)].
- [67] D.F. Litim, M. Mojaza and F. Sannino, *Vacuum stability of asymptotically safe gauge-Yukawa theories*, *JHEP* **01** (2016) 081 [[arXiv:1501.03061](#)] [[INSPIRE](#)].
- [68] O. Antipin and F. Sannino, *Conformal Window 2.0: The large  $N_f$  safe story*, *Phys. Rev. D* **97** (2018) 116007 [[arXiv:1709.02354](#)] [[INSPIRE](#)].
- [69] V. Leino, T. Rindlisbacher, K. Rummukainen, F. Sannino and K. Tuominen, *Safety versus triviality on the lattice*, *Phys. Rev. D* **101** (2020) 074508 [[arXiv:1908.04605](#)] [[INSPIRE](#)].
- [70] J. Rantaharju, T. Rindlisbacher, K. Rummukainen, A. Salami and K. Tuominen, *Spectrum of  $SU(2)$  gauge theory at large number of flavors*, *Phys. Rev. D* **104** (2021) 114504 [[arXiv:2108.10630](#)] [[INSPIRE](#)].
- [71] M.A. Metlitski and A.R. Zhitnitsky, *Theta-parameter in 2 color QCD at finite baryon and isospin density*, *Nucl. Phys. B* **731** (2005) 309 [[hep-ph/0508004](#)] [[INSPIRE](#)].
- [72] A. Amato, V. Leino, K. Rummukainen, K. Tuominen and S. Tähtinen, *From chiral symmetry breaking to conformality in  $SU(2)$  gauge theory*, [arXiv:1806.07154](#) [[INSPIRE](#)].
- [73] E. Witten, *Current Algebra Theorems for the  $U(1)$  Goldstone Boson*, *Nucl. Phys. B* **156** (1979) 269 [[INSPIRE](#)].
- [74] G. Veneziano,  *$U(1)$  Without Instantons*, *Nucl. Phys. B* **159** (1979) 213 [[INSPIRE](#)].
- [75] R.J. Crewther, P. Di Vecchia, G. Veneziano and E. Witten, *Chiral Estimate of the Electric Dipole Moment of the Neutron in Quantum Chromodynamics*, *Phys. Lett. B* **88** (1979) 123 [*Erratum* *ibid.* **91** (1980) 487] [[INSPIRE](#)].
- [76] P. Di Vecchia and F. Sannino, *The Physics of the  $\theta$ -angle for Composite Extensions of the Standard Model*, *Eur. Phys. J. Plus* **129** (2014) 262 [[arXiv:1310.0954](#)] [[INSPIRE](#)].
- [77] K. Kawarabayashi and N. Ohta, *The Problem of  $\eta$  in the Large  $N$  Limit: Effective Lagrangian Approach*, *Nucl. Phys. B* **175** (1980) 477 [[INSPIRE](#)].
- [78] M. Pospelov and A. Ritz, *Theta vacua, QCD sum rules, and the neutron electric dipole moment*, *Nucl. Phys. B* **573** (2000) 177 [[hep-ph/9908508](#)] [[INSPIRE](#)].
- [79] R. Essig et al., *Working Group Report: New Light Weakly Coupled Particles*, in proceedings of the *Community Summer Study 2013: Snowmass on the Mississippi*, Minneapolis, MN, U.S.A., 29 July–6 August 2013, [arXiv:1311.0029](#) [[INSPIRE](#)].
- [80] M.K. Gaillard, M.B. Gavela, R. Houtz, P. Quilez and R. Del Rey, *Color unified dynamical axion*, *Eur. Phys. J. C* **78** (2018) 972 [[arXiv:1805.06465](#)] [[INSPIRE](#)].

- [81] M.B. Gavela, M. Ibe, P. Quilez and T.T. Yanagida, *Automatic Peccei-Quinn symmetry*, *Eur. Phys. J. C* **79** (2019) 542 [[arXiv:1812.08174](#)] [[INSPIRE](#)].
- [82] L. Di Luzio, B. Gavela, P. Quilez and A. Ringwald, *An even lighter QCD axion*, *JHEP* **05** (2021) 184 [[arXiv:2102.00012](#)] [[INSPIRE](#)].
- [83] J.B. Kogut, M.A. Stephanov, D. Toublan, J.J.M. Verbaarschot and A. Zhitnitsky, *QCD-like theories at finite baryon density*, *Nucl. Phys. B* **582** (2000) 477 [[hep-ph/0001171](#)] [[INSPIRE](#)].
- [84] E. Witten, *Large N Chiral Dynamics*, *Annals Phys.* **128** (1980) 363 [[INSPIRE](#)].
- [85] E. Witten, *An SU(2) Anomaly*, *Phys. Lett. B* **117** (1982) 324 [[INSPIRE](#)].
- [86] R.F. Dashen, *Some features of chiral symmetry breaking*, *Phys. Rev. D* **3** (1971) 1879 [[INSPIRE](#)].
- [87] D. Gaiotto, Z. Komargodski and N. Seiberg, *Time-reversal breaking in QCD<sub>4</sub>, walls, and dualities in 2 + 1 dimensions*, *JHEP* **01** (2018) 110 [[arXiv:1708.06806](#)] [[INSPIRE](#)].
- [88] P. Di Vecchia, G. Rossi, G. Veneziano and S. Yankielowicz, *Spontaneous CP breaking in QCD and the axion potential: an effective Lagrangian approach*, *JHEP* **12** (2017) 104 [[arXiv:1709.00731](#)] [[INSPIRE](#)].
- [89] A.V. Smilga, *QCD at  $\theta \sim \pi$* , *Phys. Rev. D* **59** (1999) 114021 [[hep-ph/9805214](#)] [[INSPIRE](#)].
- [90] M.H.G. Tytgat, *QCD at  $\theta \sim \pi$  reexamined: Domain walls and spontaneous CP-violation*, *Phys. Rev. D* **61** (2000) 114009 [[hep-ph/9909532](#)] [[INSPIRE](#)].
- [91] M. Creutz, *Quark masses and chiral symmetry*, *Phys. Rev. D* **52** (1995) 2951 [[hep-th/9505112](#)] [[INSPIRE](#)].
- [92] R. Kitano, R. Matsudo, N. Yamada and M. Yamazaki, *Peeking into the  $\theta$  vacuum*, *Phys. Lett. B* **822** (2021) 136657 [[arXiv:2102.08784](#)] [[INSPIRE](#)].
- [93] H.B. Nielsen and S. Chadha, *On How to Count Goldstone Bosons*, *Nucl. Phys. B* **105** (1976) 445 [[INSPIRE](#)].
- [94] D. Orlando, S. Reffert and F. Sannino, *Charging the Conformal Window*, *Phys. Rev. D* **103** (2021) 105026 [[arXiv:2003.08396](#)] [[INSPIRE](#)].
- [95] J.M. Cline, M. Jarvinen and F. Sannino, *The Electroweak Phase Transition in Nearly Conformal Technicolor*, *Phys. Rev. D* **78** (2008) 075027 [[arXiv:0808.1512](#)] [[INSPIRE](#)].
- [96] S. Bruggisser, B. Von Harling, O. Matsedonskyi and G. Servant, *Electroweak Phase Transition and Baryogenesis in Composite Higgs Models*, *JHEP* **12** (2018) 099 [[arXiv:1804.07314](#)] [[INSPIRE](#)].
- [97] L. Del Debbio, L. Giusti and C. Pica, *Topological susceptibility in the SU(3) gauge theory*, *Phys. Rev. Lett.* **94** (2005) 032003 [[hep-th/0407052](#)] [[INSPIRE](#)].
- [98] B. Alles, M. D'Elia, A. Di Giacomo and C. Pica, *Analysis of systematic errors in the calculation of renormalization constants of the topological susceptibility on the lattice*, *Phys. Rev. D* **74** (2006) 094503 [[hep-lat/0604007](#)] [[INSPIRE](#)].
- [99] Z.-y. Duan, P.S. Rodrigues da Silva and F. Sannino, *Enhanced global symmetry constraints on  $\epsilon$  terms*, *Nucl. Phys. B* **592** (2001) 371 [[hep-ph/0001303](#)] [[INSPIRE](#)].
- [100] J.T. Lenaghan, F. Sannino and K. Splittorff, *The Superfluid and conformal phase transitions of two color QCD*, *Phys. Rev. D* **65** (2002) 054002 [[hep-ph/0107099](#)] [[INSPIRE](#)].