


# Generalized Parton Distributions (GPDs) through DVCS and DVMP

Kornelija Passek-K.

Rudjer Boskovic Institute, Croatia



*ACHT2015*

*"Strong Interactions in Quantum Field Theory"*

*Leibnitz, Oct, 8th, 2015.*

## 1 Introduction

- Resolving nucleon structure (form factors, PDFs, ...)

## 2 DVCS, DVMP, GPDs — theory

- Deeply virtual Compton scattering (DVCS)
- ..., deeply virtual meson electroproduction (DVMP)
- Generalized parton distributions (GPDs)

## 3 DVCS, DVMP, GPDs — phenomenology

- Experimental status
- Towards unravelling GPDs
- Modeling venues
- One example approach... [D. Müller (Uni. Bochum, Uni. Regensburg, IRB), K. Kumerički (Uni. Zagreb), K.P-K. (IRB), T. Lautenschlager (Uni. Regensburg), A. Schäfer (Uni. Regensburg)]

## 4 Summary

## SCATTERING

→ elastic	$(e^- p \rightarrow e^- p)$	}	exclusive
→ inelastic	$(e^- p \rightarrow e^- \gamma p, e^- p \rightarrow e^- \pi p)$		
	$(e^- p \rightarrow e^- X)$	}	inclusive


## SCATTERING

→ elastic	$(e^- p \rightarrow e^- p)$	}	exclusive
→ inelastic	$(e^- p \rightarrow e^- \gamma p, e^- p \rightarrow e^- \pi p)$		
	$(e^- p \rightarrow e^- X)$	}	inclusive

# Parton distribution functions

## ■ Deeply inelastic scattering

$$-q_1^2 \equiv Q^2 \rightarrow \infty, \quad x_{BJ} \equiv \frac{-q_1^2}{2p \cdot q_1} \rightarrow \text{cte.} \quad (\text{Bjorken limit})$$




# Parton distribution functions

■ Deeply inelastic scattering

$$-q_1^2 \equiv Q^2 \rightarrow \infty, \quad x_{BJ} \equiv \frac{-q_1^2}{2p \cdot q_1} \rightarrow \text{cte.}$$




(Bjorken limit)



# Parton distribution functions

- Deeply inelastic scattering

$$-q_1^2 \equiv Q^2 \rightarrow \infty, \quad x_{BJ} \equiv \frac{-q_1^2}{2p \cdot q_1} \rightarrow \text{cte.} \quad (\text{Bjorken limit})$$




- no information on spatial distribution of partons



# Electromagnetic form factors

- Form factors  $\rightarrow$  charge distribution

$$\Gamma^\mu(\gamma^* p \rightarrow p) = \gamma^\mu F_1(Q^2) + \frac{\kappa_p}{2M_p} i \sigma_{\nu}^\mu q_1^\nu F_2(Q^2)$$



$$q(b_\perp) \sim \int dq_1 e^{iq_1 \cdot b_\perp} F_1(t = q_1^2)$$








# Electromagnetic form factors

- Form factors  $\rightarrow$  charge distribution

$$\Gamma^\mu(\gamma^* p \rightarrow p) = \gamma^\mu F_1(Q^2) + \frac{\kappa_p}{2M_p} i \sigma_{\nu}^\mu q_1^\nu F_2(Q^2)$$

$$q(b_\perp) \sim \int dq_1 e^{iq_1 \cdot b_\perp} F_1(t = q_1^2)$$






# Electromagnetic form factors

- Form factors  $\rightarrow$  charge distribution


$$\Gamma^\mu(\gamma^* p \rightarrow p) = \gamma^\mu F_1(Q^2) + \frac{\kappa_p}{2M_p} i \sigma_{\nu}^\mu q_1^\nu F_2(Q^2)$$

$$q(b_\perp) \sim \int dq_1 e^{iq_1 \cdot b_\perp} F_1(t = q_1^2)$$



# DIS and Compton scattering

- Deeply inelastic scattering  $-q_1^2 \equiv Q^2 \rightarrow \infty$ ,  $x_{BJ} \equiv \frac{-q_1^2}{2p \cdot q_1} \rightarrow \text{cte.}$




$$\sigma_{tot}(\gamma^* p \rightarrow X) \stackrel{\text{optical theorem}}{\propto} \text{Im} \mathcal{A}(\gamma^* p \rightarrow \gamma^* p)$$

forward Compton scattering

# Probing the proton with two photons

- Deeply virtual Compton scattering (DVCS) [Müller '92, et al. '94]




$$P = P_1 + P_2$$

$$q = (q_1 + q_2)/2$$

$$\Delta = P_2 - P_1$$

# Probing the proton with two photons

- Deeply virtual Compton scattering (DVCS) [Müller '92, et al. '94]



$$P = P_1 + P_2$$

$$q = (q_1 + q_2)/2$$

$$\Delta = P_2 - P_1$$

generalized Bjorken limit:

$$-q^2 \stackrel{\text{DVCS}}{\simeq} Q^2/2 \rightarrow \infty$$

$$\xi = \frac{-q^2}{2P \cdot q} \rightarrow \text{const (as } x_B)$$

$$\vartheta = \frac{q_1^2 - q_2^2}{q_1^2 + q_2^2} \approx \frac{\eta}{\xi} \stackrel{\text{DVCS}}{=} 1$$

$$t = (P_2 - P_1)^2 = \Delta^2$$


$$\sigma \propto |\mathcal{A}(\gamma^* p \rightarrow \gamma p)|^2$$

# Factorization of DVCS $\longrightarrow$ GPDs

$\rightarrow$  cross-section can be expressed in terms of (the squares of)

Compton form factors:  ${}^a\mathcal{H}(\xi, t, Q^2), {}^a\mathcal{E}(\xi, t, Q^2), \tilde{{}^a\mathcal{H}}(\xi, t, Q^2), \tilde{{}^a\mathcal{E}}(\xi, t, Q^2), \dots$

[Collins and Freund '99]



- Compton form factor is a convolution:

$${}^a\mathcal{H}(\xi, t, Q^2) = \int dx C^a(x, \xi, Q^2/\mu^2) H^a(x, \eta = \xi, t, \mu^2)$$

# Factorization of DVCS $\longrightarrow$ GPDs

$${}^a\mathcal{H}(\xi, t, Q^2) = \int dx C^a(x, \xi, Q^2/\mu^2, \alpha_s(\mu)) H^a(x, \eta = \xi, t, \mu^2)$$

- $C^a(x, \xi, Q^2/\mu^2)$  ... hard scattering amplitude

$\rightarrow$  pQCD

# Factorization of DVCS $\longrightarrow$ GPDs

$${}^a\mathcal{H}(\xi, t, Q^2) = \int dx C^a(x, \xi, Q^2/\mu^2, \alpha_s(\mu)) H^a(x, \eta = \xi, t, \mu^2)$$

- $C^a(x, \xi, Q^2/\mu^2)$  ... hard scattering amplitude

$\rightarrow$  pQCD

- $H^a(x, \eta = \xi, t, \mu^2)$  ... Generalized parton distribution (GPD)


$\rightarrow$  nonperturbative input

$\rightarrow$  evolution  $\Leftarrow$  pQCD (limiting cases DGLAP ( $\eta = 0$ ) and ERBL ( $\eta = 1$ ) evolution equations)

$$\mu^2 \frac{d}{d\mu^2} \mathbf{F}(x, \eta, t, \mu^2) = \int_{-1}^1 \frac{dy}{2\eta} \mathbf{V}\left(\frac{\eta+x}{2\eta}, \frac{\eta+y}{2\eta}; \eta \middle| \alpha_s(\mu)\right) \cdot \mathbf{F}(y, \eta, t, \mu^2)$$




# Complementary processes



(double) DVCS

$$\boxed{\gamma^* p \rightarrow \gamma^* p}$$


$(ep \rightarrow ep l^+ l^-)$



spacelike DVCS

$$\boxed{\gamma^* p \rightarrow \gamma p}$$


$(ep \rightarrow ep \gamma)$



timelike DVCS

$$\boxed{\gamma p \rightarrow \gamma^* p}$$

$(\gamma p \rightarrow pl^+ l^-)$



Deeply virtual production of mesons (DVMP)

more difficult, but access to flavours


$$\boxed{\gamma^* p \rightarrow Mp}$$

factorization: [Collins, Frankfurt, Strikman '97]

# Hard-scattering amplitudes

## DVCS

$$\gamma^* q \rightarrow \gamma q, \gamma^* g \rightarrow \gamma g$$




NLO: [Ji et al, Belitsky et al, Mankiewicz et al, '97]

# Hard-scattering amplitudes

## DVMP

$$\gamma^* q \rightarrow (q\bar{q})q, \quad \gamma^* g \rightarrow (q\bar{q})g$$



NLO: [Belitsky and Müller '01, Ivanov et al '04]

# Definition of GPDs

- In QCD **GPDs** are defined as [Müller '92, et al. '94, Ji, Radyushkin '96]

$$\tilde{F}^q(x, \eta, t = \Delta^2) = \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle P_2 | \bar{q}(-z) \gamma^+ \gamma_5 q(z) | P_1 \rangle \Big|_{z^+=0, \mathbf{z}_\perp=0}$$


$$\tilde{F}^g(x, \eta, t = \Delta^2) = \frac{4}{P^+} \int \frac{dz^-}{2\pi} e^{ixP^+z^-} \langle P_2 | G_a^{+\mu}(-z) \tilde{G}_{a\mu}^+(z) | P_1 \rangle \Big|_{\dots}$$

- Decomposing into helicity conserving and non-conserving part:

$$F^a = \frac{\bar{u}(P_2) \gamma^+ u(P_1)}{P^+} H^a + \frac{\bar{u}(P_2) i\sigma^{+\nu} u(P_1) \Delta_\nu}{2MP^+} E^a \quad a = q, g$$

# Properties of GPDs

- Forward limit ( $\Delta \rightarrow 0, \eta \rightarrow 0$ ):  $\Rightarrow \tilde{H}$ -GPDs  $\rightarrow$  PDFs




$$P = P_1 + P_2$$

$$\Delta = P_2 - P_1, \Delta^2 = t$$

$$\eta = -\frac{\Delta^+}{P^+}$$

# Properties of GPDs

- Forward limit ( $\Delta \rightarrow 0, \eta \rightarrow 0$ ):  $\Rightarrow \tilde{H}$ -GPDs  $\rightarrow$  PDFs




- Sum rules:  $\Rightarrow$  GPD  $\rightarrow$  form factors

$$\sum_{q=u,d} Q_q \int_{-1}^1 dx \begin{cases} H^q(x, \eta, t) \\ E^q(x, \eta, t) \end{cases} = \begin{cases} F_1(t) \\ F_2(t) \end{cases}$$

# Properties of GPDs

- Forward limit ( $\Delta \rightarrow 0, \eta \rightarrow 0$ ):  $\Rightarrow \tilde{H}$ -GPDs  $\rightarrow$  PDFs



- Sum rules:  $\Rightarrow$  GPD  $\rightarrow$  form factors


$$\sum_{q=u,d} Q_q \int_{-1}^1 dx \begin{cases} H^q(x, \eta, t) \\ E^q(x, \eta, t) \end{cases} = \begin{cases} F_1(t) \\ F_2(t) \end{cases}$$

- Possibility of solution of proton spin problem

$$\frac{1}{2} \int_{-1}^1 dx x \left[ H^q(x, \eta, t) + E^q(x, \eta, t) \right] = J^q(t) \quad [\text{Ji '96}]$$

# Properties of GPDs

- Forward limit ( $\Delta \rightarrow 0, \eta \rightarrow 0$ ):  $\Rightarrow \tilde{H}$ -GPDs  $\rightarrow$  PDFs



- Sum rules:  $\Rightarrow$  GPD  $\rightarrow$  form factors

$$\sum_{q=u,d} Q_q \int_{-1}^1 dx \begin{cases} H^q(x, \eta, t) \\ E^q(x, \eta, t) \end{cases} = \begin{cases} F_1(t) \\ F_2(t) \end{cases}$$


- Possibility of solution of proton spin problem

$$\frac{1}{2} \int_{-1}^1 dx x \left[ H^q(x, \eta, t) + E^q(x, \eta, t) \right] = J^q(t) \quad [\text{Ji '96}]$$

- polynomiality (of Mellin moments in  $\eta$ ) and positivity constraints



# Contemporary hierarchy of parton distributions




W.-D. Nowak, Access to GPDs at COMPASS (Primosten/Croatia, Sept. 10-16, 2014)

32

# Experimental status

## DVCS



[from Kumericki et al. 2015]

## DVMP

- in the last decade: vector meson ( $\rho$ ,  $J/\Psi$ ,  $\phi$ ) production at H1 and ZEUS, COMPASS, pseudoscalar mesons ( $\pi$ ,  $\eta$ ) at CLAS ...

→ new results from COMPASS, JLab12 (EIC)

# Towards unravelling GPDs

DVCS: Compton form factors

$${}^a\mathcal{H}(\xi, t, Q^2) = \int dx C^a(x, \xi, Q^2/\mu^2) H^a(x, \xi, t, \mu^2)_{a=q,G} \text{ or NS,S}(\Sigma, G)$$

DVMP: transition form factors

$${}^a\mathcal{T}(\xi, t, Q^2) = \int dx \int dy T^a(x, \xi, y, Q^2/\mu^2) H^a(x, \xi, t, \mu^2) \phi(y, \mu^2)$$

- **Complete deconvolution is impossible** and to extract GPDs from the experiment we need to **model** their functional dependence, or alternatively model form factors for start.
- *"Curse of the dimensionality"*  
When the dimensionality increases, the volume of the space increases so fast that the **available data become sparse**.
- Known **GPD constraints do not help enough** (don't decrease the dimensionality of the GPD domain space).

# Modeling venues

- double distribution amplitude (DDA) satisfy automatically the polynomiality constraint so many models based on it, or specifically Radyushkin's DDA (RDDA) (VGG code, [Goeke et al. 01], BMK model [Belitsky, Muller, Kirchner 01], GK model [Goloskokov, Kroll 05]))
- 'aligned jet' model [Freund, McDermott, Strikman 02], polynomials [Belitsky et al. '98], [Liuti et al. '07], [Moutarde '09]
- 'dual model' [Polyakov, Shuvaev 02], [Guzey, Teckentrup 06], [Polyakov 07]
- **models in Mellin-Barnes integral representation** [K. Kumericki, D. Muller, K. Passek-K 08, ...]
- fitting Compton form factors with neural networks [Kumericki, Muller, Schaefer 11]

- factorization formula for singlet DVCS CFFs:

$${}^S\mathcal{H}(\xi, t, Q^2) = \int dx \mathbf{C}(x, \xi, Q^2/\mu^2, \alpha_s(\mu)) \mathbf{H}(x, \xi, t, \mu^2)$$

- factorization formula for singlet DVCS CFFs:

$${}^S\mathcal{H}(\xi, t, Q^2) = \int dx \mathbf{C}(x, \xi, Q^2/\mu^2, \alpha_s(\mu)) \mathbf{H}(x, \xi, t, \mu^2)$$

- ... in terms of **conformal moments**

(analogous to Mellin moments in DIS:  $x^n \rightarrow C_n^{3/2}(x), C_n^{5/2}(x)$ ):

$$= 2 \sum_{j=0}^{\infty} \xi^{-j-1} \mathbf{C}_j(Q^2/\mu^2, \alpha_s(\mu)) \mathbf{H}_j(\xi = \eta, t, \mu^2)$$

$$H_j^q(\eta, \dots) = \frac{\Gamma(3/2)\Gamma(j+1)}{2^{j+1}\Gamma(j+3/2)} \int_{-1}^1 dx \eta^{j-1} C_j^{3/2}(x/\eta) H^q(x, \eta, \dots)$$

...

$H_j^a$  even polynomials in  $\eta$  with maximal power  $\eta^{j+1}$

- factorization formula for singlet DVCS CFFs:

$${}^S\mathcal{H}(\xi, t, Q^2) = \int dx \mathbf{C}(x, \xi, Q^2/\mu^2, \alpha_s(\mu)) \mathbf{H}(x, \xi, t, \mu^2)$$

- ... in terms of **conformal moments**

(analogous to Mellin moments in DIS:  $x^n \rightarrow C_n^{3/2}(x), C_n^{5/2}(x)$ ):

$$= 2 \sum_{j=0}^{\infty} \xi^{-j-1} \mathbf{C}_j(Q^2/\mu^2, \alpha_s(\mu)) \mathbf{H}_j(\xi = \eta, t, \mu^2)$$

$$H_j^q(\eta, \dots) = \frac{\Gamma(3/2)\Gamma(j+1)}{2^{j+1}\Gamma(j+3/2)} \int_{-1}^1 dx \eta^{j-1} C_j^{3/2}(x/\eta) H^q(x, \eta, \dots)$$

...

$H_j^a$  even polynomials in  $\eta$  with maximal power  $\eta^{j+1}$

- series summed using **Mellin-Barnes** integral over complex  $j$ :

$$= \frac{1}{2i} \int_{c-i\infty}^{c+i\infty} dj \left[ i + \tan\left(\frac{\pi j}{2}\right) \right] \xi^{-j-1} \mathbf{C}_j(Q^2/\mu^2, \alpha_s(\mu)) \mathbf{H}_j(\xi, t, \mu^2)$$



# Advantages of conformal moments and Mellin-Barnes representation

- enables **simpler inclusion of evolution** effects
- powerful analytic methods of **complex  $j$**  plane are available (similar to complex angular momentum of Regge theory)
- opens the door for interesting **modelling of GPDs**
- possible efficient and stable numerical treatment  $\Rightarrow$  stable and fast **computer code** for evolution and fitting
- moments are equal to matrix elements of local operators and are thus **directly accessible on the lattice**
  
- **NNLO corrections for DVCS** accessible by making use of conformal OPE and known NNLO DIS results



# NLO and NNLO corrections

for generic parameters













**Thick lines:**  
 "hard" gluon  
 $N_G = 0.4$   
 $\alpha_G(0) = \alpha_\Sigma(0) + 0.05$


**Thin lines:**  
 "soft" gluon  
 $N_G = 0.3$   
 $\alpha_G(0) = \alpha_\Sigma(0) - 0.02$

$$\delta^P K = \frac{|\mathcal{H}^{N^P \text{LO}}|}{|\mathcal{H}^{N^{P-1} \text{LO}}|} - 1, \quad \delta^P \varphi = \arg\left(\frac{\mathcal{H}^{N^P \text{LO}}}{\mathcal{H}^{N^{P-1} \text{LO}}}\right)$$





# Fits (GeParD output)



GeParD parameters:  
 SPEED = 2 P = 1 NF = 4  
 SCHEME = CSBAR OZ = 4.0  
 Fit/fit parameters:  
 NS = 0.17 NO = 0.58  
 ALOS = 1.1 ALPO = 1.0  
 ALPS = 0.15 ALPO = 0.15  
 MDS = 0.75 MDC = 0.33  
 SEDS = 0.53 SEDG = 5.2  
 d.o.f = 69 - 8 = 61  $\chi^2 = 52.7$



# NNLO fit to HERA DVCS+DIS data




- Fourier transform of GPD for  $\eta = 0$  can be interpreted as probability density depending on  $x$  and transversal distance  $b$   
[Burkardt '00, '02]




$$H(x, \vec{b}) = \int \frac{d^2 \vec{\Delta}}{(2\pi)^2} e^{-i\vec{b} \cdot \vec{\Delta}} H(x, \eta = 0, \Delta^2 = -\vec{\Delta}^2),$$

# Three-dimensional image of a proton

Quarks:



Gluons:



# Summary

- Generalized parton distributions offer a unique window into hadron structure as they combine features of form factors, parton densities and distribution amplitudes.
- They are experimentally accessible via DVCS, DVMP . . . different processes offer different insight and should provide more complete picture.

- Generalized parton distributions offer a unique window into hadron structure as they combine features of form factors, parton densities and distribution amplitudes.
- They are experimentally accessible via DVCS, DVMP ... different processes offer different insight and should provide more complete picture.
- Extraction of GPDs is extremely challenging but efforts for global fits are being made.
- New data are expected from COMPASS and JLab12. DVCS and related processes have a large role in EIC proposal.

The End