

A short overview on low mass scalars at future lepton colliders

Tania Robens^{1,2}

¹Ruder Boskovic Institute, Bijenicka cesta 54, 10000 Zagreb, Croatia

²Theoretical Physics Department, CERN, 1211 Geneva 23, Switzerland

Abstract. I give a short summary on scenarios with new physics scalars that could be investigated at future e^+e^- colliders. I concentrate on cases where at least one of the additional scalar has a mass below 125 GeV, and discuss both models where this could be realized, as well as studies which focus on such scenarios. This work is based on [1], and partial results were also presented in [2].

1 Introduction

In the European Strategy report [3, 4], Higgs factories were identified as one of the high priority projects after the HL-LHC. At such machines, the properties of the Higgs particles should be measurable to utmost precision. Furthermore, new physics scalar states could also be produced in the mass range up to ~ 160 GeV depending on the collider process.

At LEP, a large variety of new physics models have been investigated, with a concise summary given in [5, 6]. However, these searches, as well as more recent bounds from direct searches for light scalars at the LHC, still leave room for new light scalar states in many new physics models. In this short proceeding, I summarize the current status of such searches and present new physics scenarios that still allow for such states, taking all current constraints into account. The work presented here is based on [1], and some of the results have additionally been presented in [2].

2 Processes at Higgs factories

At the center-of-mass (com) energies of Higgs factories, Higgs strahlung is the dominant production mode [7]. Leading-order predictions for Zh production at e^+e^- colliders for low mass scalars which are Standard Model (SM)-like, using Madgraph5 [8], are shown in figure 1 for a center-of-mass energy of 250 GeV. The VBF-type production of $e^+e^- \rightarrow h\nu_\ell\bar{\nu}_\ell$ contains contributions from Zh production, where $Z \rightarrow \nu_\ell\bar{\nu}_\ell$. While for lower masses VBF

production still plays a role, for higher scalar masses the dominant contribution stems from Zh production.




Figure 1. Leading-order production cross sections for Zh and $h\nu_\ell\bar{\nu}_\ell$ production at an e^+e^- collider with a com energy of 250 GeV using Madgraph5 for an SM-like scalar h . Shown is also the contribution of Zh to $\nu_\ell\bar{\nu}_\ell h$ using a factorized approach for the Z decay. Update of plot in [1], first presented in [2].

3 Projections for additional searches

The production of lighter scalars in scalar strahlung has already been investigated in various works [9, 10], where the latter focusses on the investigation of different detector concepts in an ILC environment. In principle two different analysis methods exist, which either use the pure Z recoil ("recoil method") or take the light scalar decay into $b\bar{b}$ into account. In [9], the sensitivity of the ILC for low-mass scalars in Zh was investigated. Figure 2 shows the reach of these two methods at 95% CL limit for agreement with a background only hypothesis, which can directly be translated into an upper bound on rescaling. The authors validate their method by reproducing LEP results [6, 11] for these channels prior to applying their method to the ILC.

Along similar lines, a more recent study that uses only the recoil method and compares different detector options has been presented in [10]. The corresponding results are shown in figure 3. The authors perform their analysis in a model where the coupling of the new resonance is rescaled by a mixing angle $\sin\theta$ and the results can therefore be directly compared to the ones displayed in figure 2.

4 Current enhancements and benchmark motivations

Recently, there has been a lot of activity in the attempt to fit the possible low-mass excesses at LEP [6] and CMS [12] within new physics scenarios. I here mainly comment on the work




Figure 2. Sensitivity predictions for an ILC with a com energy of 250 GeV from [9]. See text for details.




Figure 3. Upper bounds on the mixing angle for the model discussed in [10], in a comparison of different detector concepts and using the recoil method.

presented in [13], where several models are fitted to these excesses that contain singlet and doublet extensions of the SM scalar sector. The authors consider models with an additional doublet as well as a (complex) singlet, labelled N2HDM and 2HDMS, respectively. For both models, the authors investigate the possibility to fit the observed excesses. Out of the parameter space fitting these excesses, they then render rate predictions for a 250 GeV collider with a total luminosity of $\mathcal{L} = 2 \text{ ab}^{-1}$. The corresponding results are shown in figure 4. In particular, it is of high interest that also other final states for the h decay, as e.g. $\tau^+\tau^-$, gg , or W^+W^- can render sizeable rates. Related work concentrating on the N2HDM can be found in [14]. See also [15–23] for other models that can in principle comply with an excess in this mass range.




Figure 4. *Left:* Points in the 2HDMs that agree with both CMS and LEP excess and which can be probed at the ILC. *Right:* predicted rates in the 2HDMs and N2HDM at 250 GeV using full target luminosity.

5 Connection to electroweak phase transition

Another important topic is the connection of models with extended scalar sectors with different scenarios of electroweak phase transitions. In particular, for scenarios where the second scalar is lighter than a SM like candidate, such states can be investigated in Higgs-strahlung and the associated decay of $h \rightarrow h_i h_i$. Due to the clean environment of a lepton collider, these processes can be investigated down to relatively low rates.

A standard reference for such processes is [24], where the authors consider Higgs-strahlung at a 240 GeV e^+e^- collider, with the Higgs subsequently decays into two light scalar states corresponding to the above target signature. They give 95 % confidence level bounds for the branching ratios into the decay products of the two light scalars as a function of the light scalar masses for an integrated luminosity of $\int \mathcal{L} = 5 \text{ ab}^{-1}$ following a detailed study. We show their results for various channels in figure 5.




Figure 5. 95 % confidence bounds on branching ratios for Higgs decay into a pair of lighter particles, for a com energy of 240 GeV and $\int \mathcal{L} = 5 \text{ ab}^{-1}$. Taken from [24].

Depending on the mass, model, and decay mode, branching ratios down to $\sim 10^{-4}$ can be tested.

In simple singlet extensions it is possible to test regions in the models parameter space which can lead to a strong first-order electroweak phase transition. There has been a lot of recent activity in this field recently; here we show results from [25], where in addition several collider sensitivity projections are shown, including the bounds derived in [24]. It becomes clear that e^+e^- Higgs factories would be an ideal environment to confirm or rule out such scenarios.




Figure 6. Expected bounds on Higgs production via Higgs strahlung and subsequent decay into two light scalars, in the singlet extension scenario discussed in [25, 26]. *Left:* Taken from [25]. *Right:* For $\cos \theta = 0$ the constraints mainly stem from $h_{125} \rightarrow$ invisible searches. Depending on m_1 this scenario can be tested at current or future collider experiments.

At the LHC, searches already exist for the above process. An overview can e.g. be found in [27], where the authors present the status of current searches for

$$pp \rightarrow h_{125} \rightarrow h_i h_i \rightarrow XXY Y \tag{1}$$

which for such models can be read as a bound in

$$\sin^2 \alpha \times \text{BR}_{h_{125} \rightarrow h_i h_i \rightarrow XXY Y}.$$

The results are displayed in figure 7. Current bounds on the mixing angle for the 125 GeV-like state are around $|\sin \alpha| \lesssim 0.3$ (see e.g. [28]). This means that branching ratios $\text{BR}_{h_{125} \rightarrow h_i h_i \rightarrow XXY Y}$ down to $\mathcal{O}(10^{-5})$ can be tested. In particular the $\mu\mu\mu\mu$ final states in the low mass region give interesting constraints on the $h_{125} \rightarrow h_i h_i$ branching ratio down to ~ 0.03 .

See also [29] for more recent work in this context.




Figure 7. Limits on the process in eqn (1), taken from [27]. This displays current constraints which can especially be easily reinterpreted in extended scalar sector models, in particular models where couplings are inherited via a simple mixing angle. In this figure, the lighter scalar is denoted by s , which corresponds to h_i in the notation used in this manuscript.

6 Parameter space for some sample models

After investigating new physics signatures, we now turn to models that still allow for such low mass scalars. This is obviously only a brief overview, and more models might exist allong for low mass scalars accessible at Higgs factories; see e.g. [1] for more details.

The first model we discuss is a model that extends the scalar sector of the SM by two additional fields that transform as singlets under the electroweak gauge group [30, 31]. This model contains three CP-even neutral scalars that relate the gauge and mass eigenstates $h_{1,2,3}$ via mixing, where one of the three scalars has to have properties complying with current measurements of the SM-like scalar, while the other two can have higher or lower masses. A detailed discussion including all constraints can be found in [30, 31], with recent updates on benchmark planes also presented in [32]. In figure 8, two cases are shown where either one (high-low) or two (low-low) scalar masses are smaller than 125 GeV. On the y-axis, the respective mixing angle is shown. Decoupling here corresponds to $\sin \alpha = 0$.

Another option are e.g. two Higgs doublet models, where the SM scalar sector is augmented by a second doublet. In the so-called flavour-aligned scenario [33, 34], the authors perform a scan including bounds from theory, experimental searches and constraints, as e.g. electroweak observables, as well as B-physics. Here, the angle $\tilde{\alpha}$ parametrizes the rescaling with respect to the Standard Model couplings to gauge bosons, with $\cos \tilde{\alpha} = 0$ designating the SM decoupling. The limits on the absolute value of the cosine of rescaling angle vary between 0.05 and 0.25 [35]. In figure 10, we show this angle vs the different scalar masses,




Figure 8. Available parameter space in the TRSM, with one (high-low) or two (low-low) masses lighter than 125 GeV. *Left:* light scalar mass and mixing angle, with $\sin \alpha = 0$ corresponding to complete decoupling. *Right:* available parameter space in the (m_{h_1}, m_{h_2}) plane, with color coding denoting the rescaling parameter $\sin \alpha$ for the lighter scalar h_1 . Within the green triangle, $h_{125} \rightarrow h_2 h_1 \rightarrow h_1 h_1 h_1$ decays are kinematically allowed. Taken from [1].




Figure 9. Allowed regions in the 2HDM, from a scan presented in [36].

reproduced from [36]¹. We see that all regions for masses $\lesssim 125$ GeV can be populated, with absolute value of mixing angle ranges $|\cos(\tilde{\alpha})| \lesssim 0.1$. A more recent update of the above scan would of course be of high interest.

One can also extend the scalar sector further, e.g. by adding an additional singlet in the gauge eigenbasis. In [37], the authors consider a model where the SM scalar sector is extended by an additional doublet as well as a real singlet. The particle content of the model contains 3 CP even neutral scalar particles, out of which one, as before, needs to have the properties in compliance with LHC measurements of the 125 GeV scalar. The authors perform an extensive scan and find regions in parameter space where either one or both of the additional scalars have masses below 125 GeV. An example of the allowed parameter space

¹I thank V. Miralles for providing these plots.




Figure 10. Mixing angle and masses of different additional scalars in the aligned 2HDM, from the scan presented in [36]. For all additional scalars, regions exist where masses are $\lesssim 125$ GeV, with absolute values of mixing angles such that $|\cos(\tilde{\alpha})| \lesssim 0.1$. Taken from [1].




Figure 11. Scan results in the N2HDM, taken from [37]. There are regions in the models parameter space where either one or two of the additional scalars have masses $\lesssim 125$ GeV.

is displayed in figure 11. We see that in the CP-even sector there are regions within this model that still allow for low mass scalars.

Other models that allow for low mass scalars have e.g. been presented in [38, 39].

7 Conclusions

I very briefly discussed some aspects of searches for low mass scalars at Higgs factories, including models that allow for such low mass states, and provided references for further read-

ing. In particular, novel studies exceeding the ones presented here are highly encouraged and could be included as an input for e.g. the next European Strategy update.

References

- [1] Tania Robens. A Short Overview on Low Mass Scalars at Future Lepton Colliders. *Universe*, 8:286, 2022, 2205.09687.
- [2] Tania Natalie Robens. An overview on low mass scalars at future lepton colliders. *PoS, ICHEP2022*:1041, 11 2022, 2211.10231.
- [3] Richard Keith Ellis et al. Physics Briefing Book: Input for the European Strategy for Particle Physics Update 2020. 10 2019, 1910.11775.
- [4] 2020 Update of the European Strategy for Particle Physics (Brochure). Technical report, Geneva, 2020.
- [5] G. Abbiendi et al. Decay mode independent searches for new scalar bosons with the OPAL detector at LEP. *Eur. Phys. J. C*, 27:311–329, 2003, hep-ex/0206022.
- [6] S. Schael et al. Search for neutral MSSM Higgs bosons at LEP. *Eur. Phys. J. C*, 47:547–587, 2006, hep-ex/0602042.
- [7] H. Abramowicz et al. Higgs physics at the CLIC electron–positron linear collider. *Eur. Phys. J. C*, 77(7):475, 2017, 1608.07538.
- [8] Johan Alwall, Michel Herquet, Fabio Maltoni, Olivier Mattelaer, and Tim Stelzer. Mad-Graph 5 : Going Beyond. *JHEP*, 06:128, 2011, 1106.0522.
- [9] P. Drechsel, G. Moortgat-Pick, and G. Weiglein. Prospects for direct searches for light Higgs bosons at the ILC with 250 GeV. *Eur. Phys. J. C*, 80(10):922, 2020, 1801.09662.
- [10] Yan Wang, Mikael Berggren, and Jenny List. ILD Benchmark: Search for Extra Scalars Produced in Association with a Z boson at $\sqrt{s} = 500$ GeV. 5 2020, 2005.06265.
- [11] R. Barate et al. Search for the standard model Higgs boson at LEP. *Phys. Lett. B*, 565:61–75, 2003, hep-ex/0306033.
- [12] Albert M Sirunyan et al. Search for a standard model-like Higgs boson in the mass range between 70 and 110 GeV in the diphoton final state in proton-proton collisions at $\sqrt{s} = 8$ and 13 TeV. *Phys. Lett. B*, 793:320–347, 2019, 1811.08459.
- [13] S. Heinemeyer, C. Li, F. Lika, G. Moortgat-Pick, and S. Paasch. Phenomenology of a 96 GeV Higgs boson in the 2HDM with an additional singlet. *Phys. Rev. D*, 106(7):075003, 2022, 2112.11958.
- [14] Thomas Biekötter, Sven Heinemeyer, and Georg Weiglein. Mounting evidence for a 95 GeV Higgs boson. *JHEP*, 08:201, 2022, 2203.13180.
- [15] Thomas Biekötter, Sven Heinemeyer, and Georg Weiglein. Excesses in the low-mass Higgs-boson search and the W-boson mass measurement. *Eur. Phys. J. C*, 83(5):450, 2023, 2204.05975.
- [16] Rachid Benbrik, Mohammed Boukidi, Stefano Moretti, and Souad Semaili. Explaining the 96 GeV Di-photon anomaly in a generic 2HDM Type-III. *Phys. Lett. B*, 832:137245, 2022, 2204.07470.

- [17] Rachid Benbrik, Mohammed Boukidi, and Bouzid Manaut. W -mass and 96 GeV excess in type-III 2HDM. 4 2022, 2204.11755.
- [18] Weichao Li, Jingya Zhu, Kun Wang, Shiquan Ma, Pengfu Tian, and Haoxue Qiao. A light Higgs boson in the NMSSM confronted with the CMS di-photon and di-tau excesses. 12 2022, 2212.11739.
- [19] Duarte Azevedo, Thomas Biekötter, and P. M. Ferreira. 2HDM interpretations of the CMS diphoton excess at 95 GeV. 5 2023, 2305.19716.
- [20] Pablo Escribano, Victor Martin Lozano, and Avelino Vicente. A Scotogenic explanation for the 95 GeV excesses. 6 2023, 2306.03735.
- [21] T. Biekötter, S. Heinemeyer, and G. Weiglein. The 95.4 GeV di-photon excess at ATLAS and CMS. 6 2023, 2306.03889.
- [22] Alexander Belyaev, Rachid Benbrik, Mohammed Boukidi, Manimala Chakraborti, Stefano Moretti, and Souad Semaili. Explanation of the Hints for a 95 GeV Higgs Boson within a 2-Higgs Doublet Model. 6 2023, 2306.09029.
- [23] Saiyad Ashanujjaman, Sumit Banik, Guglielmo Coloretti, Andreas Crivellin, Bruce Mellado, and Anza-Tshildizi Mulaudzi. $SU(2)_L$ triplet scalar as the origin of the 95 GeV excess? 6 2023, 2306.15722.
- [24] Zhen Liu, Lian-Tao Wang, and Hao Zhang. Exotic decays of the 125 GeV Higgs boson at future e^+e^- lepton colliders. *Chin. Phys. C*, 41(6):063102, 2017, 1612.09284.
- [25] Jonathan Kozaczuk, Michael J. Ramsey-Musolf, and Jessie Shelton. Exotic Higgs boson decays and the electroweak phase transition. *Phys. Rev. D*, 101(11):115035, 2020, 1911.10210.
- [26] Zhen Wang, Xuliang Zhu, Elham E. Khoda, Shih-Chieh Hsu, Nikolaos Konstantinidis, Ke Li, Shu Li, Michael J. Ramsey-Musolf, Yanda Wu, and Yuwen E. Zhang. Study of Electroweak Phase Transition in Exotic Higgs Decays at the CEPC. In *Snowmass 2021*, 3 2022, 2203.10184.
- [27] Maria Cepeda, Stefania Gori, Verena Martinez Outschoorn, and Jessie Shelton. Exotic Higgs Decays. 11 2021, 2111.12751.
- [28] Tania Robens. Constraining Extended Scalar Sectors at Current and Future Colliders—An Update. *Springer Proc. Phys.*, 292:141–152, 2023, 2209.15544.
- [29] Marcela Carena, Jonathan Kozaczuk, Zhen Liu, Tong Ou, Michael J. Ramsey-Musolf, Jessie Shelton, Yikun Wang, and Ke-Pan Xie. Probing the Electroweak Phase Transition with Exotic Higgs Decays. *LHEP*, 2023:432, 2023, 2203.08206.
- [30] Tania Robens, Tim Stefaniak, and Jonas Wittbrodt. Two-real-scalar-singlet extension of the SM: LHC phenomenology and benchmark scenarios. *Eur. Phys. J. C*, 80(2):151, 2020, 1908.08554.
- [31] Tania Robens. Two-Real-Singlet-Model Benchmark Planes. *Symmetry*, 15(1):27, 2023, 2209.10996.
- [32] Tania Robens. Two-Real-Singlet Model Benchmark Planes – A Moriond Update. In *57th Rencontres de Moriond on QCD and High Energy Interactions*, 5 2023,

2305.08595.

- [33] Antonio Pich and Paula Tuzon. Yukawa Alignment in the Two-Higgs-Doublet Model. *Phys. Rev. D*, 80:091702, 2009, 0908.1554.
- [34] Antonio Pich. Flavour constraints on multi-Higgs-doublet models: Yukawa alignment. *Nucl. Phys. B Proc. Suppl.*, 209:182–187, 2010, 1010.5217.
- [35] Combined measurements of Higgs boson production and decay using up to 139 fb^{-1} of proton-proton collision data at $\sqrt{s} = 13 \text{ TeV}$ collected with the ATLAS experiment. Technical report, CERN, Geneva, Nov 2021. ATLAS-CONF-2021-053.
- [36] Otto Eberhardt, Ana Peñuelas Martínez, and Antonio Pich. Global fits in the Aligned Two-Higgs-Doublet model. *JHEP*, 05:005, 2021, 2012.09200.
- [37] Hamza Abouabid, Abdesslam Arhrib, Duarte Azevedo, Jaouad El Falaki, Pedro. M. Ferreira, Margarete Mühlleitner, and Rui Santos. Benchmarking di-Higgs production in various extended Higgs sector models. *JHEP*, 09:011, 2022, 2112.12515.
- [38] Carlos Henrique de Lima and Heather E. Logan. Unavoidable Higgs coupling deviations in the Z_2 -symmetric Georgi-Machacek model. *Phys. Rev. D*, 106(11):115020, 2022, 2209.08393.
- [39] A. Kunčinas, O. M. Ogreid, P. Osland, and M. N. Rebelo. Complex S_3 -symmetric 3HDM. *JHEP*, 07:013, 2023, 2302.07210.