Reports on Progress in **Physics**

PURPOSE-LED PUBLISHING™

PAPER • **OPEN ACCESS**

Combined measurements and interpretations of Higgs boson production and decay in proton–proton collisions at $\sqrt{s} = 13\,\text{TeV}$

To cite this article: The CMS Collaboration 2026 *Rep. Prog. Phys.* **89** 087801

View the article online for updates and enhancements.

This content was downloaded from IP address 161.53.22.193 on 27/11/2026 at 13:15

Reports on Progress in Physics

Check for updates

PAPER

Combined measurements and interpretations of Higgs boson production and decay in proton–proton collisions at $\sqrt{s} = 13\,\text{TeV}$

The CMS Collaboration[1,*]

Peking University, Beijing, People's Republic of China
[1] CERN, Geneva, Switzerland.
[*] Author to whom any correspondence should be addressed.

E-mail: cms-publication-committee-chair@cern.ch

OPEN ACCESS

RECEIVED
20 February 2026

REVISED
11 May 2026

ACCEPTED FOR PUBLICATION
26 May 2026

PUBLISHED
10 August 2026

Keywords: CMS, Higgs boson, statistical combination

Corresponding editor: Dr Lorna Brigham

Original content from this work may be used under the terms of the Creative Commons Attribution 4.0 licence.

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Abstract

Combined measurements of Higgs boson production and decay rates are reported, representing the most comprehensive study performed by the CMS Collaboration to date. The included analyses use proton–proton collision data recorded by the CMS experiment at $\sqrt{s} = 13\,\text{TeV}$ from 2016 to 2018, corresponding to an integrated luminosity of $138\,\text{fb}^{-1}$. The statistical combination is based on analyses that measure the following decay channels: $\text{H} \to \gamma\gamma$, $\text{H} \to \text{ZZ}$, $\text{H} \to \text{WW}$, $\text{H} \to \tau\tau$, $\text{H} \to \text{bb}$, $\text{H} \to \mu\mu$, and $\text{H} \to \text{Z}\gamma$. Information in the events from each decay channel is used to target multiple Higgs boson production processes. Searches for invisible Higgs boson decays are also considered, as well as an analysis that measures off-shell Higgs boson production in the $\text{H} \to \text{ZZ} \to 4\ell$ ($\ell = \text{e}, \mu$) decay channel. The best fit inclusive signal yield is measured to be $1.014^{+0.055}_{-0.053}$ times the standard model (SM) expectation, for a Higgs boson mass of 125.38 GeV. Measurements in kinematic regions defined by the simplified template cross section framework are also provided, as well as interpretations in the coupling modifier and SM effective field theory frameworks. The coupling modifier interpretation is further used to place constraints on various two-Higgs-doublet models. The results show good compatibility with the SM predictions for the majority of the measured parameters.

Contents

1. Introduction 2
2. The CMS detector 4
3. Analyses included in the combination 5
   3.1. $\text{H} \to \gamma\gamma$ 5
   3.2. $\text{H} \to \text{ZZ}$ 5
   3.3. $\text{H} \to \text{WW}$ 7
   3.4. $\text{H} \to \tau\tau$ 7
   3.5. $\text{H} \to \text{bb}$ 7
   3.6. t(t)H production with $\text{H} \to$ leptons 8
   3.7. $\text{H} \to \mu\mu$ 8
   3.8. $\text{H} \to \text{Z}\gamma$ 9
   3.9. $\text{H} \to$ invisible particles 9
   3.10. Off-shell production 9
4. Modifications to the input analyses 9
5. Statistical procedure 10
   5.1. The combined likelihood 10
   5.2. Signal yield parametrization 11
   5.3. Extraction of results 12

© CERN, for the benefit of the CMS Collaboration

6. Systematic uncertainties 13
6.1. Theoretical uncertainties 13
6.2. Experimental uncertainties 13
7. Measurements of signal strengths 15
8. Measurements of simplified template cross sections and decay branching fractions 19
8.1. Stage 0 measurements 19
8.2. Stage 1.2 measurements 24
9. Interpretation of measurements in the coupling modifier framework 31
9.1. Constraints on coupling modifiers with resolved loops 34
9.2. Constraints on coupling modifiers with effective loops 36
9.3. Constraints on coupling modifier ratios 39
9.4. Constraints on the Higgs boson self-coupling 39
10.Interpretation of measurements using benchmark two-Higgs-doublet models 44
11.Interpretation of measurements using SMEFT 47
11.1. Derivation of the SMEFT parametrization 49
11.2. Impact of Higgs boson kinematic properties 52
11.3. SMEFT parametrization summary 53
11.4. Constraints on individual Wilson coefficients 54
11.5. Constraints on linear combinations of Wilson coefficients 54
12.Summary 64
Acknowledgments 67
Data availability statement 68
Appendix. Comparison of SMEFT constraints with a simplified likelihood procedure 68
References 87

1. Introduction

In 2012, the ATLAS [1] and CMS [2] Collaborations announced the observation of a particle [3–5] with properties consistent with the Higgs boson (H) of the standard model (SM) [6–11], a neutral scalar particle that arises from the Brout–Englert–Higgs mechanism responsible for electroweak symmetry breaking. Since then, a broad set of measurements has been performed to characterize the properties and interactions of this particle with the highest achievable precision. The proton–proton collision data collected during Run 2 (2015–2018) of the CERN LHC [12] have enabled the ATLAS and CMS Collaborations to make significant advances in understanding the Higgs boson, and thus to perform precision tests of the SM.

This paper describes combined measurements of Higgs boson production and decay rates. The analyses included in the statistical combination measure the following decay channels: $\mathrm{H} \to \gamma\gamma$, $\mathrm{H} \to \mathrm{ZZ}(\to 4\ell)$, $\mathrm{H} \to \mathrm{WW}(\to \ell\nu\ell\nu$ or $\ell\nu\mathrm{qq}')$, $\mathrm{H} \to \tau\tau$, $\mathrm{H} \to \mathrm{bb}$, $\mathrm{H} \to \mu\mu$, and $\mathrm{H} \to \mathrm{Z}\gamma(\to \ell\ell\gamma)$, as shown by the Feynman diagrams in figure 1. In these expressions and throughout the paper, $\ell = \mathrm{e}$ or μ, and the distinction between a particle and its antiparticle is dropped. The $\mathrm{H} \to \mathrm{ZZ}$ and $\mathrm{H} \to \mathrm{WW}$ decay channels, where the Higgs boson is on-shell, require at least one vector boson (W, Z) to be off-shell. In addition, analyses that search for invisible decays of the Higgs boson ($\mathrm{H} \to \mathrm{inv}$) are incorporated to constrain its branching fraction into beyond-the-SM (BSM) particles. No analyses targeting the $\mathrm{H} \to \mathrm{cc}$ decay channel are included, as a result of the lack of statistical independence between them and the $\mathrm{H} \to \mathrm{bb}$ analyses included in the combination.

Information in the events from each decay channel is used to target multiple Higgs boson production mechanisms, including Higgs boson production via gluon fusion (ggH), vector boson fusion (VBF), production in association with a vector boson (VH, V = W or Z), production in association with a pair of top quarks (ttH), and production in association with a single top quark (tH). There are two ways in which ZH production occurs: the dominant production mechanism is through the quark-initiated process (qqZH), with a smaller contribution from the gluon-initiated process (ggZH). Similarly, tH production includes contributions from two processes: production in association with either a W boson (tHW) or a quark (tHq). The input analyses are not sensitive to production in association with a pair of bottom quarks (bbH). However, small contributions from this production process are accounted for in the signal modelling. Contributions from Higgs boson pair production in the input analyses are negligible compared to single Higgs boson production, and are therefore not considered. Example leading-order (LO) Feynman diagrams for each Higgs boson production process considered in the combination are shown



Figure 1. Examples of leading-order Feynman diagrams for the H → ZZ and H → WW decay channels (upper left); for the fermionic decay channels H → ff, where ff = bb, $\tau\tau$ or $\mu\mu$ (upper right); and for the H → $\gamma\gamma$ and H → Zγ decay channels (lower).



Figure 2. Examples of leading-order Feynman diagrams for the ggH (upper left), VBF (upper middle), quark-initiated VH (upper right), gluon-initiated ZH (middle left), ttH and bbH (middle right), tHq (lower left), and tHW (lower right) processes.

in figure 2. An analysis that measures off-shell Higgs boson production is also included in the combined measurement to constrain the Higgs boson total decay width directly from data.

The prediction for ggH production is taken from calculations at next-to-next-to-next-to-leading order (N^3LO) in perturbative quantum chromodynamics (QCD), including next-to-leading order (NLO) electroweak (EW) corrections. The VBF and VH cross section calculations are performed at next-to-next-to-leading order (NNLO) accuracy in QCD and NLO accuracy in EW theory. The theoretical predictions of the ttH cross section are derived at NLO accuracy in both QCD and EW, while the tH cross sections are derived at NLO QCD accuracy in the five-flavour scheme (5FS), with no EW corrections. The bbH cross section calculation includes matched predictions at NNLO QCD accuracy in the 5FS, and at NLO accuracy in the four-flavour scheme (4FS), with no EW corrections. The theoretical predictions for the various production processes and decay modes in this paper are summarized in [13], with more details provided in the references within, particularly [14–21], which describe the cross section calculations for the targeted production processes.

This paper also presents combined measurements of simplified template cross sections (STXS). In the STXS framework [13], the phase space of the different Higgs boson production processes is partitioned

into multiple nonoverlapping regions, based on the kinematic properties of the Higgs boson events. The partitions are designed to isolate regions with enhanced sensitivity to potential BSM effects, to reduce theoretical uncertainties in the SM predictions, and to limit model dependence by ensuring close correspondence with the experimental acceptance. Measurements of these phase space regions, referred to as 'STXS bins', thereby provide more granular information beyond the inclusive production and decay rate measurements. In this paper, the STXS measurements are performed at two levels of granularity. At STXS stage 0, the bins correspond closely to the main Higgs boson production processes. Subsequent stages perform a further splitting of the bins using the kinematic properties of the events. In this combined measurement, the STXS stage 1.2 binning scheme is used [22]. Interpretations of the measurements in the context of the coupling modifier [23] and SM effective field theory (SMEFT) [24] frameworks are also provided.

The ATLAS and CMS Collaborations have published combined measurements of Higgs boson production and decay rates, and couplings, with the $\sqrt{s} = 7$ and 8 TeV LHC Run 1 (2011–2012) data [25, 26]. A combination of the Run 1 ATLAS and CMS analyses has also been performed [27]. Combined measurements of Higgs boson production and decay rates, as well as Higgs boson couplings, have been conducted by both collaborations with data collected during Run 2 [28–32]. The combined measurements performed by ATLAS with data collected during Run 2 provide measurements and interpretations in the STXS framework [28–30]. This paper presents the first combined STXS measurement from CMS. With respect to the previous CMS Higgs boson combination, detailed in [32], the combination presented here uses additional input analyses: measurements of the $\mathrm{H} \to \mathrm{bb}$ decay channel via the VBF production process [33], and via production at high transverse momentum [34]; measurements of off-shell Higgs boson production in the $\mathrm{H} \to \mathrm{ZZ} \to 4\ell$ decay channel [35], and a search for the $\mathrm{H} \to$ inv decay via ttH and VH production in which the vector boson decays hadronically [36]. Furthermore, larger data sets are used for the VH ($\mathrm{H} \to \mathrm{bb}$) [37], and t(t)H ($\mathrm{H} \to \mathrm{bb}$) [38] input measurements, where t(t)H refers to analyses that target both the ttH and tH production processes.

The SM predictions for the Higgs boson production and decay rates depend on the mass of the Higgs boson m_{H}. For all measurements in this paper, the mass is fixed at $m_{\mathrm{H}} = 125.38\,\mathrm{GeV}$. This was the most precise measurement of m_{H} ($\pm 0.14\,\mathrm{GeV}$) by the CMS Collaboration at the time that the analyses entering the combination were performed [39]. Since then, a more precise measurement of $m_{\mathrm{H}} = 125.08 \pm 0.12\,\mathrm{GeV}$ has been performed by CMS in the $\mathrm{H} \to \mathrm{ZZ} \to 4\ell$ channel [35]. The ATLAS Collaboration also performed a more precise measurement of $m_{\mathrm{H}} = 125.11 \pm 0.11\,\mathrm{GeV}$, combining the $\mathrm{H} \to \mathrm{ZZ} \to 4\ell$ and $\mathrm{H} \to \gamma\gamma$ channels [40]. The small difference in m_{H} between these values has a negligible effect on the results in this paper.

This paper is organized as follows. A brief description of the CMS detector is given in section 2. Section 3 provides a summary of the analyses included in the combination, and section 4 describes the modifications made to input analyses, to ensure a common signal and uncertainty model. The statistical procedure for the combination is outlined in section 5, and the treatment of systematic uncertainties is summarized in section 6. The following sections report the results of the different signal yield parametrizations. Section 7 shows the measurements of signal strength modifiers at different levels of granularity. The measurements of the STXS and decay channel branching fractions are presented in section 8. The interpretation of these measurements in the coupling modifier framework is shown in section 9, including constraints on the Higgs boson trilinear self-coupling via NLO EW corrections. In section 10, the coupling modifier constraints are used to probe two-Higgs-doublet models, which are extensions of the SM that remain valid up to high energies. Section 11 details the interpretations in terms of SMEFT parameters. Finally, a summary of the results is presented in section 12.

The tabulated results for all measurements and interpretations are provided in the HEPData record for this paper [41].

2. The CMS detector

The CMS apparatus [2, 42] is a multipurpose, nearly hermetic detector, designed to trigger on [43–45] and identify electrons, muons, photons, and hadrons [46–48]. Its central feature is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter, each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity coverage provided by the barrel and endcap detectors. Muons are reconstructed using gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid.

More detailed descriptions of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in [2, 42].

3. Analyses included in the combination

All analyses included in the combination use a proton–proton collision data set recorded by the CMS experiment between 2016 and 2018, corresponding to an integrated luminosity of 138 fb^{-1}, unless otherwise stated.

Most analyses are designed to target a single Higgs boson decay channel. The data are split into multiple event categories to improve the sensitivity to the signal, as well as to enhance the discrimination power between different Higgs boson production processes. The latter is achieved by requiring the presence of additional leptons or jets, where a jet refers to the collimated streams of particles arising from the fragmentation of a quark or gluon. Such additional objects arise in the VBF production process, in the decay of a W or Z boson from the WH and ZH production processes, or in top quark decays from the ttH and tH processes. The properties of the final-state objects are used to further enhance the purity of the targeted Higgs boson production process. For example, the VBF topology is characterized by the presence of two jets with a large separation in pseudorapidity $\Delta\eta_{jj}$, and a large invariant mass m_{jj}. Alternatively, the identification of two additional jets as originating from the hadronization of b quarks (b tagging) is indicative of the ttH production process. Most analyses leverage machine learning (ML) algorithms to optimally use the high-dimensional event information.

A subset of input analyses perform measurements in the STXS framework [13]. To achieve this, events are categorized based on the reconstructed kinematic information. This categorization is performed either using an ML classifier, or by simply placing a selection on the equivalent reconstructed quantity, e.g. using $p_{\text{T}}^{\gamma\gamma}$ for p_{T}^{H} STXS partitions in the $\text{H} \to \gamma\gamma$ analysis. As a result of limited statistical precision, and therefore constraining power, analysis categories do not necessarily map one-to-one with the STXS bins. In the individual input measurements, STXS bins that could not be measured because of a lack of events were either fixed to the SM expectation, or, more frequently, merged with neighbouring bins. The combined measurement, however, is sensitive to all STXS partitions that were measured across the input analyses. More information regarding the STXS stage 1.2 binning scheme used in the combination is provided in section 8.2.

The input analyses are required to be statistically independent to enter the combination. As the analyses mostly target different final states, the event categories have a negligible level of overlap by construction. For analyses that target similar final states, the overlap is explicitly checked and found to be negligible.

Several small modifications, summarized in section 4, are made to the statistical models used in each analysis to make them mutually compatible for the combination. To validate this procedure, the modified statistical models are fit and the results are compared to the original publications. The fitted values of the parameters of interest (POIs) are found to be compatible with the original publications within the parameter uncertainties.

In total, up to 1951 event categories are considered in the combined fit, with over 10^4 fit parameters corresponding to the POIs and various sources of experimental and theoretical systematic uncertainties. A summary of the input analyses is shown in table 1. Additional details for each of the input analyses are provided in the remainder of this section.

3.1. $\text{H} \to \gamma\gamma$

In the $\text{H} \to \gamma\gamma$ decay channel, the ggH, VBF, VH, ttH, and tH production processes are analyzed. Despite the relatively small branching fraction (∼0.2%), this channel provides very good sensitivity in Higgs boson measurements as a result of the excellent photon energy resolution of the CMS ECAL [46, 59]. The signal extraction is performed by measuring a narrow signal peak in the diphoton invariant mass spectrum, on top of a smoothly falling background distribution. Measurements of the STXS stage 1.2 bins, described in section 8.2, are performed for all Higgs boson production processes. To improve the sensitivity of the analysis to these different production processes and STXS bins, different signal categories are defined based on the kinematic properties of the events. More details concerning the analysis and its measurements are provided in [49].

3.2. $\text{H} \to \text{ZZ}$

For the on-shell $\text{H} \to \text{ZZ}$ channel, with both Z bosons decaying leptonically ($\text{H} \to \text{ZZ} \to 4\ell$), the ggH, VBF, and VH production processes are studied with dedicated analysis categories. The analysis also

Table 1. Summary of the input analyses in the combination. The second column lists the final-state particles that are targeted in each analysis. Here, τ_h indicates a hadronically decaying τ lepton and p_T^{miss} is the magnitude of the missing transverse momentum vector. The third column lists the production processes and STXS kinematic regions targeted in each analysis. The variables NJ, p_T^H, m_{jj}, p_T^{Hjj}, and p_T^V refer to the jet multiplicity, the transverse momentum of the Higgs boson, the invariant mass of the two leading jets, the transverse momentum of the Higgs plus dijet system, and the transverse momentum of the vector boson, respectively. The fourth column indicates the granularity of the signal model used in each analysis. More information on each analysis can be found in the references listed in the last column.

Input analysis	Final states targeted	Production targeted	Signal granularity	References
$H \to \gamma\gamma$	$\gamma\gamma$	ggH, (NJ, p_T^H, m_{jj}) bins VBF, (p_T^H, m_{jj}, p_T^{Hjj}) bins VH hadronic WH leptonic, p_T^V bins ZH leptonic ttH, p_T^H bins tH	STXS stage 1.2	[49]
$H \to ZZ \to 4\ell$	4μ, $2e2\mu$, 4e	ggH, (NJ, p_T^H, m_{jj}) bins VBF, (NJ, p_T^H, m_{jj}, p_T^{Hjj}) bins VH hadronic VH leptonic, p_T^V bins ttH	STXS stage 1.2	[50]
$H \to WW$	$e\mu/\mu e$, $ee/\mu\mu$, $e\mu$+jj 3ℓ, 4ℓ	ggH, (NJ, p_T^H) bins VBF-like (p_T^H, m_{jj}) bins VH hadronic WH leptonic, p_T^V bins ZH leptonic, p_T^V bins	STXS stage 1.2	[51]
$H \to \tau\tau$	$e\mu$, $e\tau_h$, $\mu\tau_h$, $\tau_h\tau_h$ $e\tau_h$+2ℓ, $\mu\tau_h$+ℓ/2ℓ, $\tau_h\tau_h$+ℓ/2ℓ	ggH, (NJ, p_T^H, m_{jj}) bins VBF, (NJ, p_T^H, m_{jj}) bins WH leptonic, p_T^V bins ZH leptonic, p_T^V bins	Inclusive production processes or STXS stage 1.2	[52]
$H \to bb$ boosted	bb	ggH, high-p_T^H bins VBF, high-p_T^H bins	Inclusive production processes or STXS stage 1.2	[34]
VBF ($H \to bb$)	bb	VBF	Inclusive production processes	[33]
VH ($H \to bb$)	bb	WH leptonic, p_T^V bins ZH leptonic, p_T^V bins	STXS stage 1.2	[37]
t(t)H ($H \to bb$)	bb	ttH, p_T^H bins tH	Inclusive production processes or STXS stage 1.2	[38]
t(t)H ($H \to$ leptons)	2ℓ(same charge), 3ℓ, 4ℓ 1ℓ+2τ_h, 2ℓ(same charge)+1τ_h, 3ℓ+1τ_h	ttH, tH	STXS stage 1.2	[53]
$H \to \mu\mu$	$\mu\mu$	ggH, VBF, VH, ttH	Inclusive production processes	[54]
$H \to Z\gamma \to \ell\ell\gamma$	$\ell\ell\gamma$	ggH, VBF	Inclusive production processes	[55]
$H \to$ inv	Large p_T^{miss}	Monojet VBF VH leptonic VH/ttH hadronic	Inclusive production processes	[36, 56–58]
Off-shell ($H \to ZZ \to 4\ell$)	4ℓ	Off-shell	Off-shell contributions (cf equation (16))	[35]

includes ttH and tH production, although the precision with which these processes are measured is limited. This Higgs boson decay channel, despite its small branching fraction (∼0.01%), has an excellent signal-to-background ratio for the major production processes. The signal extraction makes use of multi-dimensional fits of the reconstructed Higgs boson mass and kinematic discriminators that describe the 4-lepton decay. In the analysis, the STXS stage 1.2 bins are measured. Sensitivity to the STXS bins is obtained by categorizing the events based on the kinematic properties of the Higgs boson production and the presence of additional particles in the event. More details on the analysis can be found in [50].

3.3. H → WW

The analysis of the H → WW channel is described in [51]. The analysis is restricted to final states with at least two charged leptons, thus targeting the H → WW → $\ell\nu\ell\nu$ decay for all H production processes, and additionally the H → WW → $\ell\nu$qq′ decay for VH production. Considering only leptonic final states results in the suppression of backgrounds, while retaining moderate branching fractions of ∼2% (∼9%) for H → WW → $\ell\nu\ell\nu$ (H → WW → $\ell\nu$qq′). The analysis is most sensitive to the ggH and VBF production processes. Additional categories with three and four leptons are constructed to target the production of the Higgs boson in association with a leptonically decaying vector boson. The analysis does not target ttH and tH production, which are covered by a dedicated analysis studying t(t)H production in multileptonic final states (described in section 3.6). In the analysis, the STXS stage 1.2 bins are also measured. To increase the sensitivity of these measurements, the event categories are aligned as closely as possible with the STXS bins.

3.4. H → $\tau\tau$

The analysis of the H → $\tau\tau$ decay channel measures Higgs boson production via the ggH, VBF, and VH processes, and is described in detail in [52]. As in the channels described previously, this analysis measures bins at stage 1.2 of the STXS framework. Four different di-τ final states are studied: eμ, eτ_h, $\mu\tau_h$, and $\tau_h\tau_h$, where τ_h refers to a hadronically decaying τ lepton. Although the H → $\tau\tau$ decay channel has a smaller branching fraction (∼6%) than the H → bb and H → WW modes, the leptons in the final state and the good performance of the τ_h identification [60] enable this channel to balance a moderate branching fraction with moderate background contributions. This channel is particularly sensitive to VBF and ggH production, but also has sensitivity to the VH production process, where the vector boson decays leptonically. Specific analysis categories with additional leptons are used to measure these events. The analysis does not target ttH or tH production, which are covered by the t(t)H multilepton analysis described in section 3.6.

The VH categories are defined by placing requirements on the kinematic properties of the events. To define categories that are sensitive to ggH and VBF production, a deep neural network (DNN) that takes the kinematic properties of the event as input, is used. Independent networks are trained for the measurement of the STXS bins and for the measurement of the inclusive production rates. Two different sets of analysis categories are therefore used for these two measurements. In the combination, the analysis categories that are most sensitive to the signal parametrization being fit are used, as detailed in table 2 of section 5.2 where the signal yield parametrization is introduced.

3.5. H → bb

Out of all the Higgs boson decay modes, H → bb has the largest branching fraction (∼60%). Several analyses that measure different Higgs boson production processes in the H → bb channel are included in the combination. The inputs have changed significantly since the combined measurement described in [32]. The analyses that measure VH, ttH and tH production, and Higgs boson production at high transverse momentum ($p_T^H > 450$ GeV) have all been updated, and a measurement of the VBF process in this decay channel has been included for the first time.

The analysis of VH production uses the presence of one or more leptons, or large missing transverse momentum p_T^{miss}, to target the WH and ZH production processes with the vector boson decaying leptonically. Here, p_T^{miss} is defined as the magnitude of the negative vector p_T sum of all the particle-flow candidates [61] in an event. Requiring a leptonically decaying vector boson significantly reduces the background from events where jets are produced through the strong interaction, referred to as QCD multijet production. This, combined with the large H → bb branching fraction, leads to this analysis being the most sensitive to the VH production process out of all the Higgs boson decay channels. Backgrounds from top quark pair production and vector boson production in association with jets (V+jets) are constrained in dedicated control regions (CRs). In the signal-sensitive region, DNNs are used to separate signal from background. This analysis measures the STXS stage 1.2 bins defined for the VH production process. More details are provided in [37].

A dedicated analysis of ttH and tH production in the H → bb final state is also included in the combination. All decay channels of the t(t) system are considered. Several measurements are performed, from inclusive measurements of the ttH and tH production processes, to measurements of the ttH process in bins of p_T^H according to the STXS stage 1.2 framework. The analysis employs DNNs to discriminate signal from background, categorize events in the signal region, and define background CRs. As in the case of H → $\tau\tau$, different sets of analysis categories are constructed for the measurement of inclusive quantities and for the measurement of the STXS bins. In the combination, the analysis categories that

are most sensitive to the signal yield parametrization being probed are used, as detailed in table 2. More information regarding the t(t)H ($H \to bb$) analysis is provided in [38].

An analysis measuring Higgs bosons produced with $p_T^H > 450\,GeV$, and decaying via $H \to bb$, is included in the combined measurement. The decay of a high p_T Higgs boson to a Lorentz-boosted bottom quark pair is identified by selecting large-radius jets, and employing jet substructure and heavy-flavour taggers based on advanced ML algorithms [62]. Independent regions targeting VBF and ggH production are defined using the presence or absence of forward jets. The analysis measures both the inclusive signal strengths of ggH and VBF, as well as the p_T^H and m_{jj} spectra for the ggH and VBF production processes, respectively. Following the procedure of the $H \to \tau\tau$ and t(t)H ($H \to bb$) analyses, two sets of analysis categories are constructed for the inclusive and differential measurements, and table 2 shows which set is used for the different signal yield parametrizations considered in the combination. The analysis is described in detail in [34].

Finally, an analysis of Higgs boson production via VBF in the $H \to bb$ final state is included in the combined measurement for the first time. This analysis targets the resolved final state, in which the bottom quark pair is reconstructed as two separate jets. The data set used corresponds to an integrated luminosity of 90.8 fb^{-1}, collected in 2016 and 2018. The 2017 data set is not included because no suitable triggers were available that year. A DNN is used to improve the resolution of the dijet mass, and boosted decision trees based on the event kinematics and particle identification criteria are employed to discriminate signal from major background sources. This analysis measures the inclusive VBF signal strength, and the signal templates are not split according to the STXS stage 1.2 binning. More information is provided in [33].

In recent years, the CMS Collaboration has published several searches targeting the $H \to cc$ decay channel. Analyses probing $H \to cc$ in the VH production process [63], and in events with Higgs bosons produced at high p_T ($p_T^H > 450\,GeV$) [64], are not statistically independent of the corresponding $H \to bb$ analyses included in the combination. In addition, a simultaneous measurement of the $H \to bb$ and $H \to cc$ decay channels via ttH and tH production [65] was recently performed. Although this analysis ensures statistical independence through the use of orthogonal analysis regions, and is more recent than the t(t)H ($H \to bb$) analysis included in the combination, it does not categorize events according to the STXS binning scheme. Consequently, no analyses explicitly targeting the $H \to cc$ decay channel are included in the combination.

3.6. t(t)H production with $H \to$ leptons

An analysis of ttH and tH production in final states with multiple leptons is included. As leptonic Higgs boson final states are targeted, the measurement is sensitive to the $H \to WW$, $H \to \tau\tau$, and $H \to ZZ$ decay channels. The overlap between this analysis and the dedicated $H \to ZZ \to 4\ell$, $H \to WW$, and $H \to \tau\tau$ analyses is negligible as a result of the design of the event categories. The analysis uses a categorization based on the number of leptons and τ_h candidates to target both the different Higgs boson final states and the different top quark decay processes. Categories with at least two leptons, or one lepton and two τ_h candidates, probe events where at least one top quark decays via a leptonically decaying W boson. The categories with one lepton and one τ_h, or with no leptons and two τ_h's, probe events in which both top quarks decay hadronically. In the analysis, ML algorithms are employed to better separate events from the ttH and tH production processes. More details on this analysis can be found in [53]. The STXS bins are not explicitly measured in this analysis. However, the signal contributions in the statistical model are defined with the granularity of the STXS stage 1.2 to enable this channel to enter the combined STXS measurement.

3.7. $H \to \mu\mu$

The analysis of the $H \to \mu\mu$ decay channel is described in [54]. Similarly to the case of $H \to \gamma\gamma$, the challenge posed by the small branching fraction of the $H \to \mu\mu$ decay ($\sim$0.02%) is somewhat mitigated by the excellent dimuon mass resolution of the CMS detector [47]. The ggH, VBF, VH, and ttH production processes are probed by constructing categories based on the kinematic properties of the event. The highest sensitivity is obtained for the ggH and VBF production processes. ML algorithms are used to improve the sensitivity of the measurement. For the ggH production process, categories are defined based on the output of a boosted decision tree to identify regions with increased signal purity. Candidate events are then characterized by a peak in the dimuon mass distribution, over a smoothly falling background. A similar procedure is employed for VH and ttH production. For the VBF process, a DNN is instead used to separate signal and background. The DNN output is used directly in a template fit for the signal extraction.

3.8. H → Zγ

The analysis of the H → Zγ channel targets the decay of the Z boson into two electrons or two muons (H → Zγ → ℓℓγ), which has a total branching fraction of ∼0.01%. Events are divided into different categories based on the production process to increase the sensitivity to the signal. ML algorithms are used to further divide these regions into categories with higher and lower signal-to-background ratios. The signal extraction is performed by fitting the dilepton-plus-photon invariant mass distribution, where the signal appears as a narrow peak over a smoothly falling background. No STXS bins are measured in this analysis. More details can be found in [55].

3.9. H → invisible particles

Several analyses that search for the decay of the Higgs boson into particles that cannot be detected are included in the combined measurement. These analyses provide a constraint on the invisible Higgs boson decay branching fraction, and their inclusion in the relevant measurements of the coupling modifier framework is described in section 9. The analyses primarily select events with large $p_{\mathrm{T}}^{\mathrm{miss}}$, but use additional event information to identify Higgs boson production via ggH, VBF, VH, and ttH. The VBF channel is the most sensitive to the invisible Higgs boson decay branching fraction, and more details concerning this analysis can be found in [57]. The analyses probing ggH production and VH production in which the vector boson decays leptonically are described in [56, 58], respectively. An analysis that probes ttH and VH production in the fully hadronic final state [36] is included in the combined measurement for the first time.

3.10. Off-shell production

For the first time, analysis regions targeting off-shell Higgs boson production are included in the combined results. Off-shell contributions enable the Higgs boson total decay width to be constrained directly from data, as a result of the different ways in which the on-shell and off-shell production depend on the Higgs boson couplings (described in section 9.2). This means that various assumptions on the Higgs boson couplings can be alleviated when considering BSM contributions in Higgs boson decays. The input analysis, described in more detail in [35], includes categories targeting both on-shell and off-shell production of the Higgs boson in the H → ZZ → 4ℓ decay. Only the off-shell event categories are included in the combination, as the on-shell categories are not statistically independent from the H → ZZ → 4ℓ analysis described in section 3.2. Statistical independence is ensured for the off-shell categories by selecting events with a four-lepton invariant mass $m_{4\ell} > 220\,\mathrm{GeV}$, well above the range used in the on-shell H → ZZ → 4ℓ analysis ($105 < m_{4\ell} < 140\,\mathrm{GeV}$).

4. Modifications to the input analyses

A number of small modifications are made to the statistical models of the input analyses to ensure a coherent treatment of the signal contributions, as well as the systematic uncertainties.

In some of the input analyses, Higgs boson production processes or decay channels that were not targeted by the measurements were treated as backgrounds and fixed to the SM prediction. In the combination, all single Higgs boson contributions are treated as signal, in that their yields are parametrized in terms of the POIs, as described in section 5.2.

The inclusive production process cross sections and branching fractions recommended by the LHC Higgs Working Group [13] are used to determine the SM predictions. For the STXS measurements and the subsequent interpretations, it is necessary to define the SM predictions of the STXS bin cross sections in the statistical model. The relative fraction of each STXS bin for each production process was taken from Monte Carlo (MC) simulation. As different input analyses made use of different event generators to simulate the signal processes, the relative fractions could vary between analyses. The discrepancies have been addressed by rescaling the production cross sections in the STXS bins to reference values. The reference values are calculated at NLO accuracy in QCD with MadGraph5_aMC@NLO (v2.4.2) for ggH, VBF, ttH, and tH production [66]. The ggH events are weighted as a function of $p_{\mathrm{T}}^{\mathrm{H}}$ and the number of jets to match the predictions from the NNLOPS program [67]. The POWHEG v2 generator is used to obtain the reference values for the VH production processes, where the MiNLO procedure is used for quark-initiated processes (WH and qqZH) [68, 69] and an LO generation is used for ggZH production. The VH processes are corrected to NLO EW accuracy as a function of $p_{\mathrm{T}}^{\mathrm{V}}$. All reference parton-level samples are interfaced with PYTHIA 8.230 [70] for parton showering and hadronization, using the CP5 tune [71]. The parton distribution functions (PDFs) are taken from the NNPDF 3.1 set at NNLO accuracy [72].

The sample of Higgs boson events produced through ggH does not provide an accurate description at high $p_{\mathrm{T}}^{\mathrm{H}}$ (>450 GeV). The reference cross section values for the ggH $450 < p_{\mathrm{T}}^{\mathrm{H}} < 650$ GeV and ggH $p_{\mathrm{T}}^{\mathrm{H}} > 650$ GeV STXS bins are taken from the dedicated boosted Higgs boson predictions in [73]. The reference SM predictions for all measured STXS bins are provided in section 8.2, where the combined STXS stage 1.2 measurements are presented.

The theoretical uncertainties in the Higgs boson production cross sections, including those related to the kinematic boundaries introduced in the STXS framework, are aligned between input analyses. All input analyses incorporate the STXS uncertainty scheme used in [49] for ggH and VBF production, the scheme used in [38] for ttH production, and the scheme used in [37] for VH production. These correspond to the most recent uncertainty schemes for each production process. For the tH and bbH production processes, which are not split in kinematic bins within the STXS framework, the inclusive theoretical uncertainties provided in [13] are used.

A pruning strategy is applied to reduce the complexity of the statistical model for some of the more memory-intensive input analyses, as follows. Starting from the smallest contribution in an analysis category, signal processes (split by production process and STXS bin) are removed in order of increasing yield, such that the sum of yields of the removed signal processes does not exceed 0.5% of the total signal yield in that category. The associated experimental uncertainties of a signal process are removed if the process contributes less than 1% of the total signal yield in its corresponding category. Additionally, the sum of yields of signal processes for which the experimental systematic uncertainties are removed must not exceed 5% of the total signal yield in the category. This pruning strategy has been developed to ensure negligible changes in the results for a significant decrease in the complexity of the model, and thus in the run time and memory usage of the fitting procedure. The best fit values in the individual channel STXS measurements change by less than 1% of their 68% confidence intervals in the majority of cases, with no changes greater than 3% of the 68% confidence intervals.

5. Statistical procedure

The statistical methodology developed by the ATLAS and CMS Collaborations, as described in [27], is employed in this paper. This section provides an overview of the methodology, further details on which can be found in [5, 74–76].

5.1. The combined likelihood

Each analysis is defined by a set of event categories that partition the data according to different selection criteria. In this paper, the event categories are referred to as analysis regions. Two types of analysis region exist: signal regions (SRs), which are constructed to be enriched in Higgs boson events, and CRs, which are constructed to constrain the background components in the SRs. The likelihood function for analysis region r is constructed for a data set $\mathcal{D}_r$ consisting of observations $x_d \in \mathcal{D}_r$, based on probability density functions (pdfs) $p(x_d;\vec{\alpha},\vec{\theta})$. The parameters $\vec{\alpha}$ and $\vec{\theta}$ are the POIs and nuisance parameters (NPs), respectively. For a binned analysis region, the data set consists of the observed event counts n_d in bin d of analysis region r. The likelihood function becomes

$$L_r\left(\mathcal{D}_r;\vec{\alpha},\vec{\theta}\right) = \prod_d \mathrm{Poisson}\left(n_d;\sum_{if} s_r^{if}\left(\vec{\alpha},\vec{\theta}\right)\rho_{\mathrm{sig},rd}^{if}\left(\vec{\alpha},\vec{\theta}\right) + b_r\left(\vec{\theta}\right)\rho_{\mathrm{bkg},rd}\left(\vec{\theta}\right)\right), \tag{1}$$

where the product runs over all bins d, $s_r^{if}(\vec{\alpha},\vec{\theta})$ is the expected signal yield in region r, for Higgs boson production process i and decay channel f, and the pdf $\rho_{\mathrm{sig},rd}^{if}(\vec{\alpha},\vec{\theta})$ represents the fraction of signal events that are contained in bin d. In principle, the function ρ_{sig} depends on both the POIs and the NPs, i.e. the model POIs can change the expected distribution of the signal across observable bins. For the combination, modifications to ρ_{sig} caused by the POIs are not considered, except for a short discussion on those induced by SMEFT operators (described in section 11.2). The signal contributions from each production process i, and decay channel f, in analysis region r are summed. For the most part, the analysis regions are defined to target the different Higgs boson decay channels separately to ensure orthogonality between input analyses. Therefore, the contributions in each region r are typically dominated by a single decay channel, f. The term $b_r(\vec{\theta})$ is the expected background yield in region r, and $\rho_{\mathrm{bkg},rd}(\vec{\theta})$ is the fraction of background events that are contained in bin d. The normalization and shape dependence of the background estimate is limited to the NPs. The background estimate consists of the sum over different processes, e.g. QCD multijet, top quark pair production, and V+jets.

The pdfs ρ_{sig} and ρ_{bkg} are constructed using histograms for template-based fits, e.g. in the $\text{H} \to \text{WW}$ [51] and $\text{H} \to \tau\tau$ [52] channels, and analytic functions for binned parametric fits, e.g. in the $\text{H} \to \gamma\gamma$ [49] and $\text{H} \to \text{Z}\gamma \to \ell\ell\gamma$ [55] inputs. The pdfs are estimated using simulation for the signal and both data and simulation for the background. For analyses performed using parametric models, the analytic functions are integrated over the bin dimensions to obtain the fraction of events expected in that bin.

For analyses based on unbinned fits, which only concerns the on-shell $\text{H} \to \text{ZZ} \to 4\ell$ channel [50] in the combination, the probability density term in the likelihood function can be written as

$$p(x;\vec{\alpha},\vec{\theta}) = \frac{1}{\sum_{if} s_r^{if}\left(\vec{\alpha},\vec{\theta}\right) + b_r\left(\vec{\theta}\right)} \left(\sum_{if} s_r^{if}\left(\vec{\alpha},\vec{\theta}\right) \rho_{\text{sig},r}^{if}\left(x;\vec{\alpha},\vec{\theta}\right) + b_r\left(\vec{\theta}\right) \rho_{\text{bkg},r}\left(x;\vec{\theta}\right) \right), \tag{2}$$

where $\rho_{\text{sig},r}^{if}(x;\vec{\alpha},\vec{\theta})$ is the signal pdf for Higgs boson production i and decay channel f in analysis region r, over observable x. The corresponding background pdfs for analysis region r are given by $\rho_{\text{bkg},r}(x;\vec{\alpha},\vec{\theta})$. For a data set with n entries, $x_d \in \mathcal{D}_r$, where d runs from 1 to n, a Poisson probability term is included in the definition of L_r,

$$L_r\left(\mathcal{D}_r;\vec{\alpha},\vec{\theta}\right) = \text{Poisson}\left(n; \sum_{if} s_r^{if}\left(\vec{\alpha},\vec{\theta}\right) + b_r\left(\vec{\theta}\right) \right) \prod_d p\left(x_d;\vec{\alpha},\vec{\theta}\right). \tag{3}$$

The combined likelihood function L is defined over the full data set $\mathcal{D}$ as the product of likelihoods over all mutually exclusive analysis regions L_r, multiplied by an additional constraint term for each NP,

$$L\left(\mathcal{D};\vec{\alpha},\vec{\theta}\right) = \prod_r L_r\left(\mathcal{D}_r;\vec{\alpha},\vec{\theta}\right) \prod_l p_l\left(y_l;\theta_l\right), \tag{4}$$

where each constrained NP $\theta_l \in \vec{\theta}$ is paired with a corresponding auxiliary observable $y_l \in \vec{y}$. The auxiliary observables are absorbed into the definition of the full data set $\mathcal{D}$ for notational simplicity, such that $\mathcal{D} = \{\mathcal{D}_r \forall r, \vec{y}\}$. The constraint term provides information about how well the NP is known *a priori*. The probability distributions for the auxiliary observables $p_l(y_l;\theta_l)$ can be normally, Poisson, or uniformly distributed, depending on the type of NP. In this likelihood function, all y_l are assumed to be statistically independent from each other and from the fitted observables. An individual NP represents a single source of systematic uncertainty, and its effect is considered fully correlated between all input analyses included in the fit.

5.2. Signal yield parametrization

For all input analyses, the expected signal yields $s_r^{if}(\vec{\alpha},\vec{\theta})$ and shapes $\rho_{\text{sig},r}^{if}(x;\vec{\alpha},\vec{\theta})$ are determined for a fixed m_{H} of 125.38 GeV. The numerous interpretations provided in this paper are obtained by fitting different parametrizations of the signal yield. These parametrizations are referred to as models. The expected signal yield $s_r^{if}(\vec{\alpha},\vec{\theta})$ from Higgs boson production process i and decay channel f, reconstructed in analysis region r, can be expressed as

$$s_r^{if}\left(\vec{\alpha},\vec{\theta}\right) = \mu^{if}(\vec{\alpha}) \left[\sigma^i \mathcal{B}^f\right]_{\text{SM,HO}}\left(\vec{\theta}_{\text{th,norm}}\right) \epsilon_{r,\text{SM}}^{if}\left(\vec{\theta}_{\text{th,acc}},\vec{\theta}_{\text{exp}}\right) \mathcal{L}\left(\vec{\theta}_{\text{lumi}}\right). \tag{5}$$

In this expression, $\mathcal{B}^f$ represents the Higgs boson branching fraction into decay channel f. The index i runs over STXS stage 1.2 bins for input analyses that implement the corresponding process splitting, and runs over the inclusive production process otherwise. The $\mu^{if}(\vec{\alpha})$ term in equation (5) describes the signal strength modifier for production process i and decay channel f. This encodes the effect of the POIs on the signal rate, and explains how the likelihood can be used to constrain the POIs from data. The term $\left[\sigma^i \mathcal{B}^f\right]_{\text{SM,HO}}(\vec{\theta}_{\text{th,norm}})$ represents the Higgs boson production cross section, in the SM, via production process i, multiplied by the SM Higgs boson branching fraction to final state f, evaluated at the highest-available orders (HO) in QCD and EW theory used in the combination, namely those specified for the inclusive cross sections in section 1, and for the relative STXS bin fractions in section 4. The theoretical uncertainties in the cross section and branching fraction estimates are encapsulated in the NPs $\vec{\theta}_{\text{th,norm}} \in \vec{\theta}$. The efficiency for production process i and decay channel f to be selected in analysis region r is encoded by $\epsilon_{r,\text{SM}}^{if}(\vec{\theta}_{\text{th,acc}},\vec{\theta}_{\text{exp}})$. These terms are estimated using samples of simulated Higgs boson events in the SM. The NPs affecting this term originate from both theoretical and experimental uncertainties, $\vec{\theta}_{\text{th,acc}},\vec{\theta}_{\text{exp}} \in \vec{\theta}$. The theoretical uncertainties, $\vec{\theta}_{\text{th,acc}}$, are defined such that the effect on the

total rate for process i and decay channel f is factored out. This means that variations in the NPs do not change the overall rate, but model the migration of events into (and out of) analysis region r. Finally, $\mathcal{L}(\vec{\theta}_{\text{lumi}})$ corresponds to the integrated luminosity estimate, with uncertainties encapsulated by the NPs $\vec{\theta}_{\text{lumi}}$.

Sections 7–11 provide further details on the different signal parametrization models. The full list of models is summarized in table 2, where the input analyses included in each fit are highlighted.

5.3. Extraction of results

The results are extracted using the CMS statistical analysis tool COMBINE [76], which is based on the data modelling and handling toolkits ROOFIT [77] and ROOSTATS [78].

The POIs for a particular model are estimated with their corresponding confidence intervals using a profile likelihood ratio test statistic $q(\mathcal{D};\vec{\alpha})$,

$$q(\mathcal{D};\vec{\alpha}) = -2\ln\left(\frac{L\left(\mathcal{D};\vec{\alpha},\hat{\hat{\vec{\theta}}}_{\vec{\alpha}}\right)}{L\left(\mathcal{D};\hat{\vec{\alpha}},\hat{\vec{\theta}}\right)}\right), \tag{6}$$

where the quantities $\hat{\vec{\alpha}}$ and $\hat{\vec{\theta}}$ denote the unconditional maximum likelihood estimates for the POIs and NPs, respectively. The quantity $\hat{\hat{\vec{\theta}}}_{\vec{\alpha}}$ corresponds to the conditional maximum likelihood estimate of $\vec{\theta}$ for fixed values of the POIs. The choice of POIs depends on the specific physics model under consideration. From this point onwards the explicit dependence on the data in the test statistic is omitted from the notation: $q(\vec{\alpha}) \equiv q(\mathcal{D};\vec{\alpha})$.

For each model considered, the maximum likelihood estimates $\hat{\vec{\alpha}}$ are identified as the best fit parameter values. The 68% and 95% confidence level (CL) intervals for one-dimensional (1D) measurements of each POI are determined under the asymptotic approximation as the union of intervals for which $q(\vec{\alpha}) < 1$ and $q(\vec{\alpha}) < 3.84$, respectively, unless otherwise stated. In models with more than one POI, these intervals are determined treating the other POIs as NPs. The differences between the boundaries of the 68% and 95% CL intervals and the best fit value yield the $\pm 68\%$ and $\pm 95\%$ uncertainties in the measurement. If the interval is restricted by a physical boundary that was imposed on the measured parameter, a truncated interval is reported and the uncertainty is derived from that interval; in such cases, the specified frequentist coverage is not expected to be preserved. Additionally, confidence intervals obtained for signal yield parametrizations with a nonlinear dependence on the POIs, e.g. the quadratic terms in the SMEFT interpretation described in section 11, may not maintain the specified frequentist coverage [79]. The two-dimensional (2D) 68% and 95% CL regions are determined from the set of parameter values for which $q(\vec{\alpha}) < 2.30$ and $q(\vec{\alpha}) < 5.99$, respectively.

The 1D CL intervals are often broken down into components according to the different sources of uncertainty (as shown in figure 3). Each component is calculated by performing a fit with the relevant NPs fixed to their best fit values, and re-deriving the CL intervals. The quadratic difference between the uncertainties is taken as the contribution to the total uncertainty from that component. This is performed for each component in turn, where the relevant NPs are sequentially fixed. The statistical component of the uncertainty is derived in the final step of this procedure, where all constrained NPs are fixed in the fit. Owing to correlations in the fit between NPs belonging to different uncertainty sources, the order in which the components are fixed to their best fit values can affect the contribution derived for each component. It should be stressed that this ordering dependence does not impact the total uncertainty, nor the size of the statistical component, but it can lead to small variations in the systematic uncertainty breakdown.

The likelihood functions are constructed with respect to either the observed data or an Asimov data set [80] to obtain the observed or expected results, respectively. The Asimov data set is constructed using the expected values of the POIs for the SM, and the NPs set to the values obtained in a fit to data with the POIs fixed at the SM values.

To compute the compatibility of results with the SM hypothesis for $m_{\text{H}} = 125.38\,\text{GeV}$, the test statistic is evaluated at the point in parameter space where all POIs take their values expected in the SM: $q(\vec{\alpha}_{\text{SM}})$. The compatibility is expressed as a p-value, p_{SM}, computed as

$$p_{\text{SM}} = 1 - F_{\chi^2_n}\left(q\left(\vec{\alpha}_{\text{SM}}\right)\right), \tag{7}$$

where $F_{\chi^2_n}$ is the cumulative function of the χ^2_n distribution with n equal to the number of POIs in the fitted model. This method also relies on the asymptotic assumption that $q(\vec{\alpha}_{\text{SM}})$ is distributed according to the χ^2_n distribution.

The correlations between fitted parameters are derived under the assumption of symmetric Gaussian uncertainties around the best fit parameter point, $\hat{\vec{\alpha}}$, by using the second derivatives of $q(\vec{\alpha})$. To extract the correlation coefficients, the second derivatives are determined numerically by stepping around the $q(\vec{\alpha})$ minimum.

6. Systematic uncertainties

For many of the POIs, the systematic uncertainties in their determination are expected to be comparable to, or larger than, the statistical uncertainties. Theoretical and experimental systematic uncertainties are incorporated as NPs in the fit, as described in section 5.1. The same sources of uncertainty in different analysis regions are modelled with common NPs, ensuring they are treated coherently for all measurements. Channel-specific uncertainties are also incorporated, and considered uncorrelated amongst channels. This section further details the different sources of systematic uncertainties that are common across channels.

6.1. Theoretical uncertainties

Theoretical uncertainties affecting the signal are among the dominant contributions to the systematic uncertainties. These arise from missing higher-order QCD corrections, PDF, α_S, underlying event and parton shower estimates, as well as the branching fraction predictions. The uncertainties affect the signal in two ways: normalization effects ($\vec{\theta}_{\text{th,norm}}$), which impact the expected signal yields via the cross section and branching fraction estimates, and acceptance effects ($\vec{\theta}_{\text{th,acc}}$), which model the migration of events between analysis regions (including in and out of experimental acceptance).

A common normalization uncertainty scheme is used across all channels that split the signal models according to the STXS stage 1.2 binning (as shown in table 1). Nuisance parameters are introduced to account for variations in the inclusive production cross sections, as well as the migration of events across the kinematic boundaries defined by the STXS framework. The ggH and qqH schemes follow the prescription described in [49], the VH scheme follows the prescription of [37], and the ttH scheme follows the prescription of [38]. The uncertainties in the tH and bbH production cross sections are defined inclusively, and are both individually correlated across channels [13]. For input analyses that only split the signal models into inclusive production components (as shown in table 1), the normalization uncertainties are correlated with the inclusive uncertainty components from the STXS stage 1.2 uncertainty scheme. Normalization uncertainties originating from the PDFs, the value of α_S, and the branching fraction predictions are correlated between channels, and match the recommendations in [13].

Acceptance uncertainties are derived by varying the relevant theoretical parameters, and calculating the change in the event rate for each analysis region, while factoring out the effect on the overall production cross section. This includes variations in the renormalization and factorization scales, the choice of PDF, the value of α_S, the modelling of initial-state and final-state radiation in the parton shower, and the underlying event tune. The acceptance uncertainty NPs for input analyses that adopt the same variation scheme are treated as correlated.

When measuring cross sections (sections 8.1 and 8.2), the $\vec{\theta}_{\text{th,norm}}$ NPs that affect the normalization of the measured POIs are ignored. For the signal strength measurements, as well as the interpretations in the coupling modifier and SMEFT frameworks, all $\vec{\theta}_{\text{th,norm}}$ are incorporated. The $\vec{\theta}_{\text{th,acc}}$ NPs are included in all fits.

For background processes that are normalized using theoretical predictions, the inclusive cross section uncertainties are correlated between channels in which the same background appears. These uncertainties are assumed to be uncorrelated with those affecting the signal estimates [74], with the exception of the uncertainties from the underlying event and parton shower model.

6.2. Experimental uncertainties

The different Higgs boson decay channels are usually sensitive to different experimental systematic uncertainties, resulting in the majority of them being uncorrelated. There are a few exceptions, which are discussed in the remainder of this section.

Uncertainties in the integrated luminosity measurement are considered year-by-year and the corresponding NPs ($\vec{\theta}_{\text{lumi}}$) are fully correlated across all input analyses. Values of 1.2, 2.3, and 2.5% are used for the 2016, 2017, and 2018 data sets, respectively [81–83]. A partial correlation scheme across the years is implemented to account for common sources of uncertainty in the luminosity measurements. The uncertainty in the modelling of additional collisions in the event (pileup) is considered correlated across all input analyses and years.

Table 2. A list of the different signal parametrization models provided in the combination, and the channels that are included in each fit. The second column indicates the number of POIs in the model. The last column indicates whether the model is included in this paper (no checkmark) or is provided as supplementary material (checkmark). The remaining columns show ticks for each channel entering the respective model fit. All input channels, except off-shell $H \to ZZ \to 4\ell$, are labelled as 'incl.' or 'STXS' according to the granularity of the signal processes. As described in section 3, the $H \to \tau\tau$, $H \to bb$ boosted, and ttH ($H \to bb$) inputs implement two sets of analysis regions.

Interpretation	n_{POI}	$H \to \gamma\gamma$, STXS	$H \to ZZ \to 4\ell$, STXS	$H \to WW$, STXS	$H \to \tau\tau$, STXS	$H \to \tau\tau$, incl.	$H \to bb$ boosted, STXS	$H \to bb$ boosted, incl.	VBF ($H \to bb$), incl.	VH ($H \to bb$), STXS	t(t)H ($H \to bb$), STXS	t(t)H ($H \to bb$), incl.	t(t)H ($H \to$ leptons), STXS	$H \to \mu\mu$, incl.	$H \to Z\gamma \to \ell\ell\gamma$, incl.	$H \to$ inv, incl.	Off-shell $H \to ZZ \to 4\ell$	Suppl. Mat.
Signal strength (inclusive)	1	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			
Signal strength (production)	6	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			
Signal strength (decay)	7	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			
Signal strength (production-times-decay)	31	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			
Cross section (STXS stage 0)	7	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			✓
Cross section (STXS stage 1.2)	32	✓	✓	✓	✓		✓			✓	✓		✓					✓
Cross section with branching fraction ratios (STXS stage 0)	13	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			
Cross section with branching fraction ratios (STXS stage 1.2)	36	✓	✓	✓	✓		✓			✓	✓		✓					
Signal strength (STXS stage 0)	7	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			✓
Signal strength (STXS stage 1.2)	32	✓	✓	✓	✓		✓			✓	✓		✓					✓
Signal strength with branching fraction ratios (STXS stage 0)	13	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			✓
Signal strength with branching fraction ratios (STXS stage 1.2)	36	✓	✓	✓	✓		✓			✓	✓		✓					✓
Signal strength with STXS times branching fraction	97	✓	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓			
Resolved coupling modifiers	6	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			
Effective coupling modifiers ($\mathcal{B}_{BSM} = 0$)	9	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			
Effective coupling modifiers ($\mathcal{B}_{inv.}$, $\mathcal{B}_{undet.}$ floating)	11	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓	✓		
Off-shell coupling model ($\mathcal{B}_{inv.}$, $\mathcal{B}_{undet.}$ floating)	11	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓	✓	✓	
Ratios of coupling modifiers	8	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			
Ratios of coupling modifiers, symmetry of fermion couplings	3	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			
Higgs self-coupling, κ_λ (κ_V and κ_F fixed)	1	✓	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓			
Higgs self-coupling, κ_λ (κ_V and κ_F floating)	3	✓	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓			
Higgs self-coupling, κ_λ-vs-κ_V (2D)	2	✓	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓			
Higgs self-coupling, κ_λ-vs-κ_F (2D)	2	✓	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓			
UV-complete extensions to SM (2HDM and hMSSM)	2	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓	✓			
SMEFT, one-POI-at-a-time, linear	1 (×43)	✓	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓			
SMEFT, one-POI-at-a-time, lin+quad	1 (×43)	✓	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓			
SMEFT, multiple POIs (PCA), linear	17	✓	✓	✓	✓		✓		✓	✓	✓		✓	✓	✓			

The uncertainties affecting the lepton efficiencies and energy scales are derived using tag-and-probe techniques on the Z boson resonance. Their uncertainty is dominated by the extraction techniques and therefore is correlated per flavour among the input channels and years, when the same NP scheme is applied. The jet energy scale and jet energy resolution uncertainties reflect the data taking conditions and calibration of the subdetectors of the CMS experiment. Dedicated jet energy scale and jet energy resolution NP schemes are introduced to account for the partial correlations between uncertainty sources and data-taking years. Uncertainties in the b tagging efficiency measurements are correlated across channels that implement the same NP scheme. The VH ($H \to bb$) channel uses a larger set of finely binned NPs, which are uncorrelated from the other channels. The uncertainty in the efficiency of the double-b tagger algorithm is taken to be uncorrelated from the single-b tagging efficiency estimates. Uncertainties in p_T^{miss} are taken as correlated between analyses.

Uncertainties arising from the finite size of simulated event samples, referred to as MC statistical uncertainties, are treated using the Barlow-Beeston lite approach [84]. In this method, an NP is introduced per bin of each histogram-based template derived from simulation, and these NPs are considered uncorrelated across all input analyses.

Several analyses determine the normalization, or both the normalization and shape parameters, of the background model by a fit to data. This includes the parameters that vary the choice of the background parametrization in the $H \to \gamma\gamma$, $H \to \mu\mu$, and $H \to Z\gamma \to \ell\ell\gamma$ analysis regions. These parameters are not associated with any additional constraint terms, and are therefore assigned to the statistical uncertainty component of a measurement.

7. Measurements of signal strengths

Signal strength measurements are defined by the POIs $\vec{\alpha} = \vec{\mu}$ such that

$$\mu^{if}(\vec{\alpha}) = \mu^{if}. \tag{8}$$

The parameters μ^{if} scale the rate of Higgs boson production process i and decay channel f relative to the SM expectation. In other words, for $\mu^{if} = 1$ the signal rate is equal to the SM expectation. The signal strengths are determined at different levels of granularity:

- *Inclusive*, μ^{incl}: introduces a single POI that scales all Higgs boson signal processes equally.
- *Per production process*, μ^{i}: introduces a separate signal strength parameter for each of the major Higgs boson production processes: ggH, VBF, WH, ZH, ttH, and tH. In this parametrization, as well as all subsequent parametrizations involving signal strengths or cross sections, the ZH production process includes contributions from both the gluon-initiated (ggZH) and quark-initiated (qqZH) processes, the tH production process includes contributions from both tHq and tHW production, and the bbH production process is scaled with the same parameter as ggH.
- *Per decay channel*, μ^{f}: introduces a separate signal strength parameter for each of the Higgs boson decay channels included in the combination. Seven decay channels are simultaneously fit: $H \to \gamma\gamma$, $H \to ZZ$, $H \to WW$, $H \to bb$, $H \to \tau\tau$, $H \to \mu\mu$, and $H \to Z\gamma$.
- *Per production and decay channel*, μ^{if}: introduces a separate signal strength parameter for each pair of Higgs boson production process and decay channel that the combination is sensitive to. A total of 31 signal strength parameters are fit simultaneously in this case.

The measured value of the inclusive signal strength modifier is

$$\begin{aligned} \mu^{incl} &= 1.014^{+0.055}_{-0.053} \\ &= 1.014^{+0.040}_{-0.039}\,(\text{theo})^{+0.025}_{-0.024}\,(\text{exp}) \pm 0.028\,(\text{stat}), \end{aligned} \tag{9}$$

where the total uncertainty has been decomposed, in order, into the theoretical, experimental, and statistical components.

A more granular breakdown of the uncertainties in the inclusive signal strength measurement is provided in figure 3. The largest component of the uncertainty originates from the theoretical uncertainty in the signal yield normalization ($\Delta\mu^{incl}/\mu^{incl} = 3.6\%$). The contributions from the experimental uncertainties are shared amongst the different sources of uncertainty, with no single dominant contribution.



Figure 3. Breakdown of the 68% CL intervals on the best fit inclusive (left), per production process (center), and per decay channel (right) signal strength modifiers, for the different sources of uncertainty. All contributions are symmetrized by taking the average of the upward and downward fluctuations. The top panel shows the uncertainty contributions as percentages relative to the best fit signal strength values. The total uncertainty is shown by the black boxes, while the combined systematic uncertainties from theoretical and experimental components are shown in purple and orange, respectively, and the statistical uncertainty is shown in blue. The middle and bottom panels show the breakdown of the theoretical and experimental systematic uncertainties into their different contributions, respectively. The NPs from each uncertainty source are sequentially fixed to their best fit values to derive the individual contributions following the order in which they are shown in the figure. The fractional contribution is then evaluated by dividing each individual contribution by the linear sum of all contributions in that panel. The 'Misc.' label absorbs all other experimental uncertainty contributions not listed explicitly in the figure, including those related to a specific trigger timing issue in the forward region of the ECAL during 2016–2017 data taking [43]. This particular uncertainty has a larger impact on the $\mu^{\tau\tau}$ parameter, whose sensitivity is dominated by VBF production which relies on forward-jet reconstruction.

The inclusive signal strength measurement shows good compatibility with the SM, with a p-value of $p_{SM} = 0.80$.

The per production process and per decay channel signal strength measurements are shown in figures 4 and 5, respectively. The upper plots in the figures show the best fit points with the corresponding 68% and 95% CL intervals, where the 68% CL intervals are decomposed into their theoretical, experimental, and statistical components. The numerical values for the best fit points and 68% CL intervals, and the corresponding expected intervals calculated using an Asimov data set, are provided in table 3. As with the inclusive signal strength, the uncertainties in the μ^i and μ^f parameters are broken down into the different uncertainty sources in figure 3.

In contrast to the inclusive measurement, the per production process measurement shows a small tension with the SM, with a compatibility p-value of $p_{SM} = 0.02$. This tension is mostly driven by μ^{tH}, for which an excess of 2.2 standard deviations above the SM expectation is seen. The μ^{WH} and μ^{ZH} parameters are also measured to be larger than the SM expectations by approximately two standard deviations. The 68% CL intervals range from ±7.5% for μ^{ggH} to ±39% for μ^{tH}, relative to their best fit values.



Figure 4. The measured inclusive (μ^{incl}) and per production process (μ^{i}) signal strength modifiers. In the upper plot, the thick (thin) black lines indicate the 68% (95%) CL intervals, with the theoretical systematic, experimental systematic, and statistical components of the 68% intervals indicated by the purple, orange, and blue bands, respectively. The grey band shows the 68% CL interval on the inclusive signal strength modifier. A separate axis is provided for the μ^{tH} parameter because of its larger uncertainty and high best fit value. The correlations between the 6 parameters in the per production process measurement are shown in the lower plot. The size of the correlations is indicated by the colour scale.

The per decay channel measurement shows a better compatibility with the SM ($p_{SM} = 0.33$). The largest deviations are observed in the $\mu^{\tau\tau}$ and $\mu^{Z\gamma}$ parameters. However, these are still compatible with the SM expectations within the 95% CL intervals. The $\mu^{\gamma\gamma}$, μ^{ZZ}, μ^{WW}, and $\mu^{\tau\tau}$ parameters are measured with 68% CL intervals of approximately ±10% relative to their best fit values. The μ^{bb} parameter is measured with a 68% CL interval of ±15%. This represents a significant improvement compared to the previous combined Higgs boson measurement by the CMS Collaboration (±21%) [32], because of the newly added H → bb channels and updated H → bb input analyses. The parameters for the rarer decay channels, $\mu^{\mu\mu}$ and $\mu^{Z\gamma}$, are measured with 68% CL intervals of ±37% and ±39%, respectively, relative to their best fit values.





Figure 5. The measured per decay channel signal strength modifiers, μ^f. In the upper plot, the thick (thin) black lines indicate the 68% (95%) CL intervals, with the theoretical systematic, experimental systematic, and statistical components of the 68% intervals indicated by the purple, orange, and blue bands, respectively. The grey band shows the 68% CL interval on the inclusive signal strength modifier. A separate axis is provided for the $\mu^{Z\gamma}$ parameter because of its larger uncertainty and high best fit value. The correlations between the 7 parameters considered in this fit are shown in the lower plot. The size of the correlations is indicated by the colour scale.

The lower plots in figures 4 and 5 show the correlations between the signal strengths for the per production process and per decay channel measurements, respectively. Overall, the correlations are small. The largest (anti)correlation in the per production process measurement is observed between the μ^{ttH} and μ^{tH} parameters (-32%), reflecting the experimental difficulty in distinguishing these processes as a result of their similar final states. The per decay channel signal strengths exhibit predominantly positive correlations, particularly between the $\mu^{\gamma\gamma}$, μ^{ZZ}, and μ^{WW} parameters. These correlations arise from the dominant contribution of ggH production to these decay channels and the resulting shared dependence on theoretical NPs affecting ggH production.

Table 3. Best fit values and 68% CL intervals for the per production process and per decay channel signal strengths. The total 68% CL intervals are decomposed into their statistical and systematic components. The expected intervals are given in parentheses.

Per production process, μ^i				Per decay channel, μ^f			
		Uncertainty				Uncertainty	
Parameter	Best fit	Stat	Syst	Parameter	Best fit	Stat	Syst
μ^{ggH}	$0.99^{+0.08}_{-0.07}$ $\left(^{+0.08}_{-0.07}\right)$	$^{+0.04}_{-0.04}$ $\left(^{+0.04}_{-0.04}\right)$	$^{+0.07}_{-0.06}$ $\left(^{+0.07}_{-0.06}\right)$	$\mu^{\gamma\gamma}$	$1.12^{+0.09}_{-0.09}$ $\left(^{+0.08}_{-0.08}\right)$	$^{+0.06}_{-0.06}$ $\left(^{+0.06}_{-0.06}\right)$	$^{+0.07}_{-0.06}$ $\left(^{+0.06}_{-0.05}\right)$
μ^{VBF}	$0.87^{+0.12}_{-0.11}$ $\left(^{+0.12}_{-0.11}\right)$	$^{+0.10}_{-0.10}$ $\left(^{+0.10}_{-0.10}\right)$	$^{+0.06}_{-0.06}$ $\left(^{+0.07}_{-0.06}\right)$	μ^{ZZ}	$1.03^{+0.11}_{-0.10}$ $\left(^{+0.11}_{-0.10}\right)$	$^{+0.08}_{-0.08}$ $\left(^{+0.08}_{-0.07}\right)$	$^{+0.08}_{-0.07}$ $\left(^{+0.07}_{-0.06}\right)$
μ^{WH}	$1.47^{+0.24}_{-0.24}$ $\left(^{+0.22}_{-0.21}\right)$	$^{+0.18}_{-0.18}$ $\left(^{+0.17}_{-0.17}\right)$	$^{+0.16}_{-0.15}$ $\left(^{+0.14}_{-0.13}\right)$	μ^{WW}	$0.99^{+0.09}_{-0.09}$ $\left(^{+0.09}_{-0.09}\right)$	$^{+0.05}_{-0.05}$ $\left(^{+0.05}_{-0.05}\right)$	$^{+0.08}_{-0.07}$ $\left(^{+0.08}_{-0.07}\right)$
μ^{ZH}	$1.35^{+0.23}_{-0.21}$ $\left(^{+0.20}_{-0.19}\right)$	$^{+0.17}_{-0.17}$ $\left(^{+0.16}_{-0.15}\right)$	$^{+0.16}_{-0.13}$ $\left(^{+0.13}_{-0.12}\right)$	$\mu^{\tau\tau}$	$0.84^{+0.10}_{-0.10}$ $\left(^{+0.11}_{-0.11}\right)$	$^{+0.06}_{-0.06}$ $\left(^{+0.06}_{-0.06}\right)$	$^{+0.09}_{-0.08}$ $\left(^{+0.10}_{-0.09}\right)$
μ^{ttH}	$0.77^{+0.18}_{-0.15}$ $\left(^{+0.20}_{-0.18}\right)$	$^{+0.12}_{-0.12}$ $\left(^{+0.12}_{-0.12}\right)$	$^{+0.13}_{-0.10}$ $\left(^{+0.15}_{-0.13}\right)$	μ^{bb}	$1.02^{+0.16}_{-0.15}$ $\left(^{+0.15}_{-0.14}\right)$	$^{+0.10}_{-0.10}$ $\left(^{+0.10}_{-0.10}\right)$	$^{+0.12}_{-0.11}$ $\left(^{+0.11}_{-0.10}\right)$
μ^{tH}	$5.87^{+2.38}_{-2.22}$ $\left(^{+2.14}_{-2.01}\right)$	$^{+1.88}_{-1.82}$ $\left(^{+1.77}_{-1.69}\right)$	$^{+1.45}_{-1.28}$ $\left(^{+1.20}_{-1.08}\right)$	$\mu^{\mu\mu}$	$1.15^{+0.44}_{-0.41}$ $\left(^{+0.44}_{-0.41}\right)$	$^{+0.42}_{-0.39}$ $\left(^{+0.40}_{-0.39}\right)$	$^{+0.15}_{-0.13}$ $\left(^{+0.18}_{-0.13}\right)$
				$\mu^{Z\gamma}$	$2.45^{+1.00}_{-0.93}$ $\left(^{+0.85}_{-0.85}\right)$	$^{+0.93}_{-0.90}$ $\left(^{+0.84}_{-0.83}\right)$	$^{+0.37}_{-0.24}$ $\left(^{+0.15}_{-0.18}\right)$

Figure 6 shows the results for the signal strength parametrization with a separate POI for each production process and decay channel combination, μ^{if}. In this fit, the tH production process is assumed to scale like ttH, such that both processes are fit with a common parameter, μ^{ttH+tH}. Given the five production processes and seven decay channels considered, this implies a model with 35 POIs. However, not all can be experimentally constrained in the combination. The WH and ZH production processes in the $H \to \mu\mu$ channel are fit together with a common signal strength $\mu^{VH,\mu\mu}$. Additionally, the WH, ZH, and ttH+tH production processes are constrained to the SM prediction for the $H \to Z\gamma$ channel. In the cases of VBF, WH, ZH, and ttH+tH production with the $H \to ZZ$ decay, the background contamination is sufficiently low that a negative signal strength can result in an overall negative event yield. Therefore, these signal strengths are restricted to nonnegative values, as indicated by the grey hatched boxes in figure 6.

The best fit values and 68% CL intervals decomposed into their systematic and statistical parts, as well as the corresponding expected intervals, are provided in table 4. The precision on a number of these measurements has been improved compared to the previous combined Higgs boson measurement by the CMS Collaboration [32]. Furthermore, several signal strength parameters are becoming dominated by systematic uncertainties, particularly those related to ggH production. This measurement shows a small tension with the SM hypothesis ($p_{SM} = 0.03$), which is driven by the $\mu^{ggH,\tau\tau}$, $\mu^{WH,WW}$, and $\mu^{ttH+tH,bb}$ parameters, as well as the signal strengths in the $H \to Z\gamma$ decay channel.

The correlations between the measured μ^{if} are shown in figure 7. Sizeable correlations are observed for parameters describing the same decay channel (blocks along the diagonal), whereas the correlations between different decay channels are typically small. The ggH parameters provide an exception to this pattern, where residual positive correlations are again induced by the common theoretical NPs affecting ggH production.

8. Measurements of simplified template cross sections and decay branching fractions

8.1. Stage 0 measurements

This section details the measurements of the production process cross sections and decay branching fractions. The cross section measurements follow the stage 0 definition of the STXS framework [13], which introduces a fiducial region defined by the rapidity of the Higgs boson, $|y_H| < 2.5$. All input analyses have a negligible acceptance for $|y_H| > 2.5$. By defining a fiducial region in this way, the theoretical



Figure 6. Summary of the signal strength modifier measurements. The empty circles and black lines represent the best fit points and 68% CL intervals, respectively. The orange lines indicate the systematic components of the total 68% CL intervals. The left-hand panel shows the measurements of the per decay channel signal strengths, while the grey band shows the 68% CL interval on the inclusive signal strength modifier. The other panels show the measurements of the per production and decay channel signal strength modifiers, μ^{if}. Within these panels, the blue bands indicate the 68% CL intervals on the per production process signal strength modifiers. The SM predicted values are indicated by the red lines. The central values and 68% CL intervals for the μ^{if} parameters are explicitly written in each panel of the plot. The H → ZZ signal strengths are constrained to be nonnegative as indicated by the grey hatched boxes. Additionally, the WH, ZH, and ttH+tH production processes for the H → Zγ channel are constrained to the SM predictions, a feature that is indicated by the black hatched boxes.

uncertainty due to extrapolating to the inclusive Higgs boson production phase space is minimized. The cross sections that are measured are:

- σ^{ggH}: ggH production. This parameter includes the contribution from bbH production. While [13] proposes to separate these two processes, they are merged because no analysis regions targeting H production in association with a b quark are included in the combination. This parameter also incorporates ggZH production with the Z boson decaying hadronically.
- σ^{VBF}: VBF production.
- $\sigma^{V(qq)H}$: quark-initiated Higgs boson production in association with a W or Z boson, where the vector boson decays hadronically.
- $\sigma^{W(\ell\nu)H}$: WH production, where the W boson decays leptonically.
- $\sigma^{Z(\ell\ell,\nu\nu)H}$: ZH production, where the Z boson decays leptonically. As the combination is not sensitive to the quark-initiated (qqZH) and gluon-initiated (ggZH) processes separately, they are merged.
- σ^{ttH}: associated production with a pair of top quarks.
- σ^{tH}: associated production with a single top quark. This includes the contributions from both tHq and tHW production.

The cross section parameters are extracted as products of the cross sections and the H → ZZ branching fraction, $\sigma^{i}\mathcal{B}^{ZZ}$, where H → ZZ is used as the reference decay as it is one of the channels with the smallest overall and systematic uncertainties. In addition, this choice of reference maintains consistency with previous combined measurements from the CMS and ATLAS Collaborations [27, 28, 31].

Table 4. Best fit values and 68% CL intervals for the per production and decay channel signal strengths. The total 68% CL intervals are decomposed into their statistical and systematic components. The expected intervals are given in parentheses. Some of the signal strengths are restricted to nonnegative values, as described in the text. Truncated intervals are reported for these parameters if the 68% CL interval is not fully contained in the positive domain.

	Production process														
	ggH			VBF			WH			ZH			ttH + tH		
Decay channel		Uncertainty			Uncertainty			Uncertainty			Uncertainty			Uncertainty	
	Best fit	Stat	Syst	Best fit	Stat	Syst	Best fit	Stat	Syst	Best fit	Stat	Syst	Best fit	Stat	Syst
$H \to \gamma\gamma$	$1.06^{+0.12}_{-0.11}$ $\left(^{+0.11}_{-0.10}\right)$	$^{+0.09}_{-0.09}$ $\left(^{+0.08}_{-0.08}\right)$	$^{+0.08}_{-0.07}$ $\left(^{+0.08}_{-0.07}\right)$	$1.05^{+0.34}_{-0.34}$ $\left(^{+0.33}_{-0.31}\right)$	$^{+0.30}_{-0.31}$ $\left(^{+0.30}_{-0.29}\right)$	$^{+0.16}_{-0.13}$ $\left(^{+0.14}_{-0.11}\right)$	$1.45^{+0.54}_{-0.49}$ $\left(^{+0.55}_{-0.44}\right)$	$^{+0.53}_{-0.48}$ $\left(^{+0.54}_{-0.44}\right)$	$^{+0.11}_{-0.10}$ $\left(^{+0.08}_{-0.04}\right)$	$1.20^{+0.69}_{-0.61}$ $\left(^{+0.68}_{-0.59}\right)$	$^{+0.69}_{-0.61}$ $\left(^{+0.68}_{-0.59}\right)$	$^{+0.00}_{-0.05}$ $\left(^{+0.08}_{-0.04}\right)$	$1.35^{+0.34}_{-0.29}$ $\left(^{+0.29}_{-0.25}\right)$	$^{+0.27}_{-0.26}$ $\left(^{+0.25}_{-0.23}\right)$	$^{+0.21}_{-0.13}$ $\left(^{+0.15}_{-0.10}\right)$
$H \to ZZ$	$1.03^{+0.13}_{-0.12}$ $\left(^{+0.13}_{-0.12}\right)$	$^{+0.10}_{-0.10}$ $\left(^{+0.10}_{-0.09}\right)$	$^{+0.09}_{-0.07}$ $\left(^{+0.08}_{-0.07}\right)$	$0.38^{+0.52}_{-0.38}$ $\left(^{+0.56}_{-0.47}\right)$	$^{+0.48}_{-0.38}$ $\left(^{+0.54}_{-0.45}\right)$	$^{+0.20}_{-0.05}$ $\left(^{+0.15}_{-0.11}\right)$	$0.00^{+2.64}_{-0.00}$ $\left(^{+1.79}_{-1.00}\right)$	$^{+1.91}_{-0.00}$ $\left(^{+1.76}_{-1.00}\right)$	$^{+1.82}_{-0.00}$ $\left(^{+0.33}_{-0.00}\right)$	$7.31^{+4.66}_{-6.12}$ $\left(^{+3.99}_{-1.00}\right)$	$^{+3.64}_{-4.42}$ $\left(^{+3.81}_{-1.00}\right)$	$^{+2.92}_{-4.23}$ $\left(^{+1.18}_{-0.00}\right)$	$0.00^{+0.89}_{-0.00}$ $\left(^{+1.43}_{-0.84}\right)$	$^{+0.85}_{-0.00}$ $\left(^{+1.38}_{-0.83}\right)$	$^{+0.25}_{-0.00}$ $\left(^{+0.40}_{-0.07}\right)$
$H \to WW$	$0.92^{+0.11}_{-0.11}$ $\left(^{+0.11}_{-0.11}\right)$	$^{+0.06}_{-0.05}$ $\left(^{+0.06}_{-0.06}\right)$	$^{+0.10}_{-0.09}$ $\left(^{+0.10}_{-0.09}\right)$	$0.82^{+0.24}_{-0.23}$ $\left(^{+0.24}_{-0.23}\right)$	$^{+0.20}_{-0.19}$ $\left(^{+0.21}_{-0.20}\right)$	$^{+0.13}_{-0.12}$ $\left(^{+0.13}_{-0.12}\right)$	$2.38^{+0.73}_{-0.69}$ $\left(^{+0.58}_{-0.55}\right)$	$^{+0.52}_{-0.50}$ $\left(^{+0.46}_{-0.44}\right)$	$^{+0.51}_{-0.47}$ $\left(^{+0.35}_{-0.32}\right)$	$1.88^{+0.74}_{-0.65}$ $\left(^{+0.63}_{-0.54}\right)$	$^{+0.66}_{-0.60}$ $\left(^{+0.58}_{-0.51}\right)$	$^{+0.35}_{-0.25}$ $\left(^{+0.25}_{-0.18}\right)$	$1.45^{+0.38}_{-0.33}$ $\left(^{+0.35}_{-0.32}\right)$	$^{+0.29}_{-0.29}$ $\left(^{+0.30}_{-0.29}\right)$	$^{+0.24}_{-0.15}$ $\left(^{+0.18}_{-0.14}\right)$
$H \to \tau\tau$	$0.61^{+0.21}_{-0.20}$ $\left(^{+0.28}_{-0.25}\right)$	$^{+0.09}_{-0.09}$ $\left(^{+0.09}_{-0.09}\right)$	$^{+0.19}_{-0.18}$ $\left(^{+0.26}_{-0.23}\right)$	$0.86^{+0.17}_{-0.16}$ $\left(^{+0.17}_{-0.17}\right)$	$^{+0.14}_{-0.13}$ $\left(^{+0.14}_{-0.14}\right)$	$^{+0.10}_{-0.09}$ $\left(^{+0.10}_{-0.09}\right)$	$1.40^{+0.59}_{-0.57}$ $\left(^{+0.57}_{-0.54}\right)$	$^{+0.50}_{-0.47}$ $\left(^{+0.48}_{-0.46}\right)$	$^{+0.32}_{-0.31}$ $\left(^{+0.30}_{-0.27}\right)$	$1.88^{+0.66}_{-0.57}$ $\left(^{+0.55}_{-0.48}\right)$	$^{+0.55}_{-0.51}$ $\left(^{+0.50}_{-0.45}\right)$	$^{+0.37}_{-0.26}$ $\left(^{+0.24}_{-0.18}\right)$	$0.33^{+0.42}_{-0.39}$ $\left(^{+0.49}_{-0.43}\right)$	$^{+0.29}_{-0.27}$ $\left(^{+0.32}_{-0.31}\right)$	$^{+0.31}_{-0.29}$ $\left(^{+0.38}_{-0.30}\right)$
$H \to bb$	$2.22^{+1.53}_{-1.39}$ $\left(^{+1.27}_{-1.18}\right)$	$^{+1.18}_{-1.17}$ $\left(^{+1.13}_{-1.12}\right)$	$^{+0.98}_{-0.75}$ $\left(^{+0.59}_{-0.38}\right)$	$1.65^{+0.76}_{-0.64}$ $\left(^{+0.50}_{-0.46}\right)$	$^{+0.48}_{-0.48}$ $\left(^{+0.44}_{-0.43}\right)$	$^{+0.59}_{-0.42}$ $\left(^{+0.25}_{-0.14}\right)$	$1.28^{+0.37}_{-0.36}$ $\left(^{+0.34}_{-0.32}\right)$	$^{+0.25}_{-0.24}$ $\left(^{+0.23}_{-0.23}\right)$	$^{+0.28}_{-0.26}$ $\left(^{+0.24}_{-0.22}\right)$	$1.11^{+0.27}_{-0.26}$ $\left(^{+0.26}_{-0.24}\right)$	$^{+0.20}_{-0.19}$ $\left(^{+0.19}_{-0.18}\right)$	$^{+0.18}_{-0.17}$ $\left(^{+0.18}_{-0.16}\right)$	$0.39^{+0.27}_{-0.26}$ $\left(^{+0.30}_{-0.28}\right)$	$^{+0.17}_{-0.17}$ $\left(^{+0.18}_{-0.18}\right)$	$^{+0.21}_{-0.20}$ $\left(^{+0.24}_{-0.21}\right)$
$H \to \mu\mu$	$0.40^{+0.67}_{-0.85}$ $\left(^{+0.75}_{-0.72}\right)$	$^{+0.65}_{-0.84}$ $\left(^{+0.73}_{-0.72}\right)$	$^{+0.14}_{-0.10}$ $\left(^{+0.17}_{-0.07}\right)$	$1.45^{+0.88}_{-0.71}$ $\left(^{+0.78}_{-0.67}\right)$	$^{+0.73}_{-0.67}$ $\left(^{+0.70}_{-0.64}\right)$	$^{+0.48}_{-0.24}$ $\left(^{+0.34}_{-0.22}\right)$		5.58	$^{+3.35}_{-3.08}$ $\left(^{+2.69}_{-2.38}\right)$	$^{+3.23}_{-2.96}$ $\left(^{+2.68}_{-2.37}\right)$	$^{+0.91}_{-0.86}$ $\left(^{+0.27}_{-0.14}\right)$		$2.93^{+2.50}_{-2.17}$ $\left(^{+2.22}_{-1.85}\right)$	$^{+2.42}_{-2.12}$ $\left(^{+2.19}_{-1.84}\right)$	$^{+0.65}_{-0.45}$ $\left(^{+0.37}_{-0.12}\right)$
$H \to Z\gamma$	$3.61^{+1.42}_{-1.21}$ $\left(^{+1.11}_{-1.12}\right)$	$^{+1.24}_{-1.18}$ $\left(^{+1.10}_{-1.11}\right)$	$^{+0.69}_{-0.25}$ $\left(^{+0.11}_{-0.19}\right)$	$-3.36^{+3.10}_{-3.06}$ $\left(^{+3.26}_{-2.97}\right)$	$^{+3.03}_{-2.86}$ $\left(^{+3.14}_{-2.96}\right)$	$^{+0.64}_{-1.07}$ $\left(^{+0.88}_{-0.27}\right)$		— —			— —			— —	



Figure 7. The correlations between the 31 parameters considered in the per production and decay channel fit. The size of the correlations is indicated by the colour scale. Correlations with an absolute magnitude smaller than 0.005 are not shown.

Modifications in the other Higgs boson branching fractions are incorporated by also floating the ratios of branching fractions with respect to $\mathcal{B}^{ZZ}$. In total this defines 13 POIs, with seven of the form

$$\mu^{i,ZZ} = \frac{\left[\sigma^i \mathcal{B}^{ZZ}\right]_{\text{obs}}}{\left[\sigma^i \mathcal{B}^{ZZ}\right]_{\text{SM,HO}}\left(\vec{\theta}'_{\text{th,norm}}\right)}. \tag{10}$$

The other six correspond to the ratios of branching fractions,

$$R^{f/ZZ} = \frac{\mathcal{B}^f_{\text{obs}}/\mathcal{B}^f_{\text{SM,HO}}\left(\vec{\theta}'_{\text{th,norm}}\right)}{\mathcal{B}^{ZZ}_{\text{obs}}/\mathcal{B}^{ZZ}_{\text{SM,HO}}\left(\vec{\theta}'_{\text{th,norm}}\right)}, \tag{11}$$

such that the signal contributions in the $H \to ZZ$ decay channel are scaled by $\mu^{i,ZZ}$, while signal contributions in the other decay channels are scaled by $\mu^{i,ZZ}R^{f/ZZ}$.

This measurement differs from the signal strength measurement by the treatment of the subset of theoretical NPs $\vec{\theta}'_{\text{th,norm}}$ that affect the normalization of the measured cross sections and branching fractions. These terms are included in the denominators of the POI definitions and therefore cancel the corresponding terms in the signal yield parametrization, equation (5). Consequently, these theoretical uncertainties do not enter the cross section measurements, and are instead attributed to the uncertainties in the SM predictions of the measured quantities. It should be noted that all theoretical uncertainties that account for the migration of events between STXS stage 1.2 bins are still included, as they do not impact the inclusive production process cross sections. In addition, the theoretical uncertainties affecting the signal acceptance in each analysis region are still included in the fit. The observed cross sections $\left[\sigma^i \mathcal{B}^{ZZ}\right]_{\text{obs}}$ are obtained through the multiplication of the fitted POIs by the predictions from the highest-available order SM calculation.



Figure 8. The measured STXS stage 0 cross sections and branching fraction ratios. Theoretical uncertainties affecting the normalizations of the measured parameters are not included in the fit. In the upper plot, the empty circles indicate the best fit values and the vertical lines with caps indicate the 68% CL intervals. The lower 68% CL interval for the $\sigma^{V(qq)H}\mathcal{B}^{ZZ}$ parameter lies outside of the plotted range, as indicated by the arrow. The wider boxes show the systematic uncertainty components of the 68% CL intervals. Each measured quantity is compared with the SM prediction in red, where the grey bands indicate the theoretical uncertainty in the respective prediction. The lower plot shows the correlations between the 13 parameters considered in the fit. The size of the correlations is indicated by the colour scale.

The best fit points and 68% CL intervals for the cross sections and ratios of branching fractions are shown in the upper plot of figure 8. Each measured quantity is compared to the SM prediction and its corresponding uncertainty, as shown by the red lines and grey bands, respectively. The numerical values

of the best fit points and 68% CL intervals, as well as the expected intervals calculated using an Asimov data set under the SM hypothesis, are provided in table 5. The table also contains the SM predictions of the fit parameters and the corresponding theoretical uncertainties in these predictions.

This measurement shows a similar tension with the SM as the per production process signal strength fit in figure 4. The tension is driven by the excess in the tH production process measurement. Additionally, the measurements of $\sigma^{\mathrm{W}(\ell\nu)\mathrm{H}}\mathcal{B}^{\mathrm{ZZ}}$ and $\sigma^{\mathrm{Z}(\ell\ell,\nu\nu)\mathrm{H}}\mathcal{B}^{\mathrm{ZZ}}$ mirror the observed deviations in the signal strengths μ^{WH} and μ^{ZH}. The ratios of branching fractions also present some deviations from the SM predictions, most notably for the $R^{\mathrm{bb/ZZ}}$ and $R^{\mathrm{Z}\gamma/\mathrm{ZZ}}$ parameters. As a result, the overall compatibility is measured to be $p_{\mathrm{SM}} = 0.012$. Note that this result does not incorporate the theoretical uncertainties affecting the normalizations of the measured parameters, and this can lead to an underestimation of p_{SM}.

The correlations between the fit parameters are displayed in the lower plot of figure 8. In this parametrization, sizeable correlations arise because the branching fractions are expressed as ratios to $\mathcal{B}^{\mathrm{ZZ}}$, introducing a common denominator that directly couples the decay channels. As a result, scaling the signal yields in the other decay channels by the product $\mu^{i,\mathrm{ZZ}}R^{f/\mathrm{ZZ}}$ induces anticorrelations between $\mu^{i,\mathrm{ZZ}}$ and the ratios $R^{f/\mathrm{ZZ}}$, which in turn leads to positive correlations among the ratios themselves.

The results from a measurement in which the SM Higgs boson branching fractions are assumed can be found in the supplementary material to this paper [41]. This measurement considers only the cross sections as POIs, and incorporates the theoretical uncertainties associated with the Higgs boson branching fractions. Furthermore, the measurements detailed in this section are also performed in a signal strength formalism, where all theoretical uncertainties are folded into the measurement. As a result, the confidence intervals in the POIs are slightly larger compared with the measurements in the cross section formalism, and the p_{SM} values increase. These results are also provided as supplementary material [41], and can be used for reinterpretation without the need to define an external theoretical uncertainty scheme.

8.2. Stage 1.2 measurements

This section describes cross section measurements in kinematic regions defined by the STXS framework [13]. The bins are defined according to the STXS stage 1.2 binning scheme [22]. Only the input analyses that split the signal models according to the STXS stage 1.2 granularity (as shown in table 1) are included in this measurement.

As with the production cross section measurements described in section 8.1, bbH production and ggZH production in which the Z boson decays hadronically are grouped with ggH production. The ZH leptonic bins include contributions from both qqZH and ggZH production, and the tH bin includes both tHq and tHW production. Furthermore, the qqH bins include contributions from both VBF production and quark-initiated VH production where the vector boson decays hadronically. All STXS cross sections are defined in the fiducial region of $|y_{\mathrm{H}}| < 2.5$.

The Higgs boson production processes are partitioned according to the kinematic properties of the Higgs boson, and if present, the kinematic properties of associated jets and vector bosons that are independent of the Higgs boson decay. Here, jets are built using the anti-k_{T} algorithm [85, 86] with a jet distance parameter of 0.4, from all stable particles with a lifetime greater than 10 ps, excluding particles originating from the Higgs boson decay and leptons from vector boson decays. All jets are required to have a transverse momentum $p_{\mathrm{T}}^{\mathrm{jet}} > 30\,\mathrm{GeV}$.

In the STXS stage 1.2 binning scheme, ggH production is partitioned according to the jet multiplicity (0, 1, or $\geqslant$2 jets), and the transverse momentum of the Higgs boson $p_{\mathrm{T}}^{\mathrm{H}}$. Events with $\geqslant$2 jets are further split according to the invariant mass of the dijet system m_{jj}, and the transverse momentum of the Higgs boson plus dijet system $p_{\mathrm{T}}^{\mathrm{Hjj}}$, where the dijet system is formed from the two highest-p_{T} jets in the event. Similarly, qqH production is partitioned firstly by the jet multiplicity, and events with $\geqslant$2 jets are further split according to $p_{\mathrm{T}}^{\mathrm{H}}$, m_{jj}, and $p_{\mathrm{T}}^{\mathrm{Hjj}}$. For WH and ZH production, in which the vector boson decays leptonically, the bins are defined by the transverse momentum of the vector boson $p_{\mathrm{T}}^{\mathrm{V}}$, with an additional partition for events with $p_{\mathrm{T}}^{\mathrm{V}}$ between 150 and 250 GeV according to the jet multiplicity. The ttH production process is partitioned into five $p_{\mathrm{T}}^{\mathrm{H}}$ bins, and the tH production process is measured inclusively. In the figures and tables that present the results, STXS bins with N jets are labelled 'NJ'.

Several bins from the nominal STXS stage 1.2 binning scheme must be merged to avoid either very large uncertainties in the measured cross sections, or very strong correlations between them. A criterion based on the expected correlations between measured cross sections is used to determine the merging, such that absolute correlations remain below 0.75. In total, 32 STXS regions are fit simultaneously, and these are shown by the filled boxes in figure 9. The nominal STXS stage 1.2 bin boundaries that are merged in the measurement are indicated by the dashed lines in the figure. Of the 32 measured cross sections, 13 are associated with ggH production, 5 with qqH production, 4 each with WH and

Table 5. Best fit values and 68% CL intervals for the parameters in the fit of production process cross sections and branching fraction ratios. The cross sections are defined in the fiducial region $|y_H| < 2.5$. The values are normalized to the SM predictions. The total 68% CL intervals are decomposed into their statistical and systematic components, and the expected intervals are given in parentheses. The SM predictions for each of the measured parameters are also provided, along with the corresponding theoretical uncertainties in the predictions. The $\sigma^{ggH}\mathcal{B}^{ZZ}$ prediction includes contributions from bbH production and ggZH production with the Z boson decaying to hadrons, while $\sigma^{Z(\ell\ell,\nu\nu)H}\mathcal{B}^{ZZ}$ includes contributions from ggZH production with the Z boson decaying to leptons.

Parameters	SM prediction ($m_H = 125.38$ GeV)	Best fit/SM pred.	Stat	Syst
$\sigma^{ggH}\mathcal{B}^{ZZ}$	1221^{+78}_{-78} fb	$1.02^{+0.10}_{-0.09}\,(^{+0.10}_{-0.09})$	$^{+0.08}_{-0.08}\,(^{+0.08}_{-0.08})$	$^{+0.06}_{-0.05}\,(^{+0.05}_{-0.05})$
$\sigma^{VBF}\mathcal{B}^{ZZ}$	$95.6^{+2.4}_{-2.4}$ fb	$0.94^{+0.18}_{-0.16}\,(^{+0.17}_{-0.15})$	$^{+0.14}_{-0.13}\,(^{+0.14}_{-0.13})$	$^{+0.10}_{-0.08}\,(^{+0.11}_{-0.09})$
$\sigma^{V(qq)H}\mathcal{B}^{ZZ}$	$34.4^{+0.8}_{-0.8}$ fb	$0.70^{+0.74}_{-0.75}\,(^{+0.73}_{-0.68})$	$^{+0.63}_{-0.63}\,(^{+0.61}_{-0.58})$	$^{+0.38}_{-0.40}\,(^{+0.41}_{-0.35})$
$\sigma^{W(\ell\nu)H}\mathcal{B}^{ZZ}$	$10.6^{+0.2}_{-0.3}$ fb	$1.86^{+0.42}_{-0.36}\,(^{+0.28}_{-0.24})$	$^{+0.33}_{-0.30}\,(^{+0.22}_{-0.20})$	$^{+0.26}_{-0.20}\,(^{+0.16}_{-0.13})$
$\sigma^{Z(\ell\ell,\nu\nu)H}\mathcal{B}^{ZZ}$	$6.45^{+0.28}_{-0.23}$ fb	$1.88^{+0.47}_{-0.40}\,(^{+0.29}_{-0.24})$	$^{+0.38}_{-0.33}\,(^{+0.23}_{-0.20})$	$^{+0.28}_{-0.22}\,(^{+0.17}_{-0.13})$
$\sigma^{ttH}\mathcal{B}^{ZZ}$	$13.5^{+0.9}_{-1.3}$ fb	$0.95^{+0.22}_{-0.20}\,(^{+0.20}_{-0.18})$	$^{+0.18}_{-0.17}\,(^{+0.17}_{-0.15})$	$^{+0.13}_{-0.11}\,(^{+0.12}_{-0.10})$
$\sigma^{tH}\mathcal{B}^{ZZ}$	$2.44^{+0.16}_{-0.32}$ fb	$5.44^{+2.64}_{-2.42}\,(^{+2.20}_{-2.07})$	$^{+2.19}_{-2.05}\,(^{+1.86}_{-1.75})$	$^{+1.49}_{-1.28}\,(^{+1.17}_{-1.10})$
$\mathcal{B}^{bb}/\mathcal{B}^{ZZ}$	$21.2^{+0.4}_{-0.4}$	$0.66^{+0.17}_{-0.14}\,(^{+0.24}_{-0.19})$	$^{+0.13}_{-0.11}\,(^{+0.19}_{-0.16})$	$^{+0.11}_{-0.09}\,(^{+0.15}_{-0.11})$
$\mathcal{B}^{WW}/\mathcal{B}^{ZZ}$	$8.11^{+0.15}_{-0.15}$	$0.95^{+0.12}_{-0.11}\,(^{+0.12}_{-0.11})$	$^{+0.09}_{-0.08}\,(^{+0.09}_{-0.08})$	$^{+0.08}_{-0.07}\,(^{+0.08}_{-0.07})$
$\mathcal{B}^{\tau\tau}/\mathcal{B}^{ZZ}$	$2.29^{+0.04}_{-0.04}$	$0.80^{+0.14}_{-0.12}\,(^{+0.17}_{-0.14})$	$^{+0.09}_{-0.08}\,(^{+0.11}_{-0.10})$	$^{+0.10}_{-0.09}\,(^{+0.13}_{-0.11})$
$\mathcal{B}^{\gamma\gamma}/\mathcal{B}^{ZZ}$	$0.084^{+0.002}_{-0.002}$	$1.02^{+0.12}_{-0.11}\,(^{+0.12}_{-0.11})$	$^{+0.10}_{-0.10}\,(^{+0.10}_{-0.09})$	$^{+0.06}_{-0.05}\,(^{+0.06}_{-0.05})$
$\mathcal{B}^{Z\gamma}/\mathcal{B}^{ZZ}$	$0.058^{+0.003}_{-0.003}$	$2.41^{+1.12}_{-0.87}\,(^{+0.86}_{-0.84})$	$^{+1.09}_{-0.83}\,(^{+0.85}_{-0.83})$	$^{+0.25}_{-0.25}\,(^{+0.10}_{-0.10})$
$\mathcal{B}^{\mu\mu}/\mathcal{B}^{ZZ}$	$0.0079^{+0.0001}_{-0.0001}$	$1.19^{+0.45}_{-0.40}\,(^{+0.45}_{-0.40})$	$^{+0.43}_{-0.38}\,(^{+0.42}_{-0.38})$	$^{+0.15}_{-0.12}\,(^{+0.16}_{-0.12})$



Figure 9. A diagram showing the set of 32 STXS regions that are considered. The filled boxes represent the measured STXS regions, while the dashed lines indicate the nominal STXS stage 1.2 bin boundaries that are merged in the measurement. The units of p_T^H, m_{jj}, p_T^{Hjj}, and p_T^V are in GeV.

ZH production with the vector boson decaying leptonically, 5 with ttH production, and one with tH production. The measured STXS bins, along with their SM cross section predictions, are summarized in table 6. The table also highlights which nominal STXS stage 1.2 bins are merged in the measurement. Units of GeV are assumed for all numerical values in the STXS bin naming scheme.

Following the procedure described in section 8.1, the cross sections are measured as products with the $H \to ZZ$ branching fraction, $\sigma^i\mathcal{B}^{ZZ}$. The ratios of branching fractions, $R^{f/ZZ}$, are again included as additional parameters to account for modifications in the Higgs boson branching fractions. As this measurement only combines the input analyses with the STXS stage 1.2 process granularity, the ratios for $H \to \mu\mu$ or $H \to Z\gamma$ are not considered.

The theoretical uncertainties that affect the normalization of the fitted cross sections and branching fractions are not included in the measurement. As with the production process cross section measurement, these uncertainties are instead attributed to the SM predictions. The STXS theoretical uncertainty scheme, discussed in section 6, includes NPs that account for the migration of events across the STXS

Table 6. A summary of the STXS regions measured in the combination. All STXS regions are defined within the fiducial region $|y_H| < 2.5$. Some of the measured regions are defined by merging several neighbouring bins (in parentheses) from the nominal STXS stage 1.2 binning scheme to ensure sufficient sensitivity. The SM predictions, with theoretical uncertainties, are also provided for the measured STXS regions. Units of GeV are assumed for all numerical values related to the p_T^H, m_{jj}, p_T^{Hjj}, and p_T^V variables. The ggH (ZH lep) bins include contributions from ggZH production with the Z boson decaying hadronically (leptonically). The 'VBF-topo' or 'VH-topo' naming convention refers to events consistent with the topology of VBF or VH production, characterized by two jets with large m_{jj} (for VBF), or two jets with m_{jj} compatible with a vector boson decay (for VH).

Measured regions	STXS stage 1.2 bins (number of merged bins)	σ^i_{SM} [pb]
ggH 0J, $p_T^H < 10$	ggH 0J, $p_T^H < 10$	$6.70^{+0.82}_{-0.82}$
ggH 0J, $p_T^H > 10$ + bbH	ggH 0J, $10 < p_T^H < 200$ + bbH (2)	$19.5^{+1.4}_{-1.4}$
ggH 1J, $p_T^H < 60$	ggH 1J, $p_T^H < 60$	$7.14^{+0.95}_{-0.95}$
ggH 1J, $60 < p_T^H < 120$	ggH 1J, $60 < p_T^H < 120$	$4.95^{+0.68}_{-0.68}$
ggH 1J, $120 < p_T^H < 200$	ggH 1J, $120 < p_T^H < 200$	$0.88^{+0.15}_{-0.15}$
ggH $\geqslant$2J, $0 < m_{jj} < 350$, $p_T^H < 60$	ggH $\geqslant$2J, $0 < m_{jj} < 350$, $p_T^H < 60$	$1.24^{+0.29}_{-0.29}$
ggH $\geqslant$2J, $0 < m_{jj} < 350$, $60 < p_T^H < 120$	ggH $\geqslant$2J, $0 < m_{jj} < 350$, $60 < p_T^H < 120$	$2.00^{+0.46}_{-0.46}$
ggH $\geqslant$2J, $0 < m_{jj} < 350$, $120 < p_T^H < 200$	ggH $\geqslant$2J, $0 < m_{jj} < 350$, $120 < p_T^H < 200$	$0.93^{+0.22}_{-0.22}$
ggH VBF-topo	ggH $\geqslant$2J, $350 < m_{jj} < 700$, $p_T^{Hjj} < 25$, $p_T^H < 200$; ggH $\geqslant$2J, $350 < m_{jj} < 700$, $p_T^{Hjj} > 25$, $p_T^H < 200$; ggH $\geqslant$2J, $m_{jj} > 700$, $p_T^{Hjj} < 25$, $p_T^H < 200$; ggH $\geqslant$2J, $m_{jj} > 700$, $p_T^{Hjj} > 25$, $p_T^H < 200$ (4)	$0.98^{+0.22}_{-0.22}$
ggH $200 < p_T^H < 300$	ggH $200 < p_T^H < 300$	$0.49^{+0.12}_{-0.12}$
ggH $300 < p_T^H < 450$	ggH $300 < p_T^H < 450$	$0.12^{+0.03}_{-0.03}$
ggH $450 < p_T^H < 650$	ggH $450 < p_T^H < 650$	$0.015^{+0.004}_{-0.004}$
ggH $p_T^H > 650$	ggH $p_T^H > 650$	$0.0022^{+0.0005}_{-0.0005}$
qqH other	qqH 0J; qqH 1J; qqH $\geqslant$2J, $m_{jj} < 60$; qqH $\geqslant$2J, $120 < m_{jj} < 350$ (4)	$2.78^{+0.05}_{-0.05}$
qqH $60 < m_{jj} < 120$ (VH-topo)	qqH $\geqslant$2J, $60 < m_{jj} < 120$	$0.54^{+0.01}_{-0.01}$
qqH $350 < m_{jj} < 700$	qqH $\geqslant$2J, $350 < m_{jj} < 700$, $p_T^{Hjj} < 25$, $p_T^H < 200$; qqH $\geqslant$2J, $350 < m_{jj} < 700$, $p_T^{Hjj} > 25$, $p_T^H < 200$ (2)	$0.57^{+0.03}_{-0.03}$
qqH $m_{jj} > 700$	qqH $\geqslant$2J, $m_{jj} > 700$, $p_T^{Hjj} < 25$, $p_T^H < 200$; qqH $\geqslant$2J, $m_{jj} > 700$, $p_T^{Hjj} > 25$, $p_T^H < 200$ (2)	$0.74^{+0.02}_{-0.02}$
qqH $p_T^H > 200$	qqH $\geqslant$2J, $m_{jj} > 350$, $p_T^H > 200$	$0.16^{+0.004}_{-0.004}$
WH lep $p_T^V < 75$	WH lep $p_T^V < 75$	$0.21^{+0.01}_{-0.01}$
WH lep $75 < p_T^V < 150$	WH lep $75 < p_T^V < 150$	$0.13^{+0.01}_{-0.01}$
WH lep $150 < p_T^V < 250$	WH lep $150 < p_T^V < 250$, 0J; WH lep $150 < p_T^V < 250$, $\geqslant$1J (2)	$0.040^{+0.002}_{-0.002}$
WH lep $p_T^V > 250$	WH lep $p_T^V > 250$	$0.013^{+0.001}_{-0.001}$
ZH lep $p_T^V < 150$	ZH lep $p_T^V < 75$; ZH lep $75 < p_T^V < 150$; (2)	$0.20^{+0.01}_{-0.01}$
ZH lep $150 < p_T^V < 250$, 0J	ZH $150 < p_T^V < 250$, 0J	$0.015^{+0.002}_{-0.002}$
ZH lep $150 < p_T^V < 250$, $\geqslant$1J	ZH lep $150 < p_T^V < 250$, $\geqslant$ 1J	$0.017^{+0.003}_{-0.003}$
ZH lep $p_T^V > 250$	ZH lep $p_T^V > 250$	$0.0099^{+0.0010}_{-0.0009}$
ttH $p_T^H < 60$	ttH $p_T^H < 60$	$0.11^{+0.03}_{-0.03}$
ttH $60 < p_T^H < 120$	ttH $60 < p_T^H < 120$	$0.17^{+0.02}_{-0.03}$
ttH $120 < p_T^H < 200$	ttH $120 < p_T^H < 200$	$0.13^{+0.02}_{-0.02}$
ttH $200 < p_T^H < 300$	ttH $200 < p_T^H < 300$	$0.054^{+0.008}_{-0.009}$
ttH $p_T^H > 300$	ttH $p_T^H > 300$	$0.027^{+0.004}_{-0.005}$
tH	tHq; tHW (2)	$0.090^{+0.006}_{-0.012}$

bins. These NPs are removed from the fit unless they account for a migration of events across a kinematic boundary that is merged in the measurement, indicated by the dashed lines in figure 9. The theoretical uncertainties that affect the signal acceptance in each analysis region are still included in the fit.

The best fit values and 68% CL intervals for the fitted parameters are shown in figure 10. The observed cross section values shown in the upper panel are obtained by multiplying $\mu^{i,ZZ}$ by the SM prediction at the highest available order. The SM predictions and their associated uncertainties are shown by the red lines and grey bands, respectively. The corresponding numerical values, along with the expected 68% CL intervals, are provided in table 7. The total uncertainties are decomposed into their systematic and statistical parts, highlighting that the statistical uncertainty dominates for all parameters at this level of kinematic splitting.

The results show reasonable sensitivity to many different regions of the Higgs boson production phase space. The 68% CL intervals range from approximately ±14% of the SM $\sigma^i\mathcal{B}^{ZZ}$ prediction for



Figure 10. The measured STXS stage 1.2 cross sections and branching fraction ratios. Theoretical uncertainties that affect the normalizations of the measured parameters are not included in the fit. In the upper panel, the points indicate the best fit values while the coloured lines indicate the 68% CL intervals. The wider boxes show the systematic uncertainty components of the 68% CL intervals. Each measured quantity is compared to the SM prediction in red, where the grey bands indicate the theoretical uncertainty for the respective parameter. The lower panel plots the ratio of the measured $\sigma^i\mathcal{B}^{ZZ}$ with respect to the SM predictions. The best fit values for the ggH $450 < p_T^H < 650$ GeV and tH STXS bins lie outside of the range of the ratio panel, and are thus represented by arrows. Arrows are also used to indicate 68% CL intervals that extend beyond the plotted range.

the ggH 0J, $10 < p_T^H < 200$ GeV STXS bin, to approximately ±250% for the ggH $p_T^H > 650$ GeV STXS bin. For a number of measurements, e.g. ggH $200 < p_T^H < 300$ GeV, the precision is comparable to the theoretical uncertainty in the SM prediction, meaning the possibility of constraining theoretical inputs using experimental measurements of the Higgs boson is approaching.

There are several noticeable deviations from the SM in these results. Of particular interest are the WH leptonic $p_T^V > 250$ GeV and ZH leptonic $p_T^V > 250$ GeV regions, which are measured to be 2.2 and 2.1 standard deviations above the SM predictions, respectively. These regions of the VH production phase space at high p_T^V are particularly sensitive to BSM physics, as shown by the SMEFT parametrization discussed in section 11. The ZH leptonic $p_T^V < 150$ GeV region also exceeds the SM prediction by approximately 2.5 standard deviations. Small tensions with the SM are observed in the ggH measurements. The ggH 0J, $p_T^H < 10$ GeV STXS bin is measured to be approximately two standard deviations below the SM prediction, whereas the measurements in the remaining ggH STXS bins generally exceed their respective predictions. The largest deviation is observed in the ggH $450 < p_T^H < 650$ GeV STXS bin, which exceeds the SM prediction by more than 2.2 standard deviations. As with the previous signal yield parametrizations, the tH production process is found to have an excess of ∼2 standard deviations from the SM prediction. The overall compatibility with the SM is found to be $p_{SM} = 0.06$.

The correlations between the measured parameters are shown in figure 11. The overall structure follows the pattern observed in the STXS stage 0 measurement, shown in figure 8. As in that case, expressing the branching fractions as ratios to $\mathcal{B}^{ZZ}$ introduces a common denominator that couples the decay

Table 7. Best fit values and 68% CL intervals for the parameters in the fit to stage 1.2 simplified template cross sections and branching fraction ratios. The cross sections are defined in the fiducial region $|y_H| < 2.5$. The values are normalized to the SM predictions. The total 68% CL intervals are decomposed into their statistical and systematic components, and the expected intervals are given in parentheses. The SM prediction, with its theoretical uncertainty, for each of the measured parameters is also provided.

Parameters	SM prediction ($m_H = 125.38$ GeV)	Best fit/SM pred.	Stat	Syst
ggH 0J $p_T^H < 10$ x $\mathcal{B}^{ZZ}$	182^{+22}_{-22} fb	$0.67^{+0.17}_{-0.15}\,(^{+0.19}_{-0.17})$	$^{+0.15}_{-0.14}\,(^{+0.17}_{-0.16})$	$^{+0.07}_{-0.06}\,(^{+0.08}_{-0.07})$
ggH 0J $p_T^H > 10$ + bbH x $\mathcal{B}^{ZZ}$	529^{+39}_{-39} fb	$1.15^{+0.14}_{-0.13}\,(^{+0.13}_{-0.12})$	$^{+0.12}_{-0.11}\,(^{+0.11}_{-0.10})$	$^{+0.07}_{-0.06}\,(^{+0.07}_{-0.06})$
ggH 1J $p_T^H < 60$ x $\mathcal{B}^{ZZ}$	194^{+26}_{-26} fb	$1.15^{+0.28}_{-0.26}\,(^{+0.25}_{-0.24})$	$^{+0.23}_{-0.22}\,(^{+0.22}_{-0.21})$	$^{+0.16}_{-0.14}\,(^{+0.13}_{-0.12})$
ggH 1J $60 < p_T^H < 120$ x $\mathcal{B}^{ZZ}$	134^{+18}_{-18} fb	$1.19^{+0.28}_{-0.27}\,(^{+0.26}_{-0.24})$	$^{+0.23}_{-0.23}\,(^{+0.23}_{-0.22})$	$^{+0.16}_{-0.15}\,(^{+0.11}_{-0.10})$
ggH 1J $120 < p_T^H < 200$ x $\mathcal{B}^{ZZ}$	$24.0^{+4.0}_{-4.0}$ fb	$1.44^{+0.42}_{-0.39}\,(^{+0.37}_{-0.35})$	$^{+0.35}_{-0.34}\,(^{+0.34}_{-0.32})$	$^{+0.23}_{-0.21}\,(^{+0.16}_{-0.14})$
ggH $\geqslant$2 J $0 < m_{jj} < 350$, $p_T^H < 60$ x $\mathcal{B}^{ZZ}$	$33.7^{+7.9}_{-7.9}$ fb	$2.15^{+0.91}_{-0.89}\,(^{+0.76}_{-0.72})$	$^{+0.78}_{-0.75}\,(^{+0.68}_{-0.66})$	$^{+0.47}_{-0.46}\,(^{+0.33}_{-0.30})$
ggH $\geqslant$2 J $0 < m_{jj} < 350$, $60 < p_T^H < 120$ x $\mathcal{B}^{ZZ}$	$54.3^{+12.5}_{-12.5}$ fb	$1.62^{+0.49}_{-0.46}\,(^{+0.44}_{-0.42})$	$^{+0.43}_{-0.41}\,(^{+0.40}_{-0.39})$	$^{+0.24}_{-0.21}\,(^{+0.17}_{-0.16})$
ggH $\geqslant$2 J $0 < m_{jj} < 350$, $120 < p_T^H < 200$ x $\mathcal{B}^{ZZ}$	$25.4^{+5.9}_{-5.9}$ fb	$1.00^{+0.42}_{-0.39}\,(^{+0.42}_{-0.39})$	$^{+0.36}_{-0.35}\,(^{+0.38}_{-0.36})$	$^{+0.22}_{-0.19}\,(^{+0.17}_{-0.15})$
ggH VBF-topo x $\mathcal{B}^{ZZ}$	$26.5^{+6.1}_{-6.1}$ fb	$1.49^{+0.82}_{-0.80}\,(^{+0.75}_{-0.72})$	$^{+0.74}_{-0.72}\,(^{+0.69}_{-0.67})$	$^{+0.36}_{-0.35}\,(^{+0.29}_{-0.26})$
ggH $200 < p_T^H < 300$ x $\mathcal{B}^{ZZ}$	$13.2^{+3.3}_{-3.3}$ fb	$1.39^{+0.41}_{-0.39}\,(^{+0.37}_{-0.35})$	$^{+0.35}_{-0.34}\,(^{+0.34}_{-0.32})$	$^{+0.21}_{-0.19}\,(^{+0.15}_{-0.14})$
ggH $300 < p_T^H < 450$ x $\mathcal{B}^{ZZ}$	$3.37^{+0.85}_{-0.85}$ fb	$0.88^{+0.57}_{-0.53}\,(^{+0.55}_{-0.52})$	$^{+0.51}_{-0.48}\,(^{+0.50}_{-0.48})$	$^{+0.24}_{-0.22}\,(^{+0.22}_{-0.19})$
ggH $450 < p_T^H < 650$ x $\mathcal{B}^{ZZ}$	$0.42^{+0.11}_{-0.11}$ fb	$5.29^{+2.25}_{-1.93}\,(^{+1.38}_{-1.22})$	$^{+2.03}_{-1.81}\,(^{+1.27}_{-1.17})$	$^{+0.97}_{-0.68}\,(^{+0.52}_{-0.34})$
ggH $p_T^H > 650$ x $\mathcal{B}^{ZZ}$	$0.059^{+0.015}_{-0.015}$ fb	$4.44^{+4.85}_{-4.27}\,(^{+2.58}_{-2.23})$	$^{+4.26}_{-3.83}\,(^{+2.35}_{-2.15})$	$^{+2.32}_{-1.88}\,(^{+1.05}_{-0.62})$
qqH other x $\mathcal{B}^{ZZ}$	$75.6^{+1.7}_{-1.7}$ fb	$-1.10^{+1.53}_{-1.45}\,(^{+1.37}_{-1.20})$	$^{+1.21}_{-1.11}\,(^{+1.24}_{-1.20})$	$^{+0.94}_{-0.93}\,(^{+0.58}_{-0.12})$
qqH $350 < m_{jj} < 700$ x $\mathcal{B}^{ZZ}$	$15.4^{+0.8}_{-0.8}$ fb	$1.20^{+0.77}_{-0.73}\,(^{+0.70}_{-0.67})$	$^{+0.69}_{-0.65}\,(^{+0.64}_{-0.61})$	$^{+0.35}_{-0.34}\,(^{+0.27}_{-0.28})$
qqH $m_{jj} > 700$ x $\mathcal{B}^{ZZ}$	$20.0^{+0.6}_{-0.6}$ fb	$0.90^{+0.27}_{-0.25}\,(^{+0.27}_{-0.24})$	$^{+0.24}_{-0.22}\,(^{+0.24}_{-0.22})$	$^{+0.13}_{-0.11}\,(^{+0.12}_{-0.10})$
qqH $60 < m_{jj} < 120$ (VH-topo) x $\mathcal{B}^{ZZ}$	$14.6^{+0.4}_{-0.4}$ fb	$0.55^{+0.72}_{-0.75}\,(^{+0.71}_{-0.66})$	$^{+0.66}_{-0.70}\,(^{+0.67}_{-0.63})$	$^{+0.28}_{-0.27}\,(^{+0.23}_{-0.20})$
qqH $p_T^H > 200$ x $\mathcal{B}^{ZZ}$	$4.41^{+0.12}_{-0.12}$ fb	$0.65^{+0.30}_{-0.27}\,(^{+0.32}_{-0.28})$	$^{+0.26}_{-0.24}\,(^{+0.28}_{-0.25})$	$^{+0.15}_{-0.12}\,(^{+0.15}_{-0.12})$
WH lep $p_T^V < 75$ x $\mathcal{B}^{ZZ}$	$5.60^{+0.24}_{-0.24}$ fb	$2.02^{+1.10}_{-0.91}\,(^{+0.88}_{-0.75})$	$^{+0.97}_{-0.83}\,(^{+0.81}_{-0.72})$	$^{+0.53}_{-0.36}\,(^{+0.36}_{-0.23})$
WH lep $75 < p_T^V < 150$ x $\mathcal{B}^{ZZ}$	$3.52^{+0.16}_{-0.16}$ fb	$0.87^{+0.96}_{-0.88}\,(^{+0.91}_{-0.75})$	$^{+0.90}_{-0.85}\,(^{+0.86}_{-0.72})$	$^{+0.32}_{-0.25}\,(^{+0.30}_{-0.18})$
WH lep $150 < p_T^V < 250$ x $\mathcal{B}^{ZZ}$	$1.09^{+0.05}_{-0.05}$ fb	$1.14^{+0.89}_{-0.83}\,(^{+0.60}_{-0.52})$	$^{+0.70}_{-0.64}\,(^{+0.48}_{-0.41})$	$^{+0.54}_{-0.53}\,(^{+0.37}_{-0.32})$
WH lep $p_T^V > 250$ x $\mathcal{B}^{ZZ}$	$0.35^{+0.02}_{-0.02}$ fb	$3.40^{+1.57}_{-1.10}\,(^{+0.57}_{-0.43})$	$^{+1.14}_{-0.87}\,(^{+0.42}_{-0.34})$	$^{+1.08}_{-0.67}\,(^{+0.38}_{-0.26})$
ZH lep $p_T^V < 150$ x $\mathcal{B}^{ZZ}$	$5.30^{+0.20}_{-0.17}$ fb	$2.34^{+0.63}_{-0.54}\,(^{+0.43}_{-0.38})$	$^{+0.53}_{-0.48}\,(^{+0.36}_{-0.33})$	$^{+0.34}_{-0.26}\,(^{+0.24}_{-0.20})$
ZH lep $150 < p_T^V < 250$, 0J x $\mathcal{B}^{ZZ}$	$0.41^{+0.06}_{-0.06}$ fb	$0.70^{+0.94}_{-0.79}\,(^{+0.61}_{-0.48})$	$^{+0.71}_{-0.63}\,(^{+0.48}_{-0.39})$	$^{+0.62}_{-0.48}\,(^{+0.38}_{-0.27})$
ZH lep $150 < p_T^V < 250$, $\geqslant$ 1J x $\mathcal{B}^{ZZ}$	$0.47^{+0.08}_{-0.08}$ fb	$-0.25^{+1.45}_{-1.76}\,(^{+0.93}_{-0.83})$	$^{+1.19}_{-1.27}\,(^{+0.74}_{-0.67})$	$^{+0.82}_{-1.22}\,(^{+0.57}_{-0.49})$
ZH lep $p_T^V > 250$ x $\mathcal{B}^{ZZ}$	$0.27^{+0.03}_{-0.02}$ fb	$3.07^{+1.51}_{-0.99}\,(^{+0.50}_{-0.36})$	$^{+1.04}_{-0.79}\,(^{+0.38}_{-0.30})$	$^{+1.10}_{-0.61}\,(^{+0.32}_{-0.20})$
ttH $p_T^H < 60$ x $\mathcal{B}^{ZZ}$	$3.08^{+0.92}_{-0.94}$ fb	$0.64^{+0.76}_{-0.64}\,(^{+0.70}_{-0.61})$	$^{+0.72}_{-0.62}\,(^{+0.64}_{-0.57})$	$^{+0.26}_{-0.16}\,(^{+0.28}_{-0.22})$
ttH $60 < p_T^H < 120$ x $\mathcal{B}^{ZZ}$	$4.75^{+0.63}_{-0.72}$ fb	$1.23^{+0.54}_{-0.47}\,(^{+0.48}_{-0.43})$	$^{+0.50}_{-0.45}\,(^{+0.45}_{-0.41})$	$^{+0.18}_{-0.13}\,(^{+0.18}_{-0.14})$
ttH $120 < p_T^H < 200$ x $\mathcal{B}^{ZZ}$	$3.51^{+0.49}_{-0.55}$ fb	$0.99^{+0.49}_{-0.43}\,(^{+0.43}_{-0.38})$	$^{+0.46}_{-0.41}\,(^{+0.40}_{-0.36})$	$^{+0.17}_{-0.13}\,(^{+0.17}_{-0.14})$
ttH $200 < p_T^H < 300$ x $\mathcal{B}^{ZZ}$	$1.47^{+0.23}_{-0.25}$ fb	$1.07^{+0.68}_{-0.60}\,(^{+0.55}_{-0.48})$	$^{+0.61}_{-0.55}\,(^{+0.49}_{-0.43})$	$^{+0.29}_{-0.23}\,(^{+0.26}_{-0.21})$
ttH $p_T^H > 300$ x $\mathcal{B}^{ZZ}$	$0.73^{+0.11}_{-0.12}$ fb	$0.04^{+0.90}_{-0.85}\,(^{+0.75}_{-0.65})$	$^{+0.78}_{-0.71}\,(^{+0.64}_{-0.57})$	$^{+0.44}_{-0.47}\,(^{+0.40}_{-0.32})$
tH x $\mathcal{B}^{ZZ}$	$2.44^{+0.16}_{-0.32}$ fb	$6.17^{+2.90}_{-2.70}\,(^{+2.39}_{-2.18})$	$^{+2.29}_{-2.17}\,(^{+1.96}_{-1.89})$	$^{+1.77}_{-1.61}\,(^{+1.38}_{-1.09})$
$\mathcal{B}^{bb}/\mathcal{B}^{ZZ}$	$21.2^{+0.4}_{-0.4}$	$0.56^{+0.20}_{-0.16}\,(^{+0.33}_{-0.26})$	$^{+0.15}_{-0.12}\,(^{+0.25}_{-0.20})$	$^{+0.14}_{-0.10}\,(^{+0.22}_{-0.16})$
$\mathcal{B}^{WW}/\mathcal{B}^{ZZ}$	$8.11^{+0.15}_{-0.15}$	$0.97^{+0.13}_{-0.11}\,(^{+0.13}_{-0.11})$	$^{+0.09}_{-0.08}\,(^{+0.10}_{-0.09})$	$^{+0.08}_{-0.07}\,(^{+0.08}_{-0.07})$
$\mathcal{B}^{\tau\tau}/\mathcal{B}^{ZZ}$	$2.29^{+0.04}_{-0.04}$	$0.98^{+0.17}_{-0.16}\,(^{+0.17}_{-0.15})$	$^{+0.11}_{-0.10}\,(^{+0.11}_{-0.10})$	$^{+0.13}_{-0.12}\,(^{+0.13}_{-0.12})$
$\mathcal{B}^{\gamma\gamma}/\mathcal{B}^{ZZ}$	$0.084^{+0.002}_{-0.002}$	$1.01^{+0.12}_{-0.11}\,(^{+0.12}_{-0.11})$	$^{+0.10}_{-0.09}\,(^{+0.10}_{-0.09})$	$^{+0.06}_{-0.05}\,(^{+0.06}_{-0.05})$

channels. This leads to anticorrelations between the $\sigma^i\mathcal{B}^{ZZ}$ and $R^{f/ZZ}$ parameters, and a positive correlation structure among the ratios themselves, and among the $\sigma^i\mathcal{B}^{ZZ}$ parameters. Additional correlation patterns arise from the increased granularity of the STXS stage 1.2 measurement. In particular, strong anticorrelations are observed between the ggH STXS bins with additional jets and the 'qqH other' STXS bin. Events from the 'qqH other' region are difficult to distinguish from ggH events as the additional jets in the event are typically soft or fall outside of the experimental acceptance. Therefore, these events tend to populate the analysis regions targeting ggH production, leading to a large anticorrelation. This in turn enhances the positive correlation structure between the ggH cross sections with additional jets. Strong correlations are also observed between the high-p_T^V regions for the WH and ZH production processes, and between the high p_T^H ttH and inclusive tH regions.

In accordance with the production cross section results, a measurement is performed at this granularity, in which the SM Higgs boson branching fractions are assumed. The results are provided as supplementary material to the paper [41]. This fit considers only the 32 STXS bin cross sections as POIs, and incorporates the theoretical uncertainties associated with the Higgs boson branching fractions. Results in the signal strength formalism are provided as supplementary material [41].



Figure 11. The correlations between the 36 parameters considered in the STXS stage 1.2 measurement. The size of the correlations is indicated by the colour scale. Correlations of absolute magnitude smaller than 0.005 are not shown.

Measurements are also performed in an additional parametrization, which introduces a separate parameter of interest for each STXS bin i, in each decay channel f,

$$\mu^{if} = \frac{\left[\sigma^{i}\,\mathcal{B}^{f}\right]_{\text{obs}}}{\left[\sigma^{i}\,\mathcal{B}^{f}\right]_{\text{SM,HO}}}. \tag{12}$$

This measurement is performed for the combination of all input analyses, except the $H \to$ inv and off-shell $H \to ZZ \to 4\ell$ channels. The parameters are defined using the partitions of phase space that were used in the original publications. The channels that do not implement the STXS stage 1.2 process splittings are fit with the production process cross sections. The parameters for the $H \to \gamma\gamma$ and $H \to ZZ$ decays are restricted to nonnegative values to avoid an overall negative event yield in analysis regions where the background contamination is sufficiently low.

As each decay channel is defined with a different binning scheme, the treatment of common theoretical uncertainty NPs is ambiguous. Consequently, all theoretical uncertainties affecting the cross section normalizations and branching fractions are included in the fit. In other words, this result is provided in the signal strength formalism, and can therefore be used directly for reinterpretation without the need to define an additional theoretical uncertainty scheme.

The best fit values and 68% CL intervals are shown for the 97 POIs in figure 12. Each panel represents a different Higgs boson decay channel. The corresponding numerical values are provided, along with the expected 68% CL intervals, in table 8. This result represents the most granular measurement of the Higgs boson ever performed by the CMS Collaboration. The correlations between the measured parameters are provided as supplementary material to the paper [41].

The p-value for the compatibility with the SM follows the same trend as the other signal yield parametrizations, showing some tension with the SM hypothesis. Sizeable deviations are observed for a



Figure 12. The best fit values (empty circles) and 68% CL intervals (coloured lines) for the measurement of the products of the cross sections and branching fractions. Theoretical uncertainties affecting the cross section normalizations and branching fractions are included in the fit. The best fit values for the products of cross sections and branching fractions are obtained by multiplying the fit parameters μ^{if} by the SM predictions at the highest available order. Different panels show the measurements for the different Higgs boson decay channels. The (ttH $p_T^H > 300$ GeV, H → bb), (ggH $p_T^H > 300$ GeV, H → WW), and (ggH 0J $p_T^H < 10$ GeV, H → ττ) best fit values and 68% CL intervals are entirely contained in the negative domain, and are represented by arrows as they cannot be shown on the log-scale axes. Arrows are also used to indicate 68% CL intervals that extend into the negative domain. The H → γγ and H → ZZ parameters are restricted to nonnegative values, which is marked by the hatched grey lines in the corresponding panels.

number of kinematic regions including the high p_T^V WH leptonic STXS bins for the $H \to bb$, $H \to WW$, and $H \to \tau\tau$ channels, the ZH leptonic $p_T^V > 250$ STXS bin for the $H \to bb$ channel, the ggH 0J STXS bins for the $H \to \tau\tau$ channel, and the tH measurement in the $H \to \gamma\gamma$ channel. The overall compatibility with the SM is measured to be $p_{SM} = 0.006$.

9. Interpretation of measurements in the coupling modifier framework

The coupling modifier (κ) framework [23] introduces parameters that scale Higgs boson production cross sections and decay rates. These coupling modifiers describe deviations in the couplings of the Higgs boson to other particles.

The production cross sections σ^j, or the partial decay widths Γ^j, associated with the SM particle j, scale with a factor κ_j^2 such that

$$\kappa_j^2 = \sigma^j/\sigma^j_{SM} \text{ or } \kappa_j^2 = \Gamma^j/\Gamma^j_{SM}. \tag{13}$$

In the SM, all values of κ_j are positive and equal to unity. The higher-order accuracy of the QCD and EW corrections to the SM production cross sections and branching fractions are assumed to be preserved when κ_j does not equal unity. For the ranges of parameters considered in this study, the dominant higher-order QCD and EW corrections factorize from the coupling rescaling, validating the assumption.

Several different models within the κ-framework are probed. These include a model where only the decays of the Higgs boson to SM particles are allowed; a model that considers additional BSM decays, with $|\kappa_W|$ and $|\kappa_Z|$ constrained to be less than or equal to 1 to avoid a complete degeneracy in the total Higgs boson decay width; and a model that incorporates off-shell Higgs boson measurements to alleviate this degeneracy by directly constraining the total Higgs boson decay width from data. Measurements are performed in additional models that include ratios of coupling modifiers. Finally, a model that considers NLO EW corrections to the production and decay rates from the Higgs boson self-coupling is probed.

Under the assumption that the Higgs boson does not decay to BSM particles, the product of the production cross section and branching fraction for production process i and decay channel f can be expressed as

$$\mu^{if}(\vec{\kappa}) \left[\sigma^i \mathcal{B}^f\right]_{SM,HO} = \sigma^i(\vec{\kappa})\mathcal{B}^f(\vec{\kappa}) = \frac{\sigma^i(\vec{\kappa})\Gamma^f(\vec{\kappa})}{\Gamma_H(\vec{\kappa})}, \tag{14}$$

where $\Gamma_H(\vec{\kappa})$ is the total decay width of the Higgs boson and $\Gamma^f(\vec{\kappa})$ is the partial width of the Higgs boson decay to the final state f. The coupling modifier scaling functions $\mu^{if}(\vec{\kappa})$ enter the signal yield parametrization, as shown in equation (5).

Two different configurations of coupling modifiers are introduced. In one of these configurations, all loops in Higgs boson production and decay diagrams are resolved, and coupling modifiers are introduced for each tree-level Higgs boson coupling to another SM particle. The second configuration introduces effective coupling modifiers κ_g, κ_γ, and $\kappa_{Z\gamma}$ to describe ggH production, and the $H \to \gamma\gamma$ and $H \to Z\gamma$ decays, which only occur via loop diagrams in the SM. Table 9 shows the scaling of production cross sections, partial decay widths, and the total Higgs boson decay width for both configurations.

The interference between different diagrams gives sensitivity to the relative signs of a number of coupling modifiers. As shown in table 9, ggH production is sensitive to the relative sign of κ_b and κ_t, ggZH production is sensitive to the relative sign of κ_Z and κ_t, and tHq and tHW production, as well as the $H \to \gamma\gamma$ and $H \to Z\gamma$ decays, are sensitive to the relative sign of κ_W and κ_t.

When additional BSM decays are allowed, the Higgs boson total decay width is modified as

$$\frac{\Gamma_H}{\Gamma_{H,SM}} = \frac{\kappa_H^2}{1 - (\mathcal{B}_{inv} + \mathcal{B}_{undet})}, \tag{15}$$

where $\mathcal{B}_{inv}$ represents the Higgs boson branching fraction to invisible final states that do not appreciably interact with the detector material and thus escape detection. Such decays are constrained by the $H \to$ inv searches included in the combination. The $\mathcal{B}_{undet}$ parameter represents the Higgs boson branching fraction to undetected final states, which are not directly measured by any of the input analyses. This includes decays to BSM particles that remain undetected because they produce event topologies that are not explictly searched for, as well as modifications to branching fractions into SM final states that are not directly measured, such as $H \to cc$. Together, $\mathcal{B}_{inv}$ and $\mathcal{B}_{undet}$ make up the total BSM branching fraction.

Table 8. Best fit values and 68% CL intervals for the fit that introduces a separate POI for each STXS bin in each decay channel. The cross sections are defined in the fiducial region $|y_H| < 2.5$. The values are normalized to the SM predictions. The fit is performed in the signal strength formalism, such that the theoretical uncertainties in the SM predictions are folded into the measurement. The expected intervals are given in parentheses. Some of the parameters are restricted to nonnegative values, as described in the text. Truncated intervals are reported for these parameters if the 68% CL interval is not fully contained in the positive domain.

Parameters	SM $\sigma^i\mathcal{B}^f$ (fb)	Best fit/SM pred.
$H \to \gamma\gamma$		
ggH 0J $p_T^H < 10$	15.2	$0.58^{+0.38}_{-0.26}\,(^{+0.66}_{-0.48})$
ggH 0J $p_T^H > 10$ + bbH	44.3	$1.35^{+0.27}_{-0.20}\,(^{+0.18}_{-0.16})$
ggH 1J $p_T^H < 60$	16.2	$0.84^{+0.38}_{-0.36}\,(^{+0.40}_{-0.36})$
ggH 1J $60 < p_T^H < 120$	11.2	$1.26^{+0.34}_{-0.30}\,(^{+0.32}_{-0.28})$
ggH 1J $120 < p_T^H < 200$	2.00	$1.30^{+0.55}_{-0.46}\,(^{+0.51}_{-0.44})$
ggH $\geqslant$2J $0 < m_{jj} < 350$, $p_T^H < 60$	2.82	$1.39^{+1.40}_{-1.34}\,(^{+1.44}_{-1.00})$
ggH $\geqslant$2J $0 < m_{jj} < 350$, $60 < p_T^H < 120$	4.53	$0.00^{+0.66}_{-0.00}\,(^{+0.74}_{-0.63})$
ggH $\geqslant$2J $0 < m_{jj} < 350$, $120 < p_T^H < 200$	2.12	$0.33^{+0.58}_{-0.33}\,(^{+0.65}_{-0.55})$
ggH $200 < p_T^H < 300$	1.10	$1.01^{+0.56}_{-0.41}\,(^{+0.52}_{-0.40})$
ggH $300 < p_T^H < 450$	0.28	$0.53^{+0.76}_{-0.53}\,(^{+0.76}_{-0.62})$
ggH $p_T^H > 450$	0.040	$3.56^{+3.12}_{-2.48}\,(^{+1.94}_{-1.00})$
VBF-like $350 < m_{jj} < 700$, $p_T^{Hjj} < 25$	1.59	$0.77^{+0.82}_{-0.67}\,(^{+1.31}_{-1.00})$
VBF-like $350 < m_{jj} < 700$, $p_T^{Hjj} > 25$	1.25	$2.86^{+1.98}_{-1.32}\,(^{+1.14}_{-0.97})$
VBF-like $m_{jj} > 700$, $p_T^{Hjj} < 25$	1.60	$0.54^{+0.54}_{-0.34}\,(^{+0.91}_{-0.63})$
VBF-like $m_{jj} > 700$, $p_T^{Hjj} > 25$	0.73	$0.46^{+1.50}_{-0.46}\,(^{+0.98}_{-0.85})$
qqH $60 < m_{jj} < 120$ (VH-topo)	1.22	$1.23^{+1.03}_{-0.94}\,(^{+1.00}_{-0.94})$
qqH $p_T^H > 200$	0.37	$1.44^{+0.69}_{-0.62}\,(^{+0.66}_{-0.58})$
WH lep $p_T^V < 75$	0.47	$1.65^{+1.33}_{-1.29}\,(^{+1.42}_{-1.00})$
WH lep $75 < p_T^V < 150$	0.29	$1.54^{+1.27}_{-1.12}\,(^{+1.34}_{-0.96})$
WH lep $p_T^V > 150$	0.12	$1.02^{+0.99}_{-0.79}\,(^{+1.04}_{-0.79})$
ZH lep	0.54	$1.29^{+0.80}_{-0.61}\,(^{+0.72}_{-0.62})$
ttH $p_T^H < 60$	0.26	$0.74^{+1.09}_{-0.74}\,(^{+1.20}_{-0.82})$
ttH $60 < p_T^H < 120$	0.40	$1.30^{+0.68}_{-0.59}\,(^{+0.64}_{-0.51})$
ttH $120 < p_T^H < 200$	0.29	$0.79^{+0.60}_{-0.48}\,(^{+0.62}_{-0.50})$
ttH $200 < p_T^H < 300$	0.12	$0.99^{+0.95}_{-0.79}\,(^{+0.75}_{-0.61})$
ttH $p_T^H > 300$	0.061	$0.00^{+1.45}_{-0.00}\,(^{+1.06}_{-0.99})$
tH	0.20	$8.35^{+3.64}_{-4.56}\,(^{+4.43}_{-1.00})$
$H \to ZZ$		
ggH 0J $p_T^H < 10$	182	$0.92^{+0.33}_{-0.25}\,(^{+0.34}_{-0.28})$
ggH 0J $p_T^H > 10$ + bbH	529	$1.22^{+0.22}_{-0.20}\,(^{+0.19}_{-0.17})$
ggH 1J $p_T^H < 60$	194	$1.26^{+0.56}_{-0.46}\,(^{+0.61}_{-0.55})$
ggH 1J $60 < p_T^H < 120$	134	$0.50^{+0.38}_{-0.36}\,(^{+0.56}_{-0.52})$
ggH 1J $120 < p_T^H < 200$	24.0	$2.50^{+1.28}_{-1.06}\,(^{+1.03}_{-0.80})$
ggH $\geqslant$2J $0 < m_{jj} < 350$, $p_T^H < 60$	33.7	$0.00^{+0.81}_{-0.00}\,(^{+1.65}_{-1.00})$
ggH $\geqslant$2J $0 < m_{jj} < 350$, $60 < p_T^H < 120$	54.3	$1.34^{+0.89}_{-0.65}\,(^{+0.99}_{-0.78})$
ggH $\geqslant$2J $0 < m_{jj} < 350$, $120 < p_T^H < 200$	25.4	$0.97^{+0.97}_{-0.67}\,(^{+1.11}_{-0.82})$
ggH VBF-topo	26.5	$2.02^{+2.72}_{-2.02}\,(^{+2.09}_{-1.01})$
ggH $p_T^H > 200$	17.0	$0.41^{+0.53}_{-0.36}\,(^{+1.05}_{-0.78})$
qqH other	75.6	$0.00^{+1.63}_{-0.00}\,(^{+2.59}_{-1.00})$
qqH $60 < m_{jj} < 120$ (VH-topo)	14.6	$1.29^{+1.95}_{-1.29}\,(^{+2.05}_{-1.00})$
qqH $350 < m_{jj} < 700$, $p_T^{Hjj} < 25$	10.7	$0.32^{+2.77}_{-0.32}\,(^{+2.59}_{-1.00})$
qqH $m_{jj} > 700$, $p_T^{Hjj} < 25$	15.5	$0.00^{+0.66}_{-0.00}\,(^{+1.32}_{-0.90})$
qqH $p_T^{Hjj} > 25$	9.21	$3.03^{+5.67}_{-3.03}\,(^{+4.66}_{-0.99})$
qqH $p_T^H > 200$	4.41	$0.00^{+0.79}_{-0.00}\,(^{+2.14}_{-1.00})$
VH lep $p_T^V < 150$	14.4	$8.36^{+4.00}_{-3.65}\,(^{+1.94}_{-1.00})$
VH lep $p_T^V > 150$	2.59	$0.00^{+3.50}_{-0.00}\,(^{+4.45}_{-1.00})$
ttH + tH	16.0	$0.16^{+1.00}_{-0.16}\,(^{+1.41}_{-0.83})$
$H \to \mu\mu$		
ggH + bbH + ttH + tH	10.7	$0.69^{+0.66}_{-0.73}\,(^{+0.69}_{-0.67})$
qqH + VH	1.27	$1.71^{+0.84}_{-0.76}\,(^{+0.74}_{-0.65})$

(Continued.)

Table 8. (Continued.)

$H \to Z\gamma$		
ggH + bbH + ttH + tH	77.6	$3.67^{+1.37}_{-1.22}\,(^{+1.14}_{-1.14})$
qqH + VH	9.28	$-2.39^{+3.04}_{-2.82}\,(^{+3.18}_{-2.92})$
$H \to WW$		
ggH 0J	5670	$0.86^{+0.12}_{-0.14}\,(^{+0.14}_{-0.13})$
ggH 1J $p_T^H < 60$	1572	$1.71^{+0.45}_{-0.42}\,(^{+0.39}_{-0.36})$
ggH 1J $p_T^H > 60$	1284	$0.32^{+0.25}_{-0.26}\,(^{+0.30}_{-0.26})$
ggH $\geqslant$2J $p_T^H < 200$	1135	$1.65^{+0.85}_{-0.68}\,(^{+0.63}_{-0.54})$
ggH $200 < p_T^H < 300$	107	$2.68^{+1.42}_{-1.21}\,(^{+1.09}_{-0.94})$
ggH $p_T^H > 300$	31.2	$-4.18^{+3.27}_{-3.38}\,(^{+2.66}_{-2.41})$
qqH $350 < m_{jj} < 700$	125	$0.27^{+0.99}_{-0.81}\,(^{+0.92}_{-0.85})$
qqH $m_{jj} > 700$	162	$0.68^{+0.35}_{-0.31}\,(^{+0.35}_{-0.33})$
qqH $p_T^H > 200$	35.8	$1.18^{+0.72}_{-0.66}\,(^{+0.72}_{-0.63})$
qqH $60 < m_{jj} < 120$ (VH-topo)	118	$4.30^{+3.43}_{-3.58}\,(^{+3.13}_{-3.13})$
WH lep $p_T^V < 150$	74.0	$1.36^{+1.09}_{-0.96}\,(^{+0.83}_{-0.80})$
WH lep $p_T^V > 150$	11.7	$4.08^{+1.65}_{-1.59}\,(^{+1.31}_{-1.22})$
ZH lep $p_T^V < 150$	43.0	$2.67^{+1.05}_{-0.82}\,(^{+0.79}_{-0.64})$
ZH lep $p_T^V > 150$	9.31	$-0.38^{+1.29}_{-0.64}\,(^{+1.08}_{-0.89})$
ttH + tH	130	$1.45^{+0.37}_{-0.32}\,(^{+0.34}_{-0.31})$
$H \to \tau\tau$		
ggH 0J $p_T^H < 10$	416	$-1.67^{+0.83}_{-0.84}\,(^{+0.96}_{-0.93})$
ggH 0J $p_T^H > 10$	1182	$-0.51^{+0.54}_{-0.56}\,(^{+0.53}_{-0.52})$
ggH 1J $p_T^H < 60$	443	$0.86^{+0.65}_{-0.66}\,(^{+0.67}_{-0.64})$
ggH 1J $60 < p_T^H < 120$	307	$1.87^{+0.59}_{-0.51}\,(^{+0.53}_{-0.49})$
ggH 1J $120 < p_T^H < 200$	54.8	$1.42^{+0.64}_{-0.59}\,(^{+0.62}_{-0.56})$
ggH $\geqslant$2J $p_T^H < 200$	320	$1.63^{+0.73}_{-0.59}\,(^{+0.64}_{-0.55})$
ggH $200 < p_T^H < 300$	30.1	$1.43^{+0.76}_{-0.68}\,(^{+0.76}_{-0.65})$
ggH $p_T^H > 300$	8.79	$1.91^{+0.99}_{-0.80}\,(^{+0.89}_{-0.78})$
qqH other	206	$-0.98^{+1.20}_{-1.21}\,(^{+1.22}_{-1.18})$
qqH $350 < m_{jj} < 700$	35.1	$1.30^{+1.05}_{-1.03}\,(^{+1.02}_{-1.01})$
qqH $m_{jj} > 700$	45.7	$1.27^{+0.32}_{-0.32}\,(^{+0.31}_{-0.30})$
qqH $p_T^H > 200$	10.1	$0.20^{+0.34}_{-0.33}\,(^{+0.36}_{-0.34})$
WH lep $p_T^V < 150$	20.8	$0.61^{+0.90}_{-0.89}\,(^{+0.83}_{-0.79})$
WH lep $p_T^V > 150$	3.30	$2.55^{+1.46}_{-1.16}\,(^{+1.21}_{-1.11})$
ZH lep $p_T^V < 150$	12.1	$1.76^{+0.89}_{-0.76}\,(^{+0.78}_{-0.70})$
ZH lep $p_T^V > 150$	2.62	$1.93^{+1.00}_{-0.70}\,(^{+0.79}_{-0.62})$
ttH + tH	36.5	$0.38^{+0.41}_{-0.40}\,(^{+0.49}_{-0.42})$
$H \to bb$		
ggH $300 < p_T^H < 450$	71.5	$-1.65^{+6.74}_{-6.94}\,(^{+6.64}_{-6.45})$
ggH $450 < p_T^H < 650$	8.85	$2.36^{+2.04}_{-1.61}\,(^{+1.74}_{-1.60})$
ggH $p_T^H > 650$	1.24	$2.64^{+3.26}_{-2.67}\,(^{+2.90}_{-2.64})$
qqH	5760	$1.54^{+0.66}_{-0.57}\,(^{+0.49}_{-0.44})$
WH lep $150 < p_T^V < 250$	23.1	$0.14^{+0.66}_{-0.65}\,(^{+0.64}_{-0.63})$
WH lep $p_T^V > 250$	7.52	$1.88^{+0.50}_{-0.46}\,(^{+0.42}_{-0.40})$
ZH lep $75 < p_T^V < 150$	49.4	$1.40^{+0.75}_{-0.71}\,(^{+0.72}_{-0.69})$
ZH lep $150 < p_T^V < 250$, 0J	8.68	$0.45^{+0.48}_{-0.46}\,(^{+0.56}_{-0.51})$
ZH lep $150 < p_T^V < 250$, $\geqslant$ 1J	9.93	$-0.50^{+1.09}_{-1.09}\,(^{+1.05}_{-0.91})$
ZH lep $p_T^V > 250$	5.73	$1.65^{+0.47}_{-0.38}\,(^{+0.34}_{-0.31})$
ttH $p_T^H < 60$	65.2	$0.52^{+1.82}_{-1.74}\,(^{+2.21}_{-1.95})$
ttH $60 < p_T^H < 120$	101	$0.43^{+1.26}_{-1.41}\,(^{+1.33}_{-1.29})$
ttH $120 < p_T^H < 200$	74.5	$1.00^{+0.98}_{-0.81}\,(^{+0.91}_{-0.85})$
ttH $200 < p_T^H < 300$	31.2	$0.38^{+0.87}_{-0.92}\,(^{+0.93}_{-0.86})$
ttH $p_T^H > 300$	15.5	$-1.37^{+1.20}_{-1.04}\,(^{+1.17}_{-1.09})$

Table 9. Normalization scaling factors for all relevant production cross sections, partial decay widths, and the total Higgs boson decay width. For the κ parameters representing loop processes, the resolved scaling in terms of the fundamental SM couplings is also given. Only the dominant terms in the resolved scaling factor functions are provided in the table. The contributions are calculated for $m_{\mathrm{H}} = 125.38\,\mathrm{GeV}$.

	Loops	Interference	Effective scaling factor	Resolved scaling factor
Production				
$\sigma(\mathrm{ggH})$	✓	b-t	κ_{g}^2	$1.04\kappa_{\mathrm{t}}^2 + 0.002\kappa_{\mathrm{b}}^2 - 0.038\kappa_{\mathrm{t}}\kappa_{\mathrm{b}}$
$\sigma(\mathrm{VBF})$	—	—		$0.73\kappa_{\mathrm{W}}^2 + 0.27\kappa_{\mathrm{Z}}^2$
$\sigma(\mathrm{WH})$	—	—		κ_{W}^2
$\sigma(\mathrm{qq/qg} \to \mathrm{ZH})$	—	—		κ_{Z}^2
$\sigma(\mathrm{gg} \to \mathrm{ZH})$	✓	Z-t		$2.46\kappa_{\mathrm{Z}}^2 + 0.46\kappa_{\mathrm{t}}^2 - 1.90\kappa_{\mathrm{Z}}\kappa_{\mathrm{t}}$
$\sigma(\mathrm{ttH})$	—	—		κ_{t}^2
$\sigma(\mathrm{gb} \to \mathrm{tHW})$	—	W-t		$2.91\kappa_{\mathrm{t}}^2 + 2.31\kappa_{\mathrm{W}}^2 - 4.22\kappa_{\mathrm{t}}\kappa_{\mathrm{W}}$
$\sigma(\mathrm{qb} \to \mathrm{tHq})$	—	W-t		$2.63\kappa_{\mathrm{t}}^2 + 3.58\kappa_{\mathrm{W}}^2 - 5.21\kappa_{\mathrm{t}}\kappa_{\mathrm{W}}$
$\sigma(\mathrm{bbH})$	—	—		κ_{b}^2
Partial decay width				
Γ^{ZZ}	—	—		κ_{Z}^2
Γ^{WW}	—	—		κ_{W}^2
$\Gamma^{\gamma\gamma}$	✓	W-t	κ_{γ}^2	$1.59\kappa_{\mathrm{W}}^2 + 0.07\kappa_{\mathrm{t}}^2 - 0.67\kappa_{\mathrm{W}}\kappa_{\mathrm{t}}$
$\Gamma^{\mathrm{Z}\gamma}$	✓	W-t	$\kappa_{\mathrm{Z}\gamma}^2$	$1.118\kappa_{\mathrm{W}}^2 + 0.003\kappa_{\mathrm{t}}^2 - 0.124\kappa_{\mathrm{W}}\kappa_{\mathrm{t}}$
$\Gamma^{\tau\tau}$	—	—		κ_{τ}^2
Γ^{bb}	—	—		κ_{b}^2
$\Gamma^{\mu\mu}$	—	—		κ_{μ}^2
Total width for $\mathcal{B}_{\mathrm{BSM}} = 0$				
Γ_{H}	✓	—	κ_{H}^2	$0.58\kappa_{\mathrm{b}}^2 + 0.22\kappa_{\mathrm{W}}^2 + 0.08\kappa_{\mathrm{g}}^2 + + 0.06\kappa_{\tau}^2 + 0.027\kappa_{\mathrm{Z}}^2 + 0.029\kappa_{\mathrm{c}}^2 + + 0.0023\kappa_{\gamma}^2 + 0.0016\kappa_{\mathrm{Z}\gamma}^2 + + 0.00025\kappa_{\mathrm{s}}^2 + 0.00022\kappa_{\mu}^2$

9.1. Constraints on coupling modifiers with resolved loops

Figure 13 shows the results of the resolved coupling modifier fit. In total, six coupling parameters are probed: κ_{W}, κ_{Z}, κ_{t}, κ_{b}, κ_{τ}, and κ_{μ}, and no additional BSM decays of the Higgs boson are considered. The interference terms that enter the ggH, ggZH, tHq, and tHW production scaling factors, and the $\mathrm{H} \to \gamma\gamma$ and $\mathrm{H} \to \mathrm{Z}\gamma$ decay scaling factors, provide sensitivity to the relative signs of κ_{W}, κ_{Z}, κ_{b}, and κ_{t}. As the fit is only sensitive to the relative signs, the value of κ_{t} is restricted to the positive domain without loss of generality, while both positive and negative values of κ_{W}, κ_{Z}, and κ_{b} are allowed. The measurements are not sensitive to the signs of κ_{τ} and κ_{μ}, and therefore these parameters are restricted to be positive.

The left plot in figure 13 shows the best fit points with the corresponding 68% and 95% CL intervals. The global minimum of the fit corresponds to the domain where all of the coupling modifiers are positive. Nevertheless, as the $\mathrm{H} \to \mathrm{ZZ}$ decay and ZH production depend only on the absolute value of κ_{Z}, a secondary minimum around $\kappa_{\mathrm{Z}} = -1$ is observed. The interference between diagrams in ggZH production provides a term proportional to $\kappa_{\mathrm{Z}}\kappa_{\mathrm{t}}$ in the scaling function, and this helps break the degeneracy between the two signs. Given the ggZH contributions are small, the minimum around $\kappa_{\mathrm{Z}} = -1$ is not strongly disfavoured, and the 95% CL interval includes negative values of κ_{Z}. The sensitivity to the relative sign of κ_{b} and κ_{t} enters through a weak interference term in the ggH production scaling function, and therefore negative values of κ_{b} are included in the 68% CL interval. Negative values of κ_{W} are strongly disfavoured by the data, because of the interference term proportional to $\kappa_{\mathrm{W}}\kappa_{\mathrm{t}}$ in the $\mathrm{H} \to \gamma\gamma$ decay.

The numerical values for the best fit points and 68% CL intervals for the resolved coupling modifiers are provided in table 10. The total uncertainties are decomposed into their systematic and statistical parts, and the corresponding expected 68% CL intervals are provided in parentheses. The vector boson coupling modifiers, κ_{W} and κ_{Z}, are measured with 68% CL intervals of approximately ±6%. The coupling modifiers for third generation fermions are equally well constrained with 68% CL intervals ranging from ±6% for κ_{t} to ±12% for κ_{b}. The coupling modifier to muons is measured with a 68% CL interval



Figure 13. The coupling modifiers of the Higgs boson to fermions and gauge bosons, in the resolved coupling modifier measurement. In the upper plot, the thick (thin) black lines indicate the 68% (95%) CL intervals around the best fit points (empty circles). For this model, both positive and negative values of κ_W, κ_Z, and κ_b are considered, while κ_t is restricted to the positive domain without loss of generality, as indicated by the hatched box. In the lower plot, the measured coupling modifiers are shown as functions of the fermion or gauge boson mass, where v is the vacuum expectation value of the Brout–Englert–Higgs field. In this fit, all coupling modifiers are restricted to the positive domain. The 68% and 95% CL intervals are determined around the minimum that contains the best fit point. The b quark mass is evaluated at a scale equal to m_H. The uncertainties in the particle mass values are not shown in the figure. For gauge bosons, the square root of the coupling modifier is used to keep a linear proportionality to the mass, as predicted by the SM.

of ±20%. These measurements show good compatibility with the SM hypothesis, with an overall p-value of $p_{SM} = 0.12$.

An additional fit is performed in which all κ_j are restricted to the positive domain. The right plot in figure 13 shows the measured coupling modifiers from this fit as functions of the mass of the probed particles. For the b quark, the running mass evaluated at a scale equal to $m_H = 125.38$ GeV is used: $m_b(m_H) = 2.76$ GeV. The observed agreement over three orders of magnitude of mass is a powerful test of the validity of the SM.

Table 10. Best fit values and 68% CL intervals for the Higgs boson coupling modifiers in the resolved coupling configuration. The total 68% CL intervals are decomposed into their statistical and systematic components. The expected intervals are given in parentheses. For κ_b the 68% CL interval is not contiguous, and the quoted interval corresponds to the minimum containing the best fit point. The 68% CL interval for this parameter also includes the range $-1.11 < \kappa_b < -0.89$ ($-1.13 < \kappa_b < -0.90$) for the observed (expected) case.

Parameters	Best fit	Stat	Syst
κ_W	$1.05^{+0.06}_{-0.05}$	$^{+0.04}_{-0.04}$	$^{+0.04}_{-0.04}$
	$\left(^{+0.06}_{-0.05}\right)$	$\left(^{+0.04}_{-0.04}\right)$	$\left(^{+0.04}_{-0.03}\right)$
κ_Z	$1.07^{+0.06}_{-0.06}$	$^{+0.04}_{-0.04}$	$^{+0.04}_{-0.04}$
	$\left(^{+0.06}_{-0.06}\right)$	$\left(^{+0.04}_{-0.04}\right)$	$\left(^{+0.04}_{-0.04}\right)$
κ_t	$0.91^{+0.06}_{-0.07}$	$^{+0.04}_{-0.05}$	$^{+0.04}_{-0.04}$
	$\left(^{+0.07}_{-0.08}\right)$	$\left(^{+0.04}_{-0.05}\right)$	$\left(^{+0.05}_{-0.06}\right)$
κ_b	$0.98^{+0.12}_{-0.11}$	$^{+0.08}_{-0.08}$	$^{+0.09}_{-0.08}$
	$\left(^{+0.12}_{-0.12}\right)$	$\left(^{+0.08}_{-0.08}\right)$	$\left(^{+0.09}_{-0.08}\right)$
κ_τ	$0.90^{+0.08}_{-0.07}$	$^{+0.05}_{-0.04}$	$^{+0.06}_{-0.05}$
	$\left(^{+0.08}_{-0.07}\right)$	$\left(^{+0.05}_{-0.05}\right)$	$\left(^{+0.07}_{-0.06}\right)$
κ_μ	$1.09^{+0.20}_{-0.21}$	$^{+0.18}_{-0.19}$	$^{+0.08}_{-0.08}$
	$\left(^{+0.21}_{-0.23}\right)$	$\left(^{+0.19}_{-0.22}\right)$	$\left(^{+0.08}_{-0.08}\right)$

9.2. Constraints on coupling modifiers with effective loops

In the effective coupling modifier configuration, ggH production scales according to κ_g^2, and the sensitivity to the relative sign of κ_b and κ_t is lost. Therefore, for the fits shown in this section, both positive and negative values of κ_W and κ_Z are considered, while κ_t is restricted to the positive domain without loss of generality. The other coupling modifiers, including κ_b, are restricted to be positive as the fit is not sensitive to their signs. Results in which all κ_j are restricted to be positive are provided as supplementary material [41].

Fits are performed with three different assumptions on the total Higgs boson decay width, and the results from each of the three models are summarized in figure 14. The first model, shown in blue, assumes no additional decays of the Higgs boson to BSM particles. The second model introduces $\mathcal{B}_{inv}$ and $\mathcal{B}_{undet}$ as additional parameters. However, a constraint requiring $|\kappa_W|$ and $|\kappa_Z|$ to be less than or equal to one is imposed. This alleviates a complete degeneracy in the total Higgs boson decay width, where each coupling modifier can be scaled up equally to account for a nonzero $\mathcal{B}_{undet}$. The choice of restricting κ_W and κ_Z is motivated by the fact that this condition is satisfied for a wide class of BSM models, including those with an arbitrary number of Higgs doublets, with or without additional Higgs singlets [23]. The results are shown in orange in figure 14, where the excluded parameter space is indicated by the hatched boxes. In this fit, the $\mathcal{B}_{inv}$ and $\mathcal{B}_{undet}$ parameters are constrained to be less than 13% and 21% at the 95% CL, respectively.

In the third model, the off-shell $H \to ZZ \to 4\ell$ analysis is introduced into the combination to constrain the total Higgs boson decay width directly from data. The fit is able to leverage the different scaling behaviours of off-shell and on-shell Higgs boson production, such that the external constraint of $|\kappa_V| \leqslant 1$ can be alleviated. Crucially, the scaling behaviour for off-shell Higgs boson production is independent of the total Higgs boson decay width.

To describe the scaling behaviour of off-shell ggH production, the κ_g parameter cannot be used directly because of a dependence on the energy scale q^2. Instead, an additional coupling modifier κ_Q is introduced [87]. This modifier accounts for contributions resulting from the possible presence of an additional heavy quark in the ggH loop. In the SM, its value is 0. Importantly, the κ_Q parameter is independent of q^2.

The total binned probability density in the off-shell analysis region, r, is the sum of different contributions, each with a different scaling behaviour as a function of the coupling modifiers,

$$\begin{aligned}\mathcal{P}_r =& \kappa_Z^4 \mathcal{P}_r^{(\mathrm{VBF+VH})_Z,\mathrm{sig}} + \kappa_W^2\kappa_Z^2 \mathcal{P}_r^{(\mathrm{VBF+VH})_W,\mathrm{sig}} + \kappa_W\kappa_Z^3 \mathcal{P}_r^{(\mathrm{VBF+VH})_{WZ},\mathrm{sig}} \\ &+ \kappa_Z^2 \mathcal{P}_r^{(\mathrm{VBF+VH})_Z,\mathrm{int}} + \kappa_W\kappa_Z \mathcal{P}_r^{(\mathrm{VBF+VH})_W,\mathrm{int}} + \kappa_t^2\kappa_Z^2 \mathcal{P}_r^{(\mathrm{ggH})_t,\mathrm{sig}} \\ &+ \kappa_Q^2\kappa_Z^2 \mathcal{P}_r^{(\mathrm{ggH})_Q,\mathrm{sig}} + \kappa_t\kappa_Q\kappa_Z^2 \mathcal{P}_r^{(\mathrm{ggH})_{tQ},\mathrm{sig}} \\ &+ \kappa_t\kappa_Z \mathcal{P}_r^{(\mathrm{ggH})_t,\mathrm{int}} + \kappa_Q\kappa_Z \mathcal{P}_r^{(\mathrm{ggH})_Q,\mathrm{int}} \\ &+ \frac{\kappa_Z^4\Gamma^{\mathrm{H}}_{\mathrm{SM}}}{\Gamma^{\mathrm{H}}} \mathcal{P}_r^{\mathrm{cross}} + \mathcal{P}_r^{\mathrm{bkg}},\end{aligned} \tag{16}$$

where the notation $\mathcal{P}_r^i = s_r^i \rho_{\mathrm{sig},r}^i$ has been used to represent the normalized templates (cf equation (1)), and all explicit dependence on the observables, POIs, and NPs has been dropped for simplicity. The decay channel index f has also been dropped as all contributions are from the $\mathrm{H} \to \mathrm{ZZ} \to 4\ell$ decay. The contributions are grouped into those that exhibit the same coupling modifier scaling behaviour. The terms labelled 'sig', 'int', and 'cross' refer to the pure signal contributions, the contributions from interference between the signal and background amplitudes, and the cross-feed component, respectively. This last component concerns on-shell Higgs boson decays to $2\ell + X$, where X contains additional leptons that can mimic off-shell $\mathrm{H} \to \mathrm{ZZ} \to 4\ell$ events. These cross-feed processes are dominated by $\mathrm{Z}(\ell\ell)\mathrm{H}(2\ell + X)$, and are described in more detail in [35]. Terms that depend on κ_b are ignored, as their contribution to the ggH loop is negligible at the value of q^2 at which off-shell Higgs boson production is probed. The final term represents the background contributions in analysis region r, $\mathcal{P}_r^{\mathrm{bkg}} = b_r \rho_{\mathrm{bkg},r}$, which are independent of the coupling modifiers.

To facilitate comparisons with the previously described models, the following relation between κ_g and κ_Q for on-shell ggH production at $m_H = 125.38\,\mathrm{GeV}$ is used

$$\begin{aligned}\kappa_g^2 =& 1.035\kappa_t^2 + 0.002\kappa_b^2 - 0.038\kappa_t\kappa_b \\ &+ 0.979\kappa_Q^2 + 2.015\kappa_t\kappa_Q - 0.004\kappa_b\kappa_Q.\end{aligned} \tag{17}$$

The terms in equation (17) are derived by considering the contribution of an additional heavy quark in the ggH production loop. The calculations are performed at LO using the JHUGEN framework [87–91] interfaced with MCFM [92], following the procedure described in [87]. A K factor is applied to align the terms from the LO calculation to the corresponding terms in the ggH resolved scaling factor function in table 9. The same K factor is applied to the κ_Q-dependent terms, as these contributions are expected to receive similar higher-order corrections.

Equation (17) can be inverted to express κ_Q as a function of κ_g, κ_t, and κ_b. Substituting this expression into equation (16) allows the off-shell scaling functions to be rewritten in terms of κ_g, κ_t, and κ_b, instead of κ_Q. This procedure ensures a coherent treatment of on-shell and off-shell ggH production, and aligns the fitted POIs across all three effective coupling modifier models. However, equation (17) has two solutions as a result of the presence of the κ_Q^2 term. Among the two solutions, the positive branch is chosen as it gives $\kappa_Q = 0$ in the SM limit ($\kappa_g = \kappa_t = \kappa_b = 1$), ensuring a physical off-shell scaling behaviour.

The results from the model with the off-shell regions included are shown in purple in figure 14. It should be noted that the external constraint on $|\kappa_W|$ and $|\kappa_Z|$ has been removed. The $\mathcal{B}_{\mathrm{inv}}$ and $\mathcal{B}_{\mathrm{undet}}$ parameters are constrained to be less than 13% and 25% at the 95% CL, respectively.

The best fit values and 68% CL intervals are provided for all three effective coupling modifier fits in table 11. A notable feature of the fit that includes off-shell regions is the pronounced difference between the observed and expected upper 68% CL intervals for the κ parameters. As discussed above, on-shell signal contributions exhibit a degeneracy in the total Higgs boson decay width, where all coupling modifiers can be scaled up simultaneously to accommodate a nonzero $\mathcal{B}_{\mathrm{undet}}$. While the inclusion of off-shell processes partially lifts this degeneracy, it still leads to larger expected upper 68% CL intervals compared to the lower 68% CL intervals. However, the observed data more strongly disfavour nonzero values of $\mathcal{B}_{\mathrm{undet}}$, resulting in a tighter observed constraint on this parameter. This, in turn, yields more symmetric 68% intervals for the κ parameters, with significantly tighter upper bounds in the observed results compared to the expected.

For the two models that do not include the off-shell analysis regions, shown in blue and orange in figure 14, the best fit point for κ_W is negative. The test statistic $q(\kappa_W)$ as a function of κ_W is shown for all three models in the left plots of figure 15. While different combinations of signs for κ_W and κ_Z are



Figure 14. The measured Higgs boson coupling modifiers in the effective coupling configuration. The best fit values, and 68% and 95% CL intervals, are shown for each of the models considered. The results assuming no additional BSM contributions to the Higgs boson total decay width are shown in blue. The results for the model that allows BSM contributions but places external constraints on $|\kappa_W|$ and $|\kappa_Z|$ are shown in orange, with the constraints $|\kappa_W|, |\kappa_Z| \leqslant 1$ indicated by hatched boxes. The results for the fit in which the off-shell analysis regions are included are shown in purple. In all three models, both positive and negative values of κ_W and κ_Z are considered, while κ_t is restricted to the positive domain without loss of generality, as indicated by a hatched box. The parameters $\mathcal{B}_{inv}$ and $\mathcal{B}_{undet}$, included in the latter two models, are defined to be nonnegative, which is likewise indicated by a hatched box. The arrows shown for $\kappa_{Z\gamma}$ are used to indicate 68% CL intervals that extend beyond the plotted range.

shown, the minimum value of q across all combinations (i.e. the envelope) is used to determine the best fit point, as well as the 68% and 95% CL intervals. The preference for a negative value of κ_W is driven by the observed excess in the tH production cross section, since the tH production scaling function contains a negative interference term proportional to $\kappa_W\kappa_t$. In the effective coupling configuration, the $H \to \gamma\gamma$ decay rate scales according to κ_γ^2, such that it has no dependence on κ_W. Therefore, the preferred negative value of κ_W will not result in an excess in the $H \to \gamma\gamma$ decay rate. The second model (orange) also has a slight preference for negative values of κ_Z. The right plots in figure 15 show $q(\kappa_t)$ as a function of κ_t, demonstrating how the different combinations of signs for κ_W and κ_Z impact the results for the other coupling modifiers. For example in the first model, the additional parabola from the positive κ_W and κ_Z sign combination (blue) takes over the envelope at $\kappa_t \approx 0.9$, enlarging the upper 95% CL interval for κ_t.

For the third model, in which the off-shell $H \to ZZ \to 4\ell$ analysis is introduced into the combination, negative values of κ_W and κ_Z are excluded from the 95% CL interval. This is visible in the test statistic curves shown in the lower plots of figure 15, where only the positive sign combination contributes for $q < 10$. It can be seen from equation (16) that the scaling behaviour of certain off-shell contributions is sensitive to the relative signs of the coupling modifiers. Therefore, by including the off-shell $H \to ZZ \to 4\ell$ analysis regions in the combination, the fit is able to alleviate the degeneracy between signs, favouring positive values of κ_W and κ_Z.

9.3. Constraints on coupling modifier ratios

This section presents measurements of coupling modifier ratios,

$$\lambda_{ij} = \kappa_i/\kappa_j. \tag{18}$$

The effective coupling modifier configuration for the treatment of the ggH, $H \to \gamma\gamma$, and $H \to Z\gamma$ loop diagrams is used. A reference combined coupling modifier is defined that accounts for modifications to the total event yield of a specific combination of production process and decay channel, thereby avoiding the need for assumptions on the total Higgs boson decay width. In line with previous combinations, the reference coupling modifier is chosen to be $\kappa_{gZ} = \kappa_g\kappa_Z/\kappa_H$, as a result of the precise measurement of the ggH production process in the $H \to ZZ$ decay channel. The remaining POIs are ratios of the coupling modifiers. Both positive and negative values of λ_{WZ} and λ_{tg} are considered, while κ_{gZ} and λ_{Zg} are restricted to the positive domain without loss of generality. The measurements are not sensitive to the signs of the other coupling modifier ratios, and therefore these parameters are restricted to be positive. Neither the off-shell nor $H \to$ inv input channels are included in this fit.

The best fit values along with the 68% and 95% CL intervals are presented in figure 16, and the numerical values are provided in table 12. The best fit point for λ_{WZ} is negative. This mirrors the favoured sign combination of negative κ_W and positive κ_Z observed in the effective coupling modifier fit with no additional BSM decays (shown in blue in figure 14). The overall compatibility with the SM is measured to be $p_{SM} = 0.11$.

Additional models involving ratios of coupling modifiers are fit to test the symmetry of fermion couplings. These models probe specific BSM scenarios that predict the existence of an extended Higgs sector (discussed in section 10). In such scenarios, the couplings to up-type and down-type fermions, or to leptons and quarks, can be modified separately. Therefore, to probe such scenarios, separate coupling modifiers for up-type (κ_u) and down-type (κ_d) fermions, or for quarks (κ_q) and leptons (κ_l, where $l = e, \mu, \tau$) are introduced. A generic LO coupling modifier for the Higgs boson coupling to vector bosons κ_V, where V = W or Z, is also introduced for these models.

Figure 17 shows the results of a fit where the ratio of the couplings to up-type and down-type fermions $\lambda_{du} = \kappa_d/\kappa_u$ is determined, along with the ratio $\lambda_{Vu} = \kappa_V/\kappa_u$ and $\kappa_{uu} = \kappa_u^2/\Gamma_H$. Also shown are the results of a fit where the ratio of the couplings to leptons and quarks $\lambda_{lq} = \kappa_l/\kappa_q$ is determined, along with the ratio $\lambda_{Vq} = \kappa_V/\kappa_q$ and $\kappa_{qq} = \kappa_q^2/\Gamma_H$. In these fits, κ_{uu} and κ_{qq} are restricted to the positive domain without loss of generality, while both positive and negative values are considered for the other parameters.

The numerical values of the best fit points, along with the 68% and 95% CL intervals, are provided in table 13. The best fit points for both fits correspond to the domains where all parameters are positive. Nevertheless, the measurements are not strongly sensitive to the relative signs of λ_{du} and λ_{lq}, and negative values are included in the 68% CL intervals. The overall compatibilities with the SM are measured to be $p_{SM} = 0.04$ for both fits.

9.4. Constraints on the Higgs boson self-coupling

The Higgs boson production and decay rates can be used to constrain the Higgs boson self-coupling through modifications of the production cross sections and decay widths from NLO EW corrections [93, 94]. This analysis follows the procedure used in the combination of single and double Higgs boson production from the CMS Collaboration using LHC Run 2 data [95]. The combination detailed here updates a number of the single Higgs boson input analyses with respect to [95], most notably the VH ($H \to bb$) and t(t)H ($H \to bb$) channels. In addition, an overlap removal procedure was performed in [95] to ensure that the single and double Higgs boson analyses were statistically independent. In particular, the analysis regions targeting the STXS stage 1.2 bins for ttH production in the $H \to \gamma\gamma$ decay channel were replaced by the inclusive ttH analysis regions used in the combined ttH ($H \to \gamma\gamma$) and $HH \to \gamma\gamma bb$ measurment [96]. No such overlap removal procedure is required here, as only single Higgs boson production analyses are included in this combination.

Table 11. Best fit values and 68% CL intervals for the Higgs boson coupling modifiers in the effective coupling configuration. The results are shown for each of the models considered. The total 68% CL intervals are decomposed into their statistical and systematic components. The expected intervals are given in parentheses. The one-sided intervals represent physical boundaries in the POIs. For κ_Z in the second model, the 68% CL interval is not contiguous, and the quoted interval corresponds to the minimum containing the best fit point. The observed 68% CL interval for this parameter also includes the range $\kappa_Z > 0.98$.

	$\mathcal{B}_{BSM} = 0$			$\mathcal{B}_{inv}, \mathcal{B}_{undet}$ floating, $\lvert\kappa_V\rvert \leqslant 1$			$\mathcal{B}_{inv}, \mathcal{B}_{undet}$ floating, off-shell inc.		
Parameters	Best fit	Stat	Syst	Best fit	Stat	Syst	Best fit	Stat	Syst
κ_W	$-1.03^{+0.06}_{-0.06}$	$^{+0.04}_{-0.04}$	$^{+0.05}_{-0.05}$	$-1.00^{+0.06}$	$^{+0.04}$	$^{+0.04}$	$1.03^{+0.07}_{-0.06}$	$^{+0.07}_{-0.04}$	$^{+0.03}_{-0.04}$
	$\left(^{+0.06}_{-0.06}\right)$	$\left(^{+0.04}_{-0.04}\right)$	$\left(^{+0.04}_{-0.04}\right)$	$\left(-0.06\right)$	$\left(-0.04\right)$	$\left(-0.04\right)$	$\left(^{+0.19}_{-0.06}\right)$	$\left(^{+0.16}_{-0.04}\right)$	$\left(^{+0.08}_{-0.04}\right)$
κ_Z	$1.06^{+0.06}_{-0.06}$	$^{+0.04}_{-0.04}$	$^{+0.04}_{-0.04}$	$-0.99^{+0.06}_{-0.01}$	$^{+0.04}_{-0.01}$	$^{+0.04}_{-0.00}$	$1.09^{+0.07}_{-0.06}$	$^{+0.07}_{-0.04}$	$^{+0.03}_{-0.04}$
	$\left(^{+0.06}_{-0.06}\right)$	$\left(^{+0.04}_{-0.04}\right)$	$\left(^{+0.04}_{-0.04}\right)$	$\left(-0.06\right)$	$\left(-0.03\right)$	$\left(-0.05\right)$	$\left(^{+0.18}_{-0.06}\right)$	$\left(^{+0.16}_{-0.04}\right)$	$\left(^{+0.08}_{-0.04}\right)$
κ_t	$0.80^{+0.09}_{-0.09}$	$^{+0.06}_{-0.06}$	$^{+0.06}_{-0.06}$	$0.81^{+0.10}_{-0.09}$	$^{+0.08}_{-0.07}$	$^{+0.06}_{-0.06}$	$0.92^{+0.09}_{-0.08}$	$^{+0.07}_{-0.06}$	$^{+0.05}_{-0.05}$
	$\left(^{+0.10}_{-0.09}\right)$	$\left(^{+0.06}_{-0.06}\right)$	$\left(^{+0.08}_{-0.07}\right)$	$\left(^{+0.12}_{-0.08}\right)$	$\left(^{+0.07}_{-0.06}\right)$	$\left(^{+0.09}_{-0.05}\right)$	$\left(^{+0.20}_{-0.09}\right)$	$\left(^{+0.17}_{-0.06}\right)$	$\left(^{+0.10}_{-0.07}\right)$
κ_b	$1.06^{+0.14}_{-0.14}$	$^{+0.09}_{-0.10}$	$^{+0.10}_{-0.09}$	$0.95^{+0.09}_{-0.12}$	$^{+0.07}_{-0.08}$	$^{+0.06}_{-0.09}$	$0.96^{+0.12}_{-0.12}$	$^{+0.09}_{-0.08}$	$^{+0.08}_{-0.08}$
	$\left(^{+0.13}_{-0.12}\right)$	$\left(^{+0.09}_{-0.09}\right)$	$\left(^{+0.09}_{-0.08}\right)$	$\left(^{+0.11}_{-0.11}\right)$	$\left(^{+0.09}_{-0.07}\right)$	$\left(^{+0.06}_{-0.09}\right)$	$\left(^{+0.21}_{-0.12}\right)$	$\left(^{+0.18}_{-0.09}\right)$	$\left(^{+0.11}_{-0.08}\right)$
κ_τ	$0.93^{+0.08}_{-0.07}$	$^{+0.05}_{-0.05}$	$^{+0.06}_{-0.06}$	$0.91^{+0.07}_{-0.07}$	$^{+0.04}_{-0.05}$	$^{+0.05}_{-0.06}$	$0.91^{+0.08}_{-0.07}$	$^{+0.06}_{-0.05}$	$^{+0.05}_{-0.05}$
	$\left(^{+0.08}_{-0.07}\right)$	$\left(^{+0.05}_{-0.05}\right)$	$\left(^{+0.06}_{-0.06}\right)$	$\left(^{+0.07}_{-0.08}\right)$	$\left(^{+0.04}_{-0.04}\right)$	$\left(^{+0.06}_{-0.06}\right)$	$\left(^{+0.20}_{-0.08}\right)$	$\left(^{+0.17}_{-0.05}\right)$	$\left(^{+0.10}_{-0.06}\right)$
κ_μ	$1.11^{+0.21}_{-0.22}$	$^{+0.18}_{-0.20}$	$^{+0.09}_{-0.08}$	$1.09^{+0.19}_{-0.22}$	$^{+0.17}_{-0.20}$	$^{+0.07}_{-0.08}$	$1.09^{+0.20}_{-0.21}$	$^{+0.18}_{-0.20}$	$^{+0.09}_{-0.08}$
	$\left(^{+0.21}_{-0.24}\right)$	$\left(^{+0.19}_{-0.22}\right)$	$\left(^{+0.08}_{-0.08}\right)$	$\left(^{+0.20}_{-0.24}\right)$	$\left(^{+0.18}_{-0.22}\right)$	$\left(^{+0.09}_{-0.10}\right)$	$\left(^{+0.30}_{-0.24}\right)$	$\left(^{+0.28}_{-0.22}\right)$	$\left(^{+0.13}_{-0.08}\right)$
κ_g	$0.97^{+0.07}_{-0.07}$	$^{+0.04}_{-0.05}$	$^{+0.06}_{-0.05}$	$0.99^{+0.07}_{-0.09}$	$^{+0.05}_{-0.06}$	$^{+0.05}_{-0.06}$	$0.92^{+0.07}_{-0.06}$	$^{+0.06}_{-0.04}$	$^{+0.05}_{-0.05}$
	$\left(^{+0.07}_{-0.07}\right)$	$\left(^{+0.05}_{-0.05}\right)$	$\left(^{+0.06}_{-0.05}\right)$	$\left(^{+0.08}_{-0.07}\right)$	$\left(^{+0.04}_{-0.05}\right)$	$\left(^{+0.07}_{-0.05}\right)$	$\left(^{+0.19}_{-0.07}\right)$	$\left(^{+0.17}_{-0.05}\right)$	$\left(^{+0.09}_{-0.05}\right)$
κ_γ	$1.07^{+0.07}_{-0.07}$	$^{+0.05}_{-0.05}$	$^{+0.05}_{-0.04}$	$1.02^{+0.07}_{-0.06}$	$^{+0.06}_{-0.05}$	$^{+0.04}_{-0.04}$	$1.11^{+0.08}_{-0.07}$	$^{+0.07}_{-0.05}$	$^{+0.04}_{-0.04}$
	$\left(^{+0.06}_{-0.06}\right)$	$\left(^{+0.05}_{-0.05}\right)$	$\left(^{+0.04}_{-0.04}\right)$	$\left(^{+0.05}_{-0.06}\right)$	$\left(^{+0.04}_{-0.05}\right)$	$\left(^{+0.03}_{-0.03}\right)$	$\left(^{+0.19}_{-0.06}\right)$	$\left(^{+0.17}_{-0.05}\right)$	$\left(^{+0.09}_{-0.04}\right)$
$\kappa_{Z\gamma}$	$1.62^{+0.33}_{-0.34}$	$^{+0.30}_{-0.33}$	$^{+0.13}_{-0.09}$	$1.57^{+0.30}_{-0.34}$	$^{+0.28}_{-0.31}$	$^{+0.11}_{-0.13}$	$1.62^{+0.33}_{-0.34}$	$^{+0.31}_{-0.33}$	$^{+0.11}_{-0.09}$
	$\left(^{+0.37}_{-0.58}\right)$	$\left(^{+0.36}_{-0.57}\right)$	$\left(^{+0.09}_{-0.09}\right)$	$\left(^{+0.42}_{-0.58}\right)$	$\left(^{+0.41}_{-0.57}\right)$	$\left(^{+0.11}_{-0.11}\right)$	$\left(^{+0.45}_{-0.60}\right)$	$\left(^{+0.44}_{-0.59}\right)$	$\left(^{+0.13}_{-0.08}\right)$
$\mathcal{B}_{inv}$		—		$0.07^{+0.04}_{-0.06}$	$^{+0.02}_{-0.04}$	$^{+0.03}_{-0.05}$	$0.05^{+0.04}_{-0.04}$	$^{+0.02}_{-0.02}$	$^{+0.03}_{-0.03}$
		—		$\left(^{+0.04}\right)$	$\left(^{+0.02}\right)$	$\left(^{+0.04}\right)$	$\left(^{+0.04}\right)$	$\left(^{+0.02}\right)$	$\left(^{+0.04}\right)$
$\mathcal{B}_{undet}$		—		$0.00^{+0.11}$	$^{+0.07}$	$^{+0.07}$	$0.00^{+0.11}$	$^{+0.10}$	$^{+0.02}$
		—		$\left(^{+0.10}\right)$	$\left(^{+0.06}\right)$	$\left(^{+0.08}\right)$	$\left(^{+0.29}\right)$	$\left(^{+0.26}\right)$	$\left(^{+0.12}\right)$



Figure 15. Scans of the test statistic q as a function of κ_W (left) and κ_t (right) for the three effective coupling modifier models. The upper row shows the model that assumes no additional BSM contributions to the Higgs boson total decay width. The middle row shows the model that introduces BSM contributions, but places an external constraint on $|\kappa_W|$ and $|\kappa_Z|$, and the lower row shows the model in which the off-shell $H \to ZZ \to 4\ell$ analysis regions are included. The different coloured lines indicate the value of q for different combinations of signs for κ_W and κ_Z. The solid black line shows the minimum value of q across all combinations of signs, which is used to determine the best fit point and the 68% and 95% CL intervals. The scan in the middle left plot is truncated because of the external constraint of $|\kappa_W| \leqslant 1$, which is indicated by the hatched boxes. The plots in the lower row are shown on a different y-axis scale to emphasize the exclusion of the sign combinations that include negative values of κ_W and κ_Z.

The coupling modifier of the trilinear Higgs boson self-coupling with respect to the SM prediction is defined as $\kappa_\lambda = \lambda_3/\lambda_3^{SM}$. Deviations from the SM Higgs boson production and decay rates are parametrized using a three-parameter model: κ_λ, κ_F, and κ_V. Here, κ_F and κ_V are the LO coupling



Figure 16. The measured parameters in the ratios of coupling modifiers fit. The thick (thin) black lines indicate the 68% (95%) CL intervals, around the best fit points (empty circles). For this model, both positive and negative values of λ_{WZ} and λ_{tg} are considered, while κ_{gZ} and λ_{Zg} are restricted to the positive domain without loss of generality, as indicated by the hatched box.

Table 12. Best fit values and 68% CL intervals for the coupling modifier ratios. The total 68% CL intervals are decomposed into their statistical and systematic components. The expected intervals are given in parentheses.

Parameters	Best fit	Stat	Syst
κ_{gZ}	$1.00^{+0.05}_{-0.05}$	$^{+0.04}_{-0.04}$	$^{+0.04}_{-0.03}$
	$\left(^{+0.05}_{-0.05}\right)$	$\left(^{+0.04}_{-0.04}\right)$	$\left(^{+0.04}_{-0.03}\right)$
λ_{Zg}	$1.10^{+0.09}_{-0.08}$	$^{+0.07}_{-0.06}$	$^{+0.07}_{-0.06}$
	$\left(^{+0.09}_{-0.08}\right)$	$\left(^{+0.06}_{-0.06}\right)$	$\left(^{+0.06}_{-0.06}\right)$
λ_{tg}	$0.83^{+0.09}_{-0.09}$	$^{+0.07}_{-0.07}$	$^{+0.07}_{-0.06}$
	$\left(^{+0.10}_{-0.10}\right)$	$\left(^{+0.07}_{-0.07}\right)$	$\left(^{+0.08}_{-0.07}\right)$
λ_{WZ}	$-0.97^{+0.06}_{-0.05}$	$^{+0.04}_{-0.04}$	$^{+0.04}_{-0.04}$
	$\left(^{+0.06}_{-0.06}\right)$	$\left(^{+0.04}_{-0.04}\right)$	$\left(^{+0.04}_{-0.04}\right)$
$\lambda_{\gamma Z}$	$1.01^{+0.06}_{-0.06}$	$^{+0.05}_{-0.05}$	$^{+0.03}_{-0.03}$
	$\left(^{+0.06}_{-0.06}\right)$	$\left(^{+0.05}_{-0.05}\right)$	$\left(^{+0.03}_{-0.03}\right)$
$\lambda_{Z\gamma Z}$	$1.53^{+0.31}_{-0.33}$	$^{+0.27}_{-0.32}$	$^{+0.15}_{-0.09}$
	$\left(^{+0.37}_{-0.61}\right)$	$\left(^{+0.35}_{-0.60}\right)$	$\left(^{+0.11}_{-0.09}\right)$
λ_{bZ}	$0.99^{+0.11}_{-0.10}$	$^{+0.08}_{-0.08}$	$^{+0.07}_{-0.07}$
	$\left(^{+0.11}_{-0.10}\right)$	$\left(^{+0.08}_{-0.08}\right)$	$\left(^{+0.07}_{-0.07}\right)$
$\lambda_{\tau Z}$	$0.88^{+0.07}_{-0.07}$	$^{+0.05}_{-0.04}$	$^{+0.05}_{-0.05}$
	$\left(^{+0.08}_{-0.08}\right)$	$\left(^{+0.05}_{-0.05}\right)$	$\left(^{+0.06}_{-0.06}\right)$
$\lambda_{\mu Z}$	$1.04^{+0.19}_{-0.21}$	$^{+0.18}_{-0.19}$	$^{+0.06}_{-0.08}$
	$\left(^{+0.21}_{-0.23}\right)$	$\left(^{+0.19}_{-0.22}\right)$	$\left(^{+0.08}_{-0.09}\right)$

modifiers for the Higgs boson coupling to fermions and vector bosons, respectively. In the following, both κ_F and κ_V are restricted to the positive domain.

The NLO EW corrections from κ_λ are dependent on the kinematic properties of H production. The variations of the production cross sections as a function of κ_λ have been derived at the granularity of the STXS stage 1.2 binning scheme [97]. This parametrization is applied to the input channels that split the signal contributions into the STXS bins. Inclusive scaling functions are used for all other channels.

The test statistic as a function of κ_λ is shown in figure 18. The blue lines show the results when the other two coupling modifiers are fixed to their SM values, $\kappa_F = \kappa_V = 1$. The measurement is performed in the region $\kappa_\lambda \in [-15, 20]$, beyond which the model used here is no longer valid as NNLO effects



Figure 17. The measured parameters in the fits probing the symmetry of fermion couplings. The left plot shows the results for the model that probes the ratio of couplings to up-type and down-type fermions. In this fit, both positive and negative values of λ_{du} and λ_{Vu} are considered. The right plot shows the results for the model that probes the ratio of couplings to leptons and quarks, where both positive and negative values of λ_{lq} and λ_{Vq} are considered. In both plots, the thick (thin) black lines indicate the 68% (95%) CL intervals, around the best fit points (empty circles). The hatched boxes indicate parameters that are restricted to the positive domain.

Table 13. Best fit values and 68% CL intervals for the three parameter models that probe the symmetry of fermion couplings. The total 68% CL intervals are decomposed into their statistical and systematic components. The expected intervals are given in parentheses. For λ_{du} and λ_{lq} the 68% CL intervals are not contiguous, and the quoted intervals correspond to the minima containing the best fit points. The 68% CL intervals also include the ranges $-1.11 < \lambda_{du} < -0.99$ and $-1.04 < \lambda_{lq} < -0.91$ ($-1.09 < \lambda_{du} < -0.97$ and $-1.07 < \lambda_{lq} < -0.93$) for the observed (expected) case.

Parameters	Best fit	Stat	Syst
κ_{uu}	$0.85^{+0.08}_{-0.11}$ $\left(^{+0.09}_{-0.13}\right)$	$^{+0.05}_{-0.09}$ $\left(^{+0.06}_{-0.10}\right)$	$^{+0.06}_{-0.06}$ $\left(^{+0.07}_{-0.08}\right)$
λ_{du}	$1.02^{+0.07}_{-0.06}$ $\left(^{+0.07}_{-0.06}\right)$	$^{+0.05}_{-0.04}$ $\left(^{+0.05}_{-0.04}\right)$	$^{+0.05}_{-0.05}$ $\left(^{+0.05}_{-0.04}\right)$
λ_{Vu}	$1.15^{+0.10}_{-0.06}$ $\left(^{+0.09}_{-0.06}\right)$	$^{+0.08}_{-0.05}$ $\left(^{+0.07}_{-0.04}\right)$	$^{+0.06}_{-0.04}$ $\left(^{+0.06}_{-0.04}\right)$
κ_{qq}	$0.88^{+0.07}_{-0.06}$ $\left(^{+0.08}_{-0.08}\right)$	$^{+0.04}_{-0.05}$ $\left(^{+0.05}_{-0.05}\right)$	$^{+0.05}_{-0.04}$ $\left(^{+0.06}_{-0.06}\right)$
λ_{lq}	$0.97^{+0.07}_{-0.06}$ $\left(^{+0.07}_{-0.07}\right)$	$^{+0.05}_{-0.05}$ $\left(^{+0.05}_{-0.04}\right)$	$^{+0.04}_{-0.05}$ $\left(^{+0.05}_{-0.05}\right)$
λ_{Vq}	$1.12^{+0.06}_{-0.06}$ $\left(^{+0.06}_{-0.05}\right)$	$^{+0.05}_{-0.04}$ $\left(^{+0.05}_{-0.04}\right)$	$^{+0.04}_{-0.04}$ $\left(^{+0.04}_{-0.04}\right)$

become important. In this scenario, the measured value is $\kappa_\lambda = 2.1^{+4.0}_{-3.2}$. Compared to the result for single Higgs boson production in [95], which measured $\kappa_\lambda = 5.8^{+3.5}_{-4.2}$, the constraint is slightly improved. This improvement is a result of the updated input analyses, along with the fact that no overlap removal procedure with di-Higgs analyses has been performed here. The shift in the best fit value relative to [95] is largely driven by the updated VH ($H \to bb$) inputs, as well as the different analysis regions used to target ttH production in the $H \to \gamma\gamma$ decay channel. This shift is within the observed 68% CL interval.

The assumptions on the tree-level couplings can be relaxed by allowing κ_F, κ_V, or both to vary in addition to the Higgs boson self-coupling. Figure 19 shows the test statistic as a function of κ_λ-vs-κ_F, and κ_λ-vs-κ_V, where the other tree-level coupling modifier is fixed to 1. The orange lines in figure 18 show the test statistic as a function of κ_λ, when both κ_F and κ_V are profiled. In this scenario, the measured value is $\kappa_\lambda = 0.1^{+6.3}_{-6.0}$. It is not possible to extract a 95% CL interval within the range of validity of the model. Nevertheless, it is encouraging that the available data can simultaneously constrain κ_F, κ_V, and κ_λ, using only analyses that measure single Higgs boson production.

The numerical values of the best fit points, along with the 68% and 95% CL intervals, are provided in Table 14 for the different assumptions on the values of κ_F and κ_V. The p_{SM} values are also provided for each assumption, giving a measure of the compatibility with the SM hypothesis.

Table 14. Best fit values, 68% CL, and 95% CL intervals for κ_λ, with different assumptions on the values of κ_F and κ_V. The expected intervals are given in parentheses. The p_{SM} values, which represent the compatibility with the SM hypothesis, are also provided. A 95% CL interval for the fit in which κ_F and κ_V are floating cannot be extracted within the range of validity for this model.

Assumption	Best fit κ_λ	95% CL interval	p_{SM}
$\kappa_F = \kappa_V = 1$	$2.1^{+4.0}_{-3.2}$ $\left(^{+6.6}_{-3.3}\right)$	$[-3.3, 9.6]$ $\left([-4.6, 11.6]\right)$	0.74
$\kappa_V = 1$, κ_F floating	$2.5^{+3.5}_{-3.0}$ $\left(^{+6.7}_{-3.4}\right)$	$[-2.7, 9.3]$ $\left([-4.7, 11.6]\right)$	0.31
$\kappa_F = 1$, κ_V floating	$-2.2^{+6.6}_{-3.5}$ $\left(^{+6.7}_{-4.4}\right)$	$[-8.5, 15.7]$ $\left([-6.5, 13.3]\right)$	0.19
κ_F, κ_V floating	$0.1^{+6.3}_{-6.0}$ $\left(^{+6.9}_{-6.0}\right)$	— —	0.30



Figure 18. Profile likelihood ratio scans as functions of κ_λ for the observed data (solid lines). The expected results assuming an SM Higgs boson derived using an Asimov data set with $\kappa_\lambda = 1$ are shown by the dashed lines. The blue lines represent the case where κ_F and κ_V are fixed to 1. The orange lines represent the case where κ_F and κ_V are profiled.

10. Interpretation of measurements using benchmark two-Higgs-doublet models

The models described in section 9.3, which test the symmetry of the Higgs boson couplings to fermions, can also be interpreted in the context of UV-complete extensions of the SM that contain a second Higgs doublet (two-Higgs-doublet models, 2HDMs) [98–100]. Here, the term UV-complete refers to models that remain valid up to arbitrarily high energy scales. In a 2HDM, the SM Higgs sector—which contains one doublet of complex scalar fields Φ_1—is extended by the addition of a second complex scalar doublet Φ_2. Electroweak symmetry breaking yields five physical scalar Higgs fields: two neutral *CP*-even (h and $\mathcal{H}$), one neutral *CP*-odd (A), and two charged scalars ($\mathcal{H}^\pm$).

Only 2HDMs with *CP* conservation are considered. In addition, the models require a (softly broken) $\mathbb{Z}_2$ discrete symmetry to prevent tree-level flavour changing neutral currents [101, 102], which are strongly constrained experimentally. Four 2HDM types are possible under these conditions, referred to as the Type-I, Type-II, Lepton-specific, and Flipped scenarios, which differ in the way the two Higgs doublets couple to fermions:

- Type I: all fermions couple to just one of the Higgs doublets (Φ_2 by convention).



Figure 19. Profile likelihood ratio scans as a function of κ_λ and κ_F (upper) and as a function of κ_λ and κ_V (lower). The blue colour scale shows the value of q at each point in the scan. The black marker, and solid and dashed lines, show the best fit point, and the 68% and 95% CL contours, respectively. The red marker corresponds to the SM prediction.

- Type II: up-type quarks couple to Φ_2, while down-type quarks and right-handed leptons couple to Φ_1.
- Lepton-specific: quarks couple to Φ_2, and the right-handed leptons couple to Φ_1.
- Flipped: up-type quarks and right-handed leptons couple to Φ_2, while the down-type quarks couple to Φ_1.

Each of the 2HDM scenarios contains seven free parameters. These are the four Higgs boson masses (m_h, m_H, m_A, and $m_{H^\pm}$); the mixing angles α and β, where α diagonalizes the mass-squared matrix of the neutral scalars, and β diagonalizes the mass-squared matrix of the charged scalars and pseudoscalars;

and the soft $\mathbb{Z}_2$ symmetry breaking parameter m_{12}^2. The vacuum expectation values of the two Higgs doublets, v_1 and v_2, are related by $\tan\beta = v_2/v_1$. Under the additional assumption that the observed Higgs boson with a mass of 125.38 GeV is the lightest *CP*-even scalar h, the predicted rates for its production and decay are, at LO, only sensitive to the free parameters α and β. These two parameters are substituted by $\cos(\beta-\alpha)$ and $\tan\beta$, without loss of generality.

The 2HDM interpretations in this section are derived in the decoupling limit [99], where the observed Higgs boson is much lighter than the scale Λ_{2HDM} associated with the heavier scalar states: $m_{\mathcal{H}}$, m_A, $m_{\mathcal{H}^\pm} \sim \Lambda_{\text{2HDM}} \gg m_h$. The decoupling limit implies the alignment limit, $\cos(\beta-\alpha) \ll 1$, such that the couplings of the lightest *CP*-even scalar h approach those of the SM Higgs boson. Table 15 shows the LO coupling modifiers to vector bosons, quarks, and leptons, in the vicinity of the alignment limit, for each of the 2HDM scenarios considered. In all scenarios, the coupling of the Higgs boson to vector bosons is modified by a factor $\sin(\beta-\alpha)$, while the couplings to fermions depend on the scenario.

An additional interpretation is provided in the context of the minimal supersymmetric extension of the SM (MSSM) [103–106]. In the MSSM, the Higgs sector is a 2HDM of Type II. The additional constraints that arise from the nontrivial fermion-boson symmetry in the supersymmetric theory fix all mass relations between the Higgs bosons and the angle α at tree level. This leaves only two free parameters, usually chosen to be m_A and $\tan\beta$, to fully constrain the MSSM Higgs sector.

The specific benchmark scenario considered here is the hMSSM [107, 108]. As with the generic 2HDM scenarios, the observed Higgs boson is identified with the lightest *CP*-even scalar h. Its mass is set equal to the experimentally measured value of 125.38 GeV, which fixes the values of the dominant radiative corrections to the *CP*-even Higgs boson mass matrix. As a result, the couplings of the Higgs boson to other particles can be expressed in terms of only m_A and $\tan\beta$ [109], following the relations given in table 15. These are completed by the definitions of s_{u} and s_{d} in equations (19) and (20), respectively,

$$s_{\text{u}} = \frac{1}{\sqrt{1+\frac{\left(m_A^2+m_{\text{Z}}^2\right)^2\tan^2\beta}{\left(m_{\text{Z}}^2+m_A^2\tan^2\beta-m_{\text{H}}^2(1+\tan^2\beta)\right)^2}}} \tag{19}$$

$$s_{\text{d}} = s_{\text{u}}\frac{\left(m_A^2+m_{\text{Z}}^2\right)\tan\beta}{m_{\text{Z}}^2+m_A^2\tan^2\beta-m_{\text{H}}^2(1+\tan^2\beta)}. \tag{20}$$

Although many other MSSM benchmark scenarios have been defined [110, 111], the lack of analytic expressions for the Higgs boson couplings renders these models technically more challenging to consider, and they are beyond the scope of this paper.

To set constraints on the UV-complete model parameters, the test statistic must be evaluated over a grid of points in the relevant parameter space. The relations between the Higgs boson couplings and the 2HDM and hMSSM parameters result in complicated likelihood surfaces with flat directions and multiple minima, which make it difficult to perform a direct fit of the model parameters. Instead, three-dimensional likelihood scans are performed using the coupling modifier ratio parametrizations that probe the symmetry of fermion couplings, described at the end of section 9.3 (with suitable modifications to the lepton coupling modifiers to describe the Flipped 2HDM scenario). The resulting three-dimensional likelihood surfaces are subsequently projected onto the 2D planes of $\cos(\beta-\alpha)$ and $\tan\beta$ for the 2HDM scenarios, or m_A and $\tan\beta$ for the hMSSM scenario, following the procedure described below.

A test statistic is defined over the three-dimensional grid of points, for example in the parametrization involving λ_{du}, λ_{Vu}, and κ_{uu},

$$q(\lambda_{\text{du}},\lambda_{\text{Vu}},\kappa_{\text{uu}}) = -2\ln\frac{L(\lambda_{\text{du}},\lambda_{\text{Vu}},\kappa_{\text{uu}})}{L\left(\hat{\lambda}_{\text{du}},\hat{\lambda}_{\text{Vu}},\hat{\kappa}_{\text{uu}}\right)}, \tag{21}$$

where $\hat{\lambda}_{\text{du}}$, $\hat{\lambda}_{\text{Vu}}$, and $\hat{\kappa}_{\text{uu}}$ are the values of the POIs that maximize the likelihood, and the explicit dependence on the NPs has been dropped for simplicity. An interpolation procedure is used to obtain a continuous test statistic function over the three-dimensional space. Using the relations in table 15, the parameters can be expressed as functions of $\cos(\beta-\alpha)$ and $\tan\beta$. For example, in the Type-II 2HDM

Table 15. Modifications to the couplings of the Higgs boson to vector bosons, up-type quarks, down-type quarks, and charged leptons, in the 2HDM and hMSSM scenarios. The modifications act as multiplicative factors to the SM expectations. The expressions for s_u and s_d, which enter the hMSSM terms, are given in equations (19) and (20), respectively.

	2HDM				
Coupling	Type I	Type II	Lepton-specific	Flipped	hMSSM
W, Z	$\sin(\beta-\alpha)$	$\sin(\beta-\alpha)$	$\sin(\beta-\alpha)$	$\sin(\beta-\alpha)$	$\frac{s_d+s_u\tan\beta}{\sqrt{1+\tan^2\beta}}$
u, c, t	$\cos\alpha/\sin\beta$	$\cos\alpha/\sin\beta$	$\cos\alpha/\sin\beta$	$\cos\alpha/\sin\beta$	$s_u\frac{\sqrt{1+\tan^2\beta}}{\tan\beta}$
d, s, b	$\cos\alpha/\sin\beta$	$-\sin\alpha/\cos\beta$	$\cos\alpha/\sin\beta$	$-\sin\alpha/\cos\beta$	$s_d\sqrt{1+\tan^2\beta}$
e, μ, τ	$\cos\alpha/\sin\beta$	$-\sin\alpha/\cos\beta$	$-\sin\alpha/\cos\beta$	$\cos\alpha/\sin\beta$	$s_d\sqrt{1+\tan^2\beta}$

scenario,

$$\lambda_{du}=\frac{\kappa_d(\alpha,\beta)}{\kappa_u(\alpha,\beta)}=\frac{-\sin\alpha/\cos\beta}{\cos\alpha/\sin\beta}=\frac{\sqrt{1-\cos^2(\beta-\alpha)}-\cos(\beta-\alpha)\tan\beta}{\sqrt{1-\cos^2(\beta-\alpha)}+\cos(\beta-\alpha)/\tan\beta}. \tag{22}$$

These mapping functions (provided as supplementary material [41]) are then used to obtain the value of q as a function of $\cos(\beta-\alpha)$ and $\tan\beta$, or m_A and $\tan\beta$, for the 2HDM and hMSSM scenarios, respectively.

A second quantity q' is defined as,

$$q'=-2\ln\frac{L\left(\hat{\lambda}_{du},\hat{\lambda}_{Vu},\hat{\kappa}_{uu}\right)}{L_{max}}, \tag{23}$$

where L_{max} is the maximum likelihood value attained in the planes of $\cos(\beta-\alpha)$ and $\tan\beta$, or m_A and $\tan\beta$. The allowed regions of parameter space are determined as the points in the plane for which the difference between q and q' (Δq) is less than 5.99. This value corresponds to the 95% CL region, assuming Δq follows a χ^2 distribution with two degrees of freedom. The procedure is repeated using the model with the parameters λ_{lq}, λ_{Vq}, and κ_{qq} to determine the allowed regions for the Lepton-specific 2HDM scenario.

Figure 20 shows the observed and expected 95% CL exclusion regions for the different 2HDM scenarios. The red lines at $\cos(\beta-\alpha)=0$ indicate the alignment limit, where the Higgs boson couplings coincide with the SM predictions. In the Type-I 2HDM scenario, the observed 95% CL region has an asymmetric shape that favours negative values of $\cos(\beta-\alpha)$, excluding points along the alignment limit for large values of $\tan\beta$. This behaviour is consistent with the observed values of λ_{Vu} and κ_{uu}, which lie about two standard deviations above and below their SM expectations, respectively (shown in figure 17). In the Type-II, Lepton-specific, and Flipped 2HDM scenarios, the lobe-like structures appearing in the upper-right corner of the plots arise from negative values of λ_{du} and λ_{lq}, which remain allowed within the 95% CL regions with the current sensitivity. The constraints for the hMSSM scenario are also shown in figure 20. At low values of $\tan\beta$, the observed exclusion region exhibits a point of inflection relative to the expected. This feature is again consistent with the observed value of κ_{uu} lying below the SM expectation.

The allowed parameter spaces in the 2HDM and hMSSM scenarios have been significantly reduced compared with the previous result from the CMS Collaboration using data collected in 2016 only [31]. These constraints are complementary to those obtained from direct searches for additional Higgs bosons at the LHC, for example in [112–114]. A comparison of the hMSSM constraints to those from direct searches is provided as supplementary material [41], showing that the indirect constraints from the combination provide unique sensitivity in the intermediate $\tan\beta$ region ($2<\tan\beta<7$).

11. Interpretation of measurements using SMEFT

A consistent framework for introducing perturbations to the SM Lagrangian arising from the presence of new particles at an energy scale Λ is provided by SMEFT [24]. The scale Λ is assumed to be much higher than the EW scale, $\Lambda \gg v$. The perturbations are represented by higher-dimensional operators, which encapsulate the effects of short-distance BSM physics through contact interactions between SM fields.



Figure 20. Constraints in the $\tan\beta$ vs $\cos(\beta-\alpha)$ plane for the Type-I, Type-II, lepton-specific, and Flipped 2HDM scenarios, and constraints in the $\tan\beta$ vs. m_A plane for the hMSSM scenario. The grey regions, bounded by the solid black lines, represent the regions of parameter space that are excluded at the 95% CL, given the data observed. The dashed black lines indicate the expected boundaries of the 95% CL exclusion regions, for an SM Higgs boson. The dashed red lines in the 2HDM plots indicate the alignment limit, $\cos(\beta-\alpha)=0$, where the Higgs boson couplings coincide with the SM predictions.

In contrast to the results presented in section 10, a key feature of SMEFT is that it does not rely on the realization of any specific BSM scenario. Instead, it serves as a model-agnostic framework to indirectly probe BSM physics by examining deviations in SM processes. Under the assumption that the BSM

particles are too heavy to be produced on-shell at the LHC, and that there are no additional light particles, the SM Lagrangian can be expanded in powers of $1/\Lambda$ as

$$\mathcal{L}_{\text{SMEFT}} = \mathcal{L}_{\text{SM}} + \sum_{d}^{N} \sum_{j \in \mathcal{O}^{(d)}} \frac{c_j^{(d)}}{\Lambda^{d-4}} \mathcal{O}_j^{(d)}, \tag{24}$$

where $\mathcal{O}_j^{(d)}$ are operators of mass dimension $d \geqslant 5$, and $c_j^{(d)}$ are Wilson coefficients (WCs) controlling the size of the effect of each SMEFT operator. Under the SM expectation, the WC values are zero, whereas values significantly different from zero would indicate BSM contributions in the interaction described by the corresponding operator.

Only the effects of dimension-6 operators are considered in this interpretation. Operators with odd dimension d are not considered as they violate lepton number or baryon-minus-lepton number conservation, and higher even-dimension terms are suppressed by factors of $1/\Lambda^4$ or more. After demanding that the operators maintain the gauge symmetries conserved by the SM $(SU(3)_c \times SU(2)_L \times U(1)_Y)$, there are 2499 dimension-6 operators that together form an independent basis. The Warsaw basis implementation [115] is used to align with many other EFT constraints that have been set based on data from high-energy physics experiments.

With the current amount of data collected by the CMS experiment, it is not possible to constrain such a large number of operators simultaneously. However, the number of independent operators can be reduced significantly by imposing a flavour symmetry. In this interpretation, the `topU3l` symmetry of [116] is used. This symmetry introduces a set of operators for the heavy quarks (t and b) separate from the light quarks (u, d, c, and s), and has a common set of operators for all lepton flavours. This reduces the basis to 182 operators. In addition, the kinematic variables on which the STXS measurements are based cannot distinguish between *CP*-conserving and *CP*-violating effects. As a result, only the *CP*-conserving variants of the operators are considered, which leaves a total of 129 operators.

The Higgs boson combination detailed in this paper is sensitive to 43 of the 129 operators that satisfy the above constraints. Maximum likelihood fits are performed to extract constraints on the corresponding 43 WCs, which are listed in table 16. A value of $\Lambda = 1\,\text{TeV}$ is adopted for all results in this paper. The results for an alternative scale $\Lambda = X$ can be obtained by scaling the numerical values by a factor $(X/1\,\text{TeV})^2$.

11.1. Derivation of the SMEFT parametrization

The Higgs boson production cross sections and branching fractions are parametrized as functions of the SMEFT WCs. These scaling functions enter equation (5) as signal strength modifiers $\mu^{if}(\vec{c})$. The cross section scaling functions are derived at the granularity of the STXS stage 1.2 binning scheme. This facilitates tighter constraints on the WCs due to modifications of the kinematic properties of Higgs boson production. The VH ($\text{H} \to \text{bb}$) and t(t)H ($\text{H} \to \text{bb}$) input analyses include an additional kinematic splitting of the signal contributions in the likelihood, at $p_{\text{T}}^{\text{V}} = 400\,\text{GeV}$ for the VH leptonic production processes, and at $p_{\text{T}}^{\text{V}} = 450\,\text{GeV}$ for the ttH process. The SMEFT parametrization is derived for these finer binned kinematic regions, and applied to the VH ($\text{H} \to \text{bb}$) and t(t)H ($\text{H} \to \text{bb}$) inputs. Inclusive production process scaling functions are used for analysis regions that do not split the signal into the STXS stage 1.2 bins, namely VBF ($\text{H} \to \text{bb}$), $\text{H} \to \mu\mu$, and $\text{H} \to \text{Z}\gamma \to \ell\ell\gamma$. In this interpretation, SMEFT modifications are not considered for background processes. This assumption follows the fact that the input analyses generally have small background contributions, or the background rates are estimated directly from data in dedicated CRs.

Analytic calculations of the SMEFT contributions at NLO EW accuracy are used for the $\text{H} \to \gamma\gamma$ [117] and $\text{H} \to \text{Z}\gamma$ [118] decay rates. The SMEFT contributions for all other production and decay processes are derived numerically using two universal FeynRules output (UFO) models in MadGraph5_aMC@NLO [66]. Loop-level contributions are calculated for the ggH, ggZH, and $\text{H} \to \text{gg}$ processes using SMEFT@NLO [119], where $\text{H} \to \text{gg}$ refers to the Higgs boson decay to a pair of gluons. Tree-level contributions are calculated for all remaining processes with SMEFTsim 3.0 [116]. The `topU3l` flavour symmetry is used with the $(G_{\text{F}}, m_{\text{Z}}, m_{\text{W}})$ input parameter scheme, where $G_{\text{F}} = 1.16638 \times 10^{-5}\,\text{GeV}^{-2}$ is the Fermi constant, and $m_{\text{Z}} = 91.1876\,\text{GeV}$ and $m_{\text{W}} = 80.379\,\text{GeV}$ are the Z and W boson masses, respectively [120]. The parametrization is computed for a Higgs boson mass of 125.0 GeV. While this differs from the m_{H} value of 125.38 GeV used in the extraction of the results, the impact on the derived parametrization is negligible. The PDFs are taken from the NNPDF 3.1 set [72], where the 4FS

Table 16. A list of the 43 WCs, c_j, considered in the SMEFT interpretation, and the corresponding dimension-6 operators $\mathcal{O}_j$. The coefficients are grouped into terms of a similar structure in the SM Lagrangian expansion. The operators follow the same notation as [116], where (q, u, d) denote the quark fields of the first two generations, (Q, t, b) quark fields of the third generation, and (ℓ, e, ν) lepton fields of all three generations. The Higgs doublet is represented by H, a covariant derivative by D, and a vector boson field by $X = G, W, B$. Fermion fields are represented by ψ, with L (R) indicating left-handed (right-handed) fermion fields. The d'Alembertian operator is denoted by $\Box$.

Group	WC	Operator
X^3	c_W	$\epsilon^{ijk} W^{i\nu}_{\mu} W^{j\rho}_{\nu} W^{k\mu}_{\rho}$
	c_G	$f^{abc} G^{a\nu}_{\mu} G^{b\rho}_{\nu} G^{c\mu}_{\rho}$
$H^4 D^2$	$c_{H\Box}$	$(H^\dagger H)\Box(H^\dagger H)$
	c_{HD}	$(D^\mu H^\dagger H)(H^\dagger D^\mu H)$
$X^2 H^2$	c_{HG}	$H^\dagger H G^{a}_{\mu\nu} G^{a,\mu\nu}$
	c_{HW}	$H^\dagger H W^{i}_{\mu\nu} W^{i,\mu\nu}$
	c_{HB}	$H^\dagger H B_{\mu\nu} B^{\mu\nu}$
	c_{HWB}	$H^\dagger H W^{i}_{\mu\nu} B^{\mu\nu}$
$\psi^2 H^3$	$\mathrm{Re}(c_{eH})$	$(H^\dagger H)(\bar{l} e H)$
	$\mathrm{Re}(c_{bH})$	$(H^\dagger H)(\bar{Q} \tilde{H} t)$
	$\mathrm{Re}(c_{tH})$	$(H^\dagger H)(\bar{Q} H b)$
$\psi^2 XH$	$\mathrm{Re}(c_{tG})$	$(\bar{Q}\sigma^{\mu\nu} T^a t)\tilde{H} G^{a}_{\mu\nu}$
	$\mathrm{Re}(c_{tW})$	$(\bar{Q}\sigma^{\mu\nu} t)\sigma^i \tilde{H} W^{i}_{\mu\nu}$
	$\mathrm{Re}(c_{tb})$	$(\bar{Q}\sigma^{\mu\nu} t)\tilde{H} B_{\mu\nu}$
	$\mathrm{Re}(c_{bG})$	$(\bar{Q}\sigma^{\mu\nu} T^a b) H G^{a}_{\mu\nu}$
	$\mathrm{Re}(c_{bW})$	$(\bar{Q}\sigma^{\mu\nu} b)\sigma^i H W^{i}_{\mu\nu}$
$\psi^2 H^2 D$	$c^{(1)}_{Hl}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}\gamma^\mu l)$
	$c^{(3)}_{Hl}$	$(H^\dagger i \overleftrightarrow{D}^i_\mu H)(\bar{l}\sigma^i\gamma^\mu l)$
	$c^{(1)}_{Hq}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}\gamma^\mu q)$
	$c^{(3)}_{Hq}$	$(H^\dagger i \overleftrightarrow{D}^i_\mu H)(\bar{q}\sigma^i\gamma^\mu q)$
	$c^{(1)}_{HQ}$	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{Q}\gamma^\mu Q)$
	$c^{(3)}_{HQ}$	$(H^\dagger i \overleftrightarrow{D}^i_\mu H)(\bar{Q}\sigma^i\gamma^\mu Q)$
	c_{He}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}\gamma^\mu e)$
	c_{Hu}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}\gamma^\mu u)$
	c_{Hd}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}\gamma^\mu d)$
	$\mathrm{Re}(c_{Htb})$	$i(H^\dagger D_\mu H)(\bar{t}\gamma^\mu b)$
	c_{Ht}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{t}\gamma^\mu t)$
	c_{Hb}	$(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{b}\gamma^\mu b)$
$(\bar{L}L)(\bar{L}L)$	$c^{(1)}_{ll}$	$(\bar{l}\gamma_\mu l)(\bar{l}\gamma^\mu l)$
	$c^{(11)}_{Qq}$	$(\bar{Q}\gamma_\mu Q)(\bar{q}\gamma^\mu q)$
	$c^{(18)}_{Qq}$	$(\bar{Q}T^a\gamma_\mu Q)(\bar{q}T^a\gamma^\mu q)$
	$c^{(31)}_{Qq}$	$(\bar{Q}\sigma^i\gamma_\mu Q)(\bar{q}\sigma^i\gamma^\mu q)$
	$c^{(38)}_{Qq}$	$(\bar{Q}\sigma^i T^a\gamma_\mu Q)(\bar{q}\sigma^i T^a\gamma^\mu q)$
$(\bar{R}R)(\bar{R}R)$	$c^{(1)}_{tu}$	$(\bar{t}\gamma_\mu t)(\bar{u}\gamma^\mu u)$
	$c^{(8)}_{tu}$	$(\bar{t}T^a\gamma_\mu t)(\bar{u}T^a\gamma^\mu u)$
	$c^{(1)}_{td}$	$(\bar{t}\gamma_\mu t)(\bar{d}\gamma^\mu d)$
	$c^{(8)}_{td}$	$(\bar{t}T^a\gamma_\mu t)(\bar{d}T^a\gamma^\mu d)$
$(\bar{L}L)(\bar{R}R)$	$c^{(1)}_{qt}$	$(\bar{q}\gamma_\mu q)(\bar{t}\gamma^\mu t)$
	$c^{(8)}_{qt}$	$(\bar{q}T^a\gamma_\mu q)(\bar{t}T^a\gamma^\mu t)$
	$c^{(1)}_{Qu}$	$(\bar{Q}\gamma_\mu Q)(\bar{u}\gamma^\mu u)$
	$c^{(8)}_{Qu}$	$(\bar{Q}T^a\gamma_\mu Q)(\bar{u}T^a\gamma^\mu u)$
	$c^{(1)}_{Qd}$	$(\bar{Q}\gamma_\mu Q)(\bar{d}\gamma^\mu d)$
	$c^{(8)}_{Qd}$	$(\bar{Q}T^a\gamma_\mu Q)(\bar{d}T^a\gamma^\mu d)$

is used for the bbH and tHq production processes, and the 5FS for all other processes. All fermions are assumed to be massless when deriving the cross section scaling functions, except t and b in the 4FS, or t in the 5FS. For the decay parametrization, all fermions are taken to be massive. The light quark (u, d, s) and lepton masses are taken from [121]. Heavy quark masses (t, b, c) are aligned with the values in [120], where the masses of the c and b quarks are run up to the Higgs boson mass scale.

The SMEFT contributions do not only modify interaction vertices, but also affect the masses and decay widths of intermediate particles. These propagator corrections are included in all parametrizations derived with SMEFTSIM 3.0 [116], and are considered simultaneously with the vertex modifications. Special care is taken to ensure that double insertions, the inclusion of multiple new physics interactions in a single diagram, are avoided.

In the SMEFT framework, the matrix element for a particular Higgs boson production or decay process can be written as the sum of matrix elements originating from the SM and BSM Lagrangians,

$$\mathcal{M}_{\text{SMEFT}} = \mathcal{M}_{\text{SM}} + \mathcal{M}_{\text{BSM}}. \tag{25}$$

In this interpretation, the BSM contributions are restricted to diagrams with a single insertion of an EFT vertex such that,

$$\mathcal{M}_{\text{BSM}} = \sum_j \alpha_j c_j, \tag{26}$$

where the index j runs over all dimension-6 operators that contribute to the process, c_j are the corresponding WCs, and α_j are complex proportionality constants. Squaring the total matrix element gives,

$$\begin{aligned} |\mathcal{M}_{\text{SMEFT}}|^2 &= |\mathcal{M}_{\text{SM}}|^2 + (\mathcal{M}^*_{\text{SM}}\mathcal{M}_{\text{BSM}} + \mathcal{M}_{\text{SM}}\mathcal{M}^*_{\text{BSM}}) + |\mathcal{M}_{\text{BSM}}|^2 \\ &= |\mathcal{M}_{\text{SM}}|^2 + \sum_j \left(\mathcal{M}^*_{\text{SM}}\alpha_j + \mathcal{M}_{\text{SM}}\alpha^*_j\right) c_j + \sum_j |\alpha_j|^2 c_j^2 + \sum_{j\neq k} \left(\alpha^*_j\alpha_k + \alpha_j\alpha^*_k\right) c_j c_k, \end{aligned} \tag{27}$$

where the last sum is over all unique combinations of j and k with $j \neq k$. Dividing by $|\mathcal{M}_{\text{SM}}|^2$ yields a quadratic functional form

$$\mu = 1 + \sum_j A_j c_j + \sum_{jk} B_{jk} c_j c_k, \tag{28}$$

where A_j and B_{jk} are real constants that encode the impact of the WCs on the respective process. The A_j constants are referred to as the linear terms and account for the interference between the SM and BSM diagrams. The remaining purely BSM terms are split into two categories: B_{jj} are referred to as the quadratic terms, and B_{jk} for $j \neq k$ are the cross terms. The Higgs boson production cross sections and partial decay widths are proportional to $|\mathcal{M}_{\text{SMEFT}}|^2$ and therefore scale as

$$\begin{aligned} \mu^i_{\text{prod}}(\vec{c}) &= \frac{\sigma^i_{\text{SMEFT}}}{\sigma^i_{\text{SM}}} = 1 + \sum_j A^i_j c_j + \sum_{jk} B^i_{jk} c_j c_k, \\ \mu^f_{\text{partial}}(\vec{c}) &= \frac{\Gamma^f_{\text{SMEFT}}}{\Gamma^f_{\text{SM}}} = 1 + \sum_j A^f_j c_j + \sum_{jk} B^f_{jk} c_j c_k, \end{aligned} \tag{29}$$

where the index i refers to a particular STXS stage 1.2 bin and f refers to the Higgs boson decay channel. The decay rates also depend on the SMEFT corrections to the Higgs boson total decay width. This is calculated as the sum of the partial width scaling functions over all possible Higgs boson final states, weighted according to the SM predictions of the branching fractions at the highest available order, taken from [13],

$$\mu^{\text{H}}(\vec{c}) = \sum_f \mathcal{B}^f_{\text{SM,HO}}\, \mu^f_{\text{partial}} = 1 + \sum_j A^{\text{H}}_j c_j + \sum_{jk} B^{\text{H}}_{jk} c_j c_k. \tag{30}$$

The index f runs over all Higgs boson final states with up to four elementary particles produced, including those that are not explicitly targeted in the combination, e.g. $\text{H} \to \text{cc}$ and $\text{H} \to \text{gg}$. The linear terms A^{H}_j have been compared to the analytic solution of the Higgs boson total width provided in [122], and show excellent agreement.

As the Higgs boson is a scalar, narrow-width particle and only on-shell production is considered in this interpretation, the total scaling function for production bin i and decay channel f can be written as

$$\mu^{if}(\vec{c}) = \frac{\left[\sigma^i \mathcal{B}^f\right]_{\text{SMEFT}}}{\left[\sigma^i \mathcal{B}^f\right]_{\text{SM}}} = \mu^i_{\text{prod}}(\vec{c})\, \mu^f_{\text{partial}}(\vec{c})\, \mu^H(\vec{c})^{-1} = \frac{\left(1 + \sum_j A^i_j c_j + \sum_{jk} B^i_{jk} c_j c_k\right)}{\left(1 + \sum_j A^{\text{H}}_j c_j + \sum_{jk} B^{\text{H}}_{jk} c_j c_k\right)} \left(1 + \sum_j A^f_j c_j + \sum_{jk} B^f_{jk} c_j c_k\right). \tag{31}$$

The total scaling functions $\mu^{if}(\vec{c})$ act as multiplicative corrections to the SM predictions of the Higgs boson production cross section and branching fraction at the highest available order $\left[\sigma^i \mathcal{B}^f\right]_{\text{SM,HO}}$, as shown in equation (5). This procedure assumes that the SMEFT corrections factorize from higher order SM QCD and EW corrections.

The terms A_j and B_{jk} are derived numerically using the EFT2Obs tool [123] for all processes except the $\text{H} \to \gamma\gamma$ and $\text{H} \to \text{Z}\gamma$ partial decay widths, which are computed analytically [117, 118]. The EFT2Obs framework wraps and interfaces widely used packages to facilitate the process of deriving an EFT parametrization. It utilizes MadGraph5_aMC@NLO [66] (version 2.6.7) for event generation, and Pythia 8 [70] (version 8.306) for parton showering, hadronization, and modelling the underlying event, where the CP5 tune [71] is used. Higgs boson production events are further classified into the STXS stage 1.2 bins using the HiggsTemplateCrossSections Rivet routine [124]. A small modification is made to include events in which a Higgs boson and two leptons are produced at the same interaction vertex in the VH process. This guarantees the correct classification of events with leptons from off-shell vector boson decays, which can be enhanced significantly by SMEFT operators compared to the small SM contribution.

Events are generated under the SM hypothesis and then reweighted to different points in SMEFT parameter space using the SMEFTsim 3.0 and SMEFT@NLO UFO models. To derive the full quadratic dependence for N operators requires $2N + N(N-1)/2$ reweighting points in addition to the nominal SM hypothesis. The reweighting points are chosen to ensure that the overlap between the SM and BSM parameter spaces is sufficient to avoid large statistical uncertainties in the derived A_j and B_{jk} terms. For the four-fermion decay modes, corrections from the $\mathcal{O}_{\text{HW}}$, $\mathcal{O}_{\text{HB}}$, and $\mathcal{O}_{\text{HWB}}$ operators introduce photon-mediated diagrams that lead to a divergence at $m_{\text{ff}} \sim 0\,\text{GeV}$, where m_{ff} is the invariant mass of the subleading fermion pair. Dedicated simulated samples were generated for these operators to account for the significant difference between the SM and BSM phase spaces.

11.2. Impact of Higgs boson kinematic properties

Section 11.1 described how to parametrize SMEFT corrections to the Higgs boson production cross sections and branching fractions. However, the kinematic properties of Higgs boson events can be modified by introducing SMEFT operators, thus affecting other terms that enter the combined likelihood (equation (4)).

First, the efficiency factors $\epsilon^{if}_{r,\text{SM}}$ that encode the fraction of events from production region i and decay channel f that are selected in analysis region r, will become dependent on the SMEFT WCs, $\epsilon^{if}_r(\vec{c})$. The parametrization described in equation (31) becomes invalid when $\epsilon^{if}_r(\vec{c})$ differs significantly from the SM prediction. On the production side, the STXS stage 1.2 binning is sufficiently fine-grained to ensure that the efficiency factors are flat as functions of the WCs. In addition, the theoretical systematic uncertainties assigned to the SM efficiency factors $\vec{\theta}_{\text{th,acc}}$, are expected to cover the possible modifications induced by the SMEFT operators. As a result, this interpretation assumes that modifications to the kinematic properties of Higgs boson production do not affect the efficiency factors.

The same assumption cannot be made for the Higgs boson decay. Two-body Higgs boson decays are exempt as the kinematic properties of the final state are not affected by the SMEFT operators. However, four-body decays are particularly problematic as the STXS framework includes no fiducial phase space restriction on final-state particles. As discussed at the end of section 11.1, the effects of certain SMEFT operators are concentrated in regions of phase space (e.g. $m_{\text{ff}} \sim 0$) that are not SM-like, and lie outside of the experimental acceptance. For example, the $\text{H} \to \text{ZZ} \to 4\ell$ analysis [50] requires the invariant mass of the subleading lepton pair to be greater than 12 GeV to reduce large contributions from background processes. This selection criterion removes the parameter space that is significantly enhanced by the SMEFT operators, thus presenting a scenario where $\epsilon^{if}_r(\vec{c})$ differs significantly from the SM prediction.

A new procedure has been developed to fully account for these effects in the four-body decay channels, namely $\text{H} \to \text{ZZ} \to 4\ell$ and $\text{H} \to \text{WW} \to \ell\nu\ell\nu/\ell\nu\text{qq}'$. The EFT2Obs package [123] employs the

MadGraph5_aMC@NLO standalone reweighting feature [66] to output a reweighting module. This module can then be used to reweight existing MC samples without the computationally expensive task of generating new events. The new weights are calculated using the generator-level information, which is available in the samples, and can be used to study EFT effects after the event reconstruction and selection.

Signal events from the samples simulated under the SM hypothesis that enter the analysis regions for the $\mathrm{H} \to \mathrm{ZZ} \to 4\ell$ [50] and $\mathrm{H} \to \mathrm{WW} \to \ell\nu\ell\nu/\ell\nu\mathrm{qq}'$ [51] channels are identified. These events are subsequently reweighted to different points in SMEFT parameter space, and the corrected partial width scaling functions are derived. As these functions are derived using only events within experimental acceptance, the efficiency factor modifications are fully absorbed into the partial width scaling. The final parametrization uses the corrected partial width scaling functions for the $\mathrm{H} \to \mathrm{ZZ} \to 4\ell$ and $\mathrm{H} \to \mathrm{WW} \to \ell\nu\ell\nu/\ell\nu\mathrm{qq}'$ signal yield estimates. The total Higgs boson width is an inclusive quantity and therefore uses the partial width scaling functions without acceptance corrections in its derivation (equation (30)).

Kinematic modifications from SMEFT operators can also impact the shape of the signal distribution $\rho^{if}_{\mathrm{sig},r}$ as a function of the fitted observable. This may lead to a bias in the measured values of the WCs. For example, the $\mathrm{H} \to \mathrm{ZZ} \to 4\ell$ analysis [50] fits a discriminant $\mathcal{D}_{\mathrm{kin}}$, which depends on the kinematic properties of the four-body decay, to separate signal and background events. Using the standalone reweighting procedure, shape modifications were studied and found to be small compared with the statistical uncertainty in the observed signal yields. In other words, the potential bias induced by shape modifications is negligible compared with the range of WC values that the combination is sensitive to. As a result, all signal shapes $\rho^{if}_{\mathrm{sig},r}$ are assumed to follow the SM distributions, and to be independent of the WC values.

11.3. SMEFT parametrization summary

The parametrizations of the SMEFT corrections are computed up to quadratic order for each production cross section and partial width. As the total scaling $\mu^{if}(\vec{c})$ is the product of multiple quadratic functions, a Taylor expansion is performed and the series is truncated to remove higher-order contributions. The WC constraints are extracted for two different series truncations: the linear parametrization includes terms up to $\mathcal{O}(c/\Lambda^2)$,

$$\mu^{if}(\vec{c}) = 1 + \sum_j \left(A^i_j + A^f_j - A^{\mathrm{H}}_j\right) c_j, \tag{32}$$

and the linear-plus-quadratic parametrization includes terms up to $\mathcal{O}(c^2/\Lambda^4)$,

$$\begin{aligned}\mu^{if}(\vec{c}) = 1 &+ \sum_j \left(A^i_j + A^f_j - A^{\mathrm{tot}}_j\right) c_j + \sum_{jk} \left(B^i_{jk} + B^f_{jk} - B^{\mathrm{H}}_{jk}\right) c_j c_k \\ &+ \left(\sum_j A^i_j c_j\right)\left(\sum_j A^f_j c_j\right) - \left(\sum_j A^i_j c_j\right)\left(\sum_j A^{\mathrm{H}}_j c_j\right) \\ &- \left(\sum_j A^f_j c_j\right)\left(\sum_j A^{\mathrm{H}}_j c_j\right) + \left(\sum_j A^{\mathrm{H}}_j c_j\right)^2 .\end{aligned} \tag{33}$$

It is important to note that the $\mathcal{O}(c^2/\Lambda^4)$ terms in the linear-plus-quadratic parametrization enter at the same order in $1/\Lambda$ as the linear contributions from dimension-8 operators. This introduces an inconsistency in the linear-plus-quadratic parametrization, as only some contributions at $\mathcal{O}(\Lambda^{-4})$ are considered. As a result, the linear parametrization is considered for the main results of this interpretation. The linear-plus-quadratic parametrization is used as a comparison to indicate how much the inclusion of $\mathcal{O}(\Lambda^{-4})$ terms can impact the sensitivity of the results.

Only the A_j and B_{jk} constants with an absolute value greater than 0.01 are included in the parametrization. This threshold corresponds to a 1% deviation for WC values of 1. Any effects that fall below this threshold are deemed to be negligible, and the corresponding A_j and B_{jk} terms are dropped from the parametrization.

The relative impacts of the SMEFT operators on the Higgs boson production cross sections and branching fractions are shown in figures 21–23. The quantities show the change in cross section or branching fraction relative to the SM expectation, for the WC values indicated in the legend. These WC values represent the expected symmetrized 95% CL intervals in the linear-plus-quadratic parametrization, when considering effects in one SMEFT operator at a time. This choice therefore illustrates the size

of cross section and branching fraction modifications to which the combination is currently sensitive. The impact from the linear and linear-plus-quadratic parametrizations are shown by the filled and open histograms, respectively. The figures are split by operator groups.

The SMEFT parametrizations are provided in the POPXF format [125], which are included in the HEPData record for this paper [41].

11.4. Constraints on individual Wilson coefficients

Maximum likelihood fits are performed to extract constraints on the WCs by introducing $\mu^{if}(\vec{c})$ into the signal yield parametrization of equation (5). Individual constraints are obtained by scanning over one WC, while fixing all others to the SM expectation ($c_j = 0$). The expected and observed constraints are summarized in figure 24 and table 17. The best fit values, and 68% and 95% CL intervals, are shown for both the linear and linear-plus-quadratic parametrizations. The results are also translated into 95% CL lower limits on the energy scale of BSM physics Λ_j. This quantity is evaluated by converting the symmetrized 95% CL constraints on c_j/Λ^2 into an energy scale limit, assuming $c_j = 1$. The WCs are categorized into operators of the same form in the SMEFT Lagrangian, and then listed in order of excluded energy scale (sensitivity) in the linear-plus-quadratic parametrization.

The tightest constraints are obtained for the c_{HG} and c_{HB} WCs, which are mostly driven by the measurements of ggH production and of the $H \to \gamma\gamma$ decay channel, respectively. The 95% CL lower limit on the new physics energy scale for these operators is around 15 TeV.

Most WCs show good agreement with the SM. However, there are a number of WCs that present a sizeable deviation from zero. For instance $c_{Hq}^{(3)}$ has a p-value with respect to the SM hypothesis of $p_{\text{SM}} = 0.01$. Figure 22 shows that this operator impacts the WH and ZH leptonic cross sections, with larger SMEFT corrections for the higher p_{T}^{V} bins. Comparing with the measurements in figure 10, it is clear how the observed excesses in the WH and ZH leptonic $p_{\text{T}}^{\text{V}} > 250\,\text{GeV}$ cross sections pull $c_{Hq}^{(3)}$ away from zero. The p_{SM} values are provided for each WC in table 17.

Figure 24 also provides a comparison of the constraints obtained using the linear and linear-plus-quadratic parametrizations. The terms $\mathcal{O}(c^2/\Lambda^4)$ are shown to have a larger effect for the more weakly constrained WCs. This follows from the fact that the relative quadratic-to-linear contribution,

$$\frac{B_{jj}c_j^2}{A_j c_j} = \frac{B_{jj}c_j}{A_j}, \tag{34}$$

grows as a function of c_j. In other words, the importance of the $\mathcal{O}(c^2/\Lambda^4)$ terms increases as one moves further from the SM hypothesis. Conversely, the sensitivities for the more tightly constrained operators are dominated by the $\mathcal{O}(c/\Lambda^2)$ terms. Several WCs are only constrained when quadratic terms are considered. These include four-fermion operators that impact ttH production and are represented by the hatched lines in figure 24.

11.5. Constraints on linear combinations of Wilson coefficients

It is expected that BSM physics would induce modifications in multiple operators simultaneously. The data used in the combination do not contain sufficient information to constrain all WCs at the same time, as some of the WCs have (nearly) degenerate effects on the signal processes. This leads to flat directions in the likelihood surface, meaning that many degrees of freedom are left unconstrained.

Linear combinations of the SMEFT WCs that can be constrained simultaneously are used to define a new basis. It is possible to construct this basis purely from theoretical considerations. For example, the Higgs basis [13] provides a natural choice for the production cross section and branching fraction observables measured in the combination. However, a principal component analysis (PCA) technique is instead used to identify the sensitive linear combinations directly from data.

The Hessian matrix $\mathcal{H}_{\text{STXS}}$ from the 97 POI fit (result shown in figure 12) is computed numerically for an Asimov data set under the SM hypothesis. This is the matrix of second derivatives of the log-likelihood with respect to all fit parameters, including both POIs and NPs. Under the Gaussian approximation, the covariance matrix $\mathcal{C}_{\text{STXS}}$ is equal to the inverse of the Hessian matrix, $\mathcal{H}_{\text{STXS}}^{-1}$. A submatrix of the covariance matrix, $\widetilde{\mathcal{C}}_{\text{STXS}}$, is obtained by removing all rows and columns that do not correspond to the POIs. This reduced Hessian matrix is then approximated using the inverse of the submatrix, $\widetilde{\mathcal{H}}_{\text{STXS}} = \widetilde{\mathcal{C}}_{\text{STXS}}^{-1}$.

The $\widetilde{\mathcal{H}}_{\text{STXS}}$ matrix is translated to the SMEFT WC basis using the linear parametrization shown in equation (32). A translation matrix $\mathcal{T}$ is constructed with terms,

$$\mathcal{T}_j^{\,if} = A_j^i + A_j^f - A_j^{\text{H}}, \tag{35}$$

Table 17. Expected and observed 95% CL intervals on the WCs, when considering modifications to one WC at a time. The results for both the linear and linear-plus-quadratic parametrizations are provided. For non-contiguous intervals, only the range containing the best fit point is quoted; the full intervals are available in the HEPData record for this paper [41]. The best fit values and p_{SM} values are also given in the table.

		Linear				Linear + quadratic			
Operator group	WCs	Expected 95% CL	Observed 95% CL	Best fit	p_{SM}	Expected 95% CL	Observed 95% CL	Best fit	p_{SM}
X^3	c_G	$[-1.6, 2.3]$	$[-1.5, 2.4]$	-0.028	0.97	$[-0.33, 0.29]$	$[-0.29, 0.26]$	-0.001	1.00
	c_W	$[-0.17, 0.15]$	$[-0.32, 0.03]$	-0.14	0.11	$[-0.16, 0.16]$	$[-0.3, 0.032]$	-0.13	0.11
H^4D^2	$c_{H\Box}$	$[-0.79, 0.86]$	$[-0.57, 1.1]$	0.25	0.55	$[-0.79, 0.86]$	$[-0.57, 1.1]$	0.25	0.56
	c_{HD}	$[-4.1, 3.9]$	$[-2.2, 5.9]$	1.9	0.35	$[-4.1, 3.9]$	$[-2.3, 5.8]$	1.9	0.36
X^2H^2	c_{HG}	$[-0.0038, 0.0043]$	$[-0.0039, 0.0042]$	$-7e-05$	0.98	$[-0.0039, 0.0043]$	$[-0.0039, 0.0042]$	$-7e-05$	0.98
	c_{HW}	$[-0.012, 0.011]$	$[-0.021, 0.0043]$	-0.0075	0.22	$[-0.012, 0.012]$	$[-0.02, 0.0044]$	-0.0074	0.22
	c_{HB}	$[-0.004, 0.0037]$	$[-0.0075, 0.00086]$	-0.0033	0.13	$[-0.0039, 0.0038]$	$[-0.007, 0.00081]$	-0.0029	0.12
	c_{HWB}	$[-0.0066, 0.0072]$	$[-0.002, 0.013]$	0.0049	0.17	$[-0.0068, 0.007]$	$[-0.0022, 0.012]$	0.0048	0.19
ψ^2H^3	$\mathrm{Re}(c_{eH})$	$[-2, 1.9]$	$[-2.1, 1.8]$	-0.13	0.89	$[-2, 1.9]$	$[-2.1, 1.8]$	-0.075	0.95
	$\mathrm{Re}(c_{bH})$	$[-0.028, 0.03]$	$[-0.017, 0.043]$	0.012	0.44	$[-0.03, 0.028]$	$[-0.018, 0.039]$	0.011	0.43
	$\mathrm{Re}(c_{tH})$	$[-1.3, 1.1]$	$[-1, 1.3]$	0.22	0.71	$[-1.3, 1.1]$	$[-1, 1.3]$	0.21	0.72
ψ^2XH	$\mathrm{Re}(c_{tG})$	$[-0.15, 0.14]$	$[-0.16, 0.13]$	-0.0092	0.90	$[-0.14, 0.14]$	$[-0.15, 0.13]$	-0.0028	0.97
	$\mathrm{Re}(c_{tW})$	$[-0.14, 0.13]$	$[-0.26, 0.023]$	-0.12	0.11	$[-0.13, 0.13]$	$[-0.25, 0.025]$	-0.11	0.11
	$\mathrm{Re}(c_{tb})$	$[-0.075, 0.068]$	$[-0.14, 0.018]$	-0.056	0.14	$[-0.073, 0.071]$	$[-0.13, 0.018]$	-0.054	0.14
	$\mathrm{Re}(c_{bG})$	—				$[-0.3, 0.3]$	$[-0.39, 0.35]$	-0.21	0.32
	$\mathrm{Re}(c_{bW})$	—				$[-0.42, 0.42]$	$[-0.67, -0.23]$	-0.48	0.01
ψ^2H^2D	$c_{Hl}^{(1)}$	$[-1.3, 1.6]$	$[-1.3, 1.7]$	0.13	0.81	$[-1.6, 1.4]$	$[-1.5, 1.5]$	0.14	0.88
	$c_{Hl}^{(3)}$	$[-0.39, 0.36]$	$[-0.62, 0.17]$	-0.21	0.29	$[-0.37, 0.37]$	$[-0.59, 0.17]$	-0.21	0.28
	$c_{Hq}^{(1)}$	$[-0.32, 0.25]$	$[-0.63, 0.047]$	-0.25	0.10	$[-0.075, 0.098]$	$[-0.11, 0.13]$	-0.065	0.13
	$c_{Hq}^{(3)}$	$[-0.035, 0.039]$	$[0.012, 0.097]$	0.055	0.01	$[-0.058, 0.032]$	$[0.0099, 0.064]$	0.039	0.01
	$c_{HQ}^{(1)}$	$[-6.5, 7.5]$	$[-2.5, 12]$	4.1	0.23	$[-2.1, 1.8]$	$[-2.8, 2.4]$	1.5	0.14
	$c_{HQ}^{(3)}$	$[-10, 10]$	$[-7.2, 13]$	3.6	0.49	$[-2, 1.9]$	$[-2.9, 2.6]$	-2	0.06
	c_{He}	$[-2.2, 1.9]$	$[-2.8, 1.5]$	-0.47	0.67	$[-2.2, 1.9]$	$[-2.8, 1.5]$	-0.48	0.66
	c_{Hu}	$[-0.21, 0.25]$	$[-0.018, 0.5]$	0.22	0.07	$[-0.17, 0.1]$	$[-0.21, 0.16]$	0.091	0.08
	c_{Hd}	$[-0.52, 0.48]$	$[-1.2, -0.075]$	-0.58	0.02	$[-0.18, 0.21]$	$[-0.26, -0.0021]$	-0.17	0.05
	$\mathrm{Re}(c_{Htb})$	—				$[-0.77, 0.77]$	$[0.43, 1.2]$	0.86	0.01
	c_{Ht}	$[-29, 19]$	$[-37, 14]$	-8.5	0.43	$[-19, 30]$	$[13, 56]$	30	0.03
	c_{Hb}	—				$[-42, 41]$	$[-45, 46]$	19	0.64

(Continued.)

Table 17. (Continued.)

$(\bar{L}L)(\bar{L}L)$	$c_{ll}^{(1)}$	$[-0.65, 0.7]$	$[-0.3, 1.1]$	0.39	0.27	$[-0.65, 0.7]$	$[-0.3, 1.1]$	0.38	0.27
	$c_{Qq}^{(11)}$	—				$[-0.96, 0.94]$	$[-0.81, 0.78]$	−0.018	0.97
	$c_{Qq}^{(18)}$	$[-2.2, 3.1]$	$[-2.5, 2.5]$	−0.43	0.74	$[-2.6, 1.5]$	$[-2.2, 1.2]$	−0.45	0.69
	$c_{Qq}^{(31)}$	$[-5.2, 6.8]$	$[-2.3, 12]$	4.4	0.23	$[-0.64, 0.61]$	$[-0.92, 0.84]$	−0.6	0.06
	$c_{Qq}^{(38)}$	$[-9.8, 14]$	$[-12, 8]$	−3.8	0.47	$[-1.4, 1.3]$	$[-1.9, -0.13]$	−1.3	0.05
$(\bar{R}R)(\bar{R}R)$	$c_{tu}^{(1)}$	—				$[-1.2, 1.3]$	$[-1.1, 1.1]$	−0.018	0.99
	$c_{tu}^{(8)}$	$[-3.4, 5]$	$[-3.5, 4.3]$	−0.33	0.88	$[-3.2, 2.1]$	$[-2.8, 1.8]$	−0.15	0.87
	$c_{td}^{(1)}$	—				$[-1.8, 1.8]$	$[-1.5, 1.5]$	0.015	0.99
	$c_{td}^{(8)}$	$[-5.9, 8.4]$	$[-6.6, 7]$	−0.93	0.72	$[-4.6, 3.1]$	$[-3.9, 2.6]$	−0.45	0.85
$(\bar{L}L)(\bar{R}R)$	$c_{qt}^{(1)}$	—				$[-1.1, 1.2]$	$[-0.86, 0.84]$	−0.006	1.00
	$c_{qt}^{(8)}$	$[-2.2, 3.2]$	$[-2.4, 2.6]$	−0.37	0.77	$[-3.3, 1.7]$	$[-2.2, 1.3]$	−0.17	0.85
	$c_{Qu}^{(1)}$	—				$[-1.4, 1.3]$	$[-1.2, 1.2]$	−0.018	0.99
	$c_{Qu}^{(8)}$	$[-3.5, 5.1]$	$[-3.7, 4.4]$	−0.42	0.82	$[-3.7, 2.3]$	$[-3.3, 2]$	−0.33	0.87
	$c_{Qd}^{(1)}$	—				$[-1.7, 1.6]$	$[-1.3, 1.3]$	0.009	0.99
	$c_{Qd}^{(8)}$	$[-5.7, 8.1]$	$[-6.3, 6.5]$	−1.1	0.73	$[-4.1, 2.9]$	$[-3.2, 2.3]$	−0.3	0.84



Figure 21. Impact of the SMEFT operators on the Higgs boson cross sections and branching fractions. Units of GeV are assumed for all numerical values related to the p_T^H, m_{jj}, p_T^{Hjj}, and p_T^V variables. The impacts are shown for operators from the following groups: X^3, H^4D^2, X^2H^2, ψ^2H^3, and ψ^2XH. The WCs are set to the expected symmetrized 95% CL interval value in the linear-plus-quadratic parametrization, assuming all other WCs are set to zero (SM). The impacts are shown relative to the SM predictions for the linear parametrization in the filled histograms, and the linear-plus-quadratic parametrization in the open histograms.

where the indices (i,f) run over the 97 μ^{if} POIs, and A_j are the corresponding linear terms for WC c_j. If multiple STXS stage 1.2 bins are merged in the POI definition, the linear term for production is computed as the mean of the individual STXS stage 1.2 bin A_j^i constants, weighted according to their SM



Figure 22. Impact of the SMEFT operators on the Higgs boson production cross sections and branching fractions. Units of GeV are assumed for all numerical values related to the $p_{\mathrm{T}}^{\mathrm{H}}$, m_{jj}, $p_{\mathrm{T}}^{\mathrm{Hjj}}$, and $p_{\mathrm{T}}^{\mathrm{V}}$ variables. The impacts are shown for operators from the $\psi^2 H^2 D$ group. The WCs are set to the expected symmetrized 95% CL interval value in the linear-plus-quadratic parametrization, assuming all other WCs are set to zero (SM). The impacts are shown relative to the SM predictions for the linear parametrization in the filled histograms, and the linear-plus-quadratic parametrization in the open histograms.

cross section predictions. The reduced Hessian matrix in terms of the SMEFT WCs is then obtained by performing the translation,

$$\widetilde{\mathcal{H}}_{\mathrm{SMEFT}} = \mathcal{T}^{T} \widetilde{\mathcal{H}}_{\mathrm{STXS}} \mathcal{T}, \tag{36}$$



Figure 23. Impact of the SMEFT operators on the Higgs boson production cross sections and branching fractions. Units of GeV are assumed for all numerical values related to the p_T^H, m_{jj}, p_T^{Hjj}, and p_T^V variables. The impacts are shown for operators from the four-fermion groups. The WCs are set to the expected symmetrized 95% CL interval value in the linear-plus-quadratic parametrization, assuming all other WCs are set to zero (SM). The impacts are shown relative to the SM predictions for the linear parametrization in the filled histograms, and the linear-plus-quadratic parametrization in the open histograms.

where $\mathcal{T}^T$ is the transpose of matrix $\mathcal{T}$. An eigendecomposition of this matrix $\widetilde{\mathcal{H}}_{\text{SMEFT}} = \mathcal{R}^T D \mathcal{R}$, is performed. The rotation matrix $\mathcal{R}$ provides linear combinations of the SMEFT WCs (eigenvectors, $\vec{\text{EV}}$), and the diagonal matrix D of eigenvalues λ indicating their constraining power. The scaling functions



Figure 24. Individual constraints on the SMEFT WCs. The left panel shows the best fit values, and the 68% and 95% CL intervals, when considering modifications in one SMEFT operator at a time. The results from the linear and linear-plus-quadratic parametrizations are shown in blue and orange, respectively. Arrows are used to indicate where the 95% CL interval extends beyond the plotted range. In the right panel, the results are translated into a 95% lower limit on the BSM physics energy scale. The WCs are categorized into operators from the same group, and then listed in order of the probed energy scale using the linear-plus-quadratic parametrization. The hatched lines represent WCs that cannot be constrained in the linear-only parametrization.

$\mu^{if}(\vec{\mathrm{EV}})$ that enter the combined likelihood are now parametrized as functions of the linear combinations by rotating the nominal SMEFT WCs, $\vec{c} = \mathcal{R}^T\vec{\mathrm{EV}}$.

This procedure provides as many eigenvectors as there are WCs (43). The constraining power for eigenvector $\mathrm{EV}_j \in \vec{\mathrm{EV}}$ is provided by the quantity $1/\sqrt{\lambda_j}$, which is an estimate of the (symmetrized) expected 68% CL interval. Eigenvectors with $1/\sqrt{\lambda_j} > 10$ are fixed to the SM (i.e. 0) in the fit, as the data are deemed not to be sensitive to these directions. This choice of threshold is arbitrary, but it ensures that the fit converges as the curvature of the likelihood is sufficiently high. In addition, this

Table 18. Expected and observed 95% CL intervals on the linear combinations of SMEFT WCs. The results are provided for the linear parametrization. The best fit values are also given.

Eigenvector	Expected 95% CL	Observed 95% CL	Best fit
EV_0	$[-0.0034, 0.0038]$	$[-0.00097, 0.007]$	0.0028
EV_1	$[-0.0054, 0.0051]$	$[-0.0053, 0.0056]$	0.00 033
EV_2	$[-0.053, 0.055]$	$[-0.024, 0.1]$	0.036
EV_3	$[-0.06, 0.062]$	$[0.035, 0.17]$	0.1
EV_4	$[-0.061, 0.065]$	$[-0.031, 0.11]$	0.036
EV_5	$[-0.48, 0.47]$	$[-0.71, 0.32]$	-0.18
EV_6	$[-0.53, 0.56]$	$[-0.38, 0.84]$	0.21
EV_7	$[-0.65, 0.68]$	$[-1.2, 0.2]$	-0.52
EV_8	$[-0.94, 0.95]$	$[-1.6, 0.35]$	-0.64
EV_9	$[-1.4, 1.4]$	$[-0.55, 2.3]$	0.85
EV_{10}	$[-2.5, 2.5]$	$[-2.4, 2.8]$	0.16
EV_{11}	$[-2.7, 2.3]$	$[-1.3, 3.6]$	1.2
EV_{12}	$[-2.6, 2.6]$	$[-2, 4.4]$	1.2
EV_{13}	$[-8.3, 7.4]$	$[-17, 0.82]$	-7.4
EV_{14}	$[-8.2, 11]$	$[-5, 16]$	4.5
EV_{15}	$[-11, 9.8]$	$[-20, 2.5]$	-8.1
EV_{16}	$[-14, 12]$	$[-19, 16]$	-2.6

threshold avoids probing regions of parameter space where the SMEFT perturbative expansion breaks down. In total, 17 eigenvectors are included in the fit, shown by the truncated $\mathcal{R}$ matrix in figure 25.

The linear effects of the eigenvectors on the $\mu^{if} = [\sigma^i \mathcal{B}^f]_{\text{obs}}/[\sigma^i \mathcal{B}^f]_{\text{SM,HO}}$ POIs are summarized in figures 26 and 27. The figures show the change in $\sigma^i \mathcal{B}^f$ relative to the SM prediction, for EV_j set to the boundary of the expected (symmetrized) 95% CL interval on each parameter. The eigenvectors are ordered in terms of sensitivity, with the most sensitive (EV_0) at the top and least sensitive (EV_{16}) at the bottom. The most sensitive direction is driven by the measurement of ggH in the $H \to \gamma\gamma$ decay channel.

A maximum likelihood fit is performed to extract constraints on the eigenvectors by using $\mu^{if}(\vec{EV})$ to parametrize the signal yields. In this fit, all 17 eigenvectors are varied simultaneously. These constraints are provided for a linear parametrization with terms up to $\mathcal{O}(EV/\Lambda^2)$. Introducing quadratic terms means that the eigenvectors obtained via the PCA procedure are no longer orthogonal. The reason for this is that the matrix $\mathcal{T}$, used to map the 97 POIs to the SMEFT WCs, is defined using only the linear terms. In addition, the introduction of quadratic terms can lead to the likelihood fit converging to one of many local minima which arise when allowing multiple directions in SMEFT parameter space to vary simultaneously. As a result, no linear-plus-quadratic results are provided for the rotated basis fits.

The expected and observed constraints are summarized in figure 28 and table 18. The best fit values, as well as the 68% and 95% CL intervals, are shown. Again, the results are translated into 95% CL lower limits on the energy scale of BSM physics Λ_j, assuming the corresponding $EV_j = 1$. The eigenvectors are ordered according to their respective excluded energy scale. The 95% CL intervals are in the range of approximately ± 20 for the least constrained eigenvector, to ± 0.005 for the most constrained direction (assuming $\Lambda = 1\,\text{TeV}$).

Overall the results show good agreement with the SM, with $p_{\text{SM}} = 0.11$. The largest deviation from the SM is observed in EV_3, where the best fit value and 68% CL interval are $EV_3 = 0.101^{+0.036}_{-0.034}$. As can be seen in figure 25, this eigenvector is dominated by the contribution from $c_{Hq}^{(3)}$, which matches the tension observed in the individual constraint for this WC.

The correlation matrix for the fitted eigenvectors is shown in figure 29. Overall, an adequate level of decorrelation is achieved, which helps the fit to converge. However, some correlations are present (up to 83%). The primary source of these correlations is the different splitting of signal contributions in the PCA procedure and in the full combined likelihood. The 97 μ^{if} POIs used in the PCA are defined with channel-dependent merging of STXS bins, whereas the full combined likelihood splits signal contributions according to the full STXS stage 1.2 binning scheme. Since the SMEFT parametrization is defined at the STXS stage 1.2 granularity, bins that are merged in the 97 POI fit can scale differently with the eigenvectors, thereby inducing correlations. Additional correlations arise because the eigenvectors are derived using an Asimov data set under the SM hypothesis, and are therefore not guaranteed to be orthogonal at the observed best fit point. Finally, correlations can also be induced in the maximum likelihood fit due to the correlated effects of NPs.



Figure 25. Truncated rotation matrix $\mathcal{R}$, derived using the PCA procedure. Only the 17 eigenvectors that are included in the fit are shown. Each row represents a different eigenvector EV_j ordered by constraining power. The left panel shows $1/\sqrt{\lambda_j}$ for each eigenvector, where λ_j is the corresponding eigenvalue. This quantity provides an estimate of the expected 68% CL intervals. The vertical red line identifies the threshold beyond which the eigenvectors are not considered in the fit. The right panel shows the rotation matrix elements, $\mathcal{R}_{jk}$, for each Wilson coefficient c_k, such that $\mathrm{EV}_j = \sum_k \mathcal{R}_{jk} c_k$. The size of each element is represented by a colour scale with red meaning large positive values and blue meaning large negative values. The value of each element is also shown in the plot.



Figure 26. Impact of the eigenvectors EV_j on the μ^{if} parameters from the 97 POI fit (figure 12), for the $H \to \gamma\gamma$, $H \to ZZ \to 4\ell$, $H \to \mu\mu$, and $H \to Z\gamma \to \ell\ell\gamma$ decay channels. Units of GeV are assumed for all numerical values related to the p_T^H, m_{jj}, p_T^{Hjj}, and p_T^V variables. The EV_j are individually set to their expected symmetrized 95% CL interval value in the fully profiled fit. When varying one eigenvector, all other eigenvectors are set to zero (SM). The impacts are shown relative to the SM predictions for the linear parametrization. The eigenvectors are ordered from most sensitive (upper) to least sensitive (lower). All eigenvectors included in the fit are shown.



Figure 27. Impact of the eigenvectors EV_j on the μ^{if} parameters from the 97 POI fit (figure 12), for the H → WW, H → ττ, and H → bb decay channels. Units of GeV are assumed for all numerical values related to the p_T^H, m_{jj}, p_T^{Hjj}, and p_T^V variables. The EV_j are individually set to their expected symmetrized 95% CL interval value in the fully profiled fit. When varying one eigenvector, all other eigenvectors are set to zero (SM). The impacts are shown relative to the SM predictions for the linear parametrization. The eigenvectors are ordered from most sensitive (upper) to least sensitive (lower). All eigenvectors included in the fit are shown.

12. Summary

The most comprehensive combined measurements of Higgs boson (H) production and decay rates performed by the CMS Collaboration to date are reported. The combination uses proton–proton collision



Figure 28. Constraints on the linear combinations of SMEFT WCs extracted with the PCA procedure. The left panel shows the observed best fit values, and 68% and 95% CL intervals, as well as the expected 95% CL intervals, for a linear parametrization with terms up to $\mathcal{O}(\mathrm{EV}/\Lambda^2)$. An arrow is used to indicate where the 95% CL interval for EV_2 extends beyond the plotted range. In the right panel, the results are translated into a 95% lower limit on the BSM physics energy scale, assuming $\mathrm{EV}_j = 1$. The eigenvectors are listed in order of the expected excluded energy scale.

data recorded by the CMS experiment at $\sqrt{s} = 13\,\mathrm{TeV}$ between 2016 and 2018, corresponding to an integrated luminosity of $138\,\mathrm{fb}^{-1}$. The decay channels included in the combined measurements are $\mathrm{H} \to \gamma\gamma$, $\mathrm{H} \to \mathrm{ZZ}$, $\mathrm{H} \to \mathrm{WW}$, $\mathrm{H} \to \tau\tau$, $\mathrm{H} \to \mathrm{bb}$, $\mathrm{H} \to \mu\mu$, and $\mathrm{H} \to \mathrm{Z}\gamma \to \ell\ell\gamma$ ($\ell = \mathrm{e}$ or μ). Information in the events from each decay channel is used to target several Higgs boson production processes, namely gluon fusion (ggH), vector boson fusion, W (WH) and Z (ZH) boson-associated production, Higgs boson production in association with a top quark pair, and Higgs boson production in association with a top quark (tH). Results are provided for various assumptions on the scaling behaviour of Higgs boson production and decay. This includes measurements of signal strengths, production cross sections, branching fractions, and, for the first time, a combined measurement by the CMS Collaboration of kinematic regions defined by the simplified template cross section (STXS) framework.

The inclusive signal strength is measured to be $1.014^{+0.055}_{-0.053}$ times the standard model (SM) expectation, for a Higgs boson mass of 125.38 GeV, where the largest contribution to the total uncertainty originates from the theoretical systematic component. The per production process signal strengths have a p-value of $p_{\mathrm{SM}} = 0.02$. The leading contribution to this low p-value is the observed excess of 2.2 standard deviations in the tH production process. The corresponding 68% confidence level (CL) intervals range from 7.5% for ggH production to 39% for tH production, relative to the best fit value. The per decay channel signal strengths exhibit a good level of compatibility with the SM, with a p-value of $p_{\mathrm{SM}} = 0.33$. The 68% CL intervals range from 8% for $\mathrm{H} \to \gamma\gamma$ to 39% for $\mathrm{H} \to \mathrm{Z}\gamma$.

Production cross section measurements are performed at two levels of granularity. The STXS stage 0 measurements correspond to the different Higgs boson production processes, while the STXS stage



Figure 29. Correlation matrix for the linear combinations of SMEFT WCs extracted with the PCA procedure.

1.2 measurements further partition the phase space into nonoverlapping kinematic regions. In total, 32 kinematic regions are measured simultaneously, of which 13 are associated with ggH production, five with vector boson fusion production, four with each of WH and ZH production, where the vector boson decays leptonically, five with Higgs boson production in association with a top quark pair, and one with tH production. The production cross sections are extracted as products with the $H \to ZZ$ branching fraction, and the ratios of branching fractions are included as additional parameters in the fit to account for modifications to the SM Higgs boson decay rates. Overall, the STXS stage 1.2 measurements show reasonable agreement with the SM predictions ($p_{SM} = 0.06$), with some tensions observed in the high transverse momentum (p_T^V) regions of the WH and ZH leptonic production processes, as well as for tH production. The 68% CL intervals relative to the SM predicted values range from ∼14% for the ggH STXS bin with no additional jets and Higgs boson transverse momentum between 10 and 200 GeV to ∼260% for the ggH STXS bin with Higgs boson transverse momentum greater than 650 GeV. A measurement considering a separate parameter for each STXS bin in each decay channel is also performed. In total 97 parameters are fitted, which constitutes the most granular measurement of the Higgs boson ever performed by the CMS Collaboration. The compatibility of this measurement with the SM is $p_{SM} = 0.006$.

Several interpretations of the measurements are also provided. Higgs boson coupling modifiers are probed in the κ-framework for both the resolved and effective configurations. In the resolved coupling modifier configuration, the measurements exhibit a good level of compatibility with the SM ($p_{SM} = 0.12$). The vector boson and third generation fermion coupling modifiers are measured with 68% CL intervals ranging from 6% for κ_W to 12% for κ_b. The Higgs boson coupling modifier to muons is measured with a 68% CL interval of 20%. Models with different assumptions on the total Higgs boson decay width are provided for the effective coupling modifier configuration, including models with beyond-the-SM (BSM) decays of the Higgs boson. In order to constrain the branching fraction to invisible final states that are allowed in BSM models, analyses targeting the $H \to$ inv decay are included. One model

also includes off-shell analysis regions in the combination to provide a constraint on the total Higgs boson decay width directly from data. In this fit, the invisible (undetected) branching fraction is constrained to be less than 13% (25%) at the 95% CL. Ratios of coupling modifiers are also extracted, including models that test the symmetry of the couplings of the Higgs boson to fermions. These models are further used to extract constraints on specific extensions of the SM that contain a second Higgs doublet. No significant deviations from the SM are observed and the allowed parameter spaces are significantly reduced with respect to previous indirect constraints from CMS. In addition, constraints on the trilinear Higgs boson self-coupling are provided. These constraints arise from next-to-leading order electroweak corrections to the Higgs boson production and decay rates. Assuming other Higgs boson couplings are as predicted in the SM, the trilinear self-coupling is measured to be $2.14^{+3.95}_{-3.16}$, relative to the SM prediction.

An interpretation in the SM effective field theory (SMEFT) framework is also performed. This framework provides a model-agnostic tool to indirectly probe BSM physics via measurements of Higgs boson production and decay, assuming that the new particles of the BSM theory exist at an energy scale larger than the electroweak scale. A SMEFT parametrization of STXS measurements is derived using simulation. Individual constraints are extracted on the 43 SMEFT Wilson coefficients to which the combination is sensitive. To assess the compatibility of data with the SM, a p-value is computed for each of the Wilson coefficients. These p-values generally indicate good agreement with the SM predictions, with the majority being above 0.1. The largest discrepancy from the SM is observed in the $c_{Hq}^{(3)}$ parameter ($p_{\mathrm{SM}} = 0.01$), which is driven by the observed excesses in the high-$p_{\mathrm{T}}^{\mathrm{V}}$ WH and ZH leptonic STXS measurements. In addition, simultaneous constraints are provided on linear combinations of SMEFT operators, where the constrained directions in parameter space are extracted using a principal component analysis procedure. A total of 17 independent directions in SMEFT parameter space are constrained, and the measurements show reasonable compatibility with the SM hypothesis ($p_{\mathrm{SM}} = 0.11$).

In conclusion, the results show good compatibility with the SM predictions for the majority of the measured parameters.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG and PSG, TARISTU24-TK10 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMFTR, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIT and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES, NSC, and NAWA (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI (Thailand); TUBITAK and TENMAK (Türkiye); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, Contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, 101001205, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, Project No. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA) and Fonds voor Wetenschappelijk Onderzoek Contract No. 1228724 N (Belgium); the Beijing Municipal Science & Technology Commission, No. Z191100007219010, the Fundamental Research Funds for the Central Universities, the Ministry of Science and Technology of China under Grant No. 2023YFA1605804, the Natural Science Foundation of China under Grant No. 12535004, and USTC Research Funds of the Double First-Class Initiative No. YD2030002017 (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli

National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy—EXC 2121 'Quantum Universe' – 390833306, and under Project No. 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project No. 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2025-1.1.5-NEMZ_KI-2025-00004 (Hungary); the Council of Science and Industrial Research, India; ICSC—National Research Center for High Performance Computing, Big Data and Quantum Computing, FAIR—Future Artificial Intelligence Research, and CUP I53D23001070006 (Mission 4 Component 1), funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, Project No. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369, 2021/43/B/ST2/01552, 2023/49/B/ST2/03273, and the NAWA Contract BPN/PPO/2021/1/00011 (Poland); the Fundação para a Ciência e a Tecnologia (Portugal); the National Priorities Research Program by Qatar National Research Fund; MICIU/AEI/10.13039/501100011033, ERDF/EU, 'European Union NextGenerationEU/PRTR', and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, the National Science, Research and Innovation Fund program IND_FF_68_369_2300_097, and the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, Grant B39G680009 (Thailand); the Eric & Wendy Schmidt Fund for Strategic Innovation through the CERN Next Generation Triggers Project under Grant Agreement No. SIF-2023-004; the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, Contract C-1845; and the Weston Havens Foundation (USA).

Data availability statement

Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the CMS data preservation, re-use and open access policy.

Author contribution

The CMS Collaboration
Conceptualization (equal), Data curation (equal), Formal analysis (equal), Funding acquisition (equal), Investigation (equal), Methodology (equal), Project administration (equal), Resources (equal), Software (equal), Supervision (equal), Validation (equal), Visualization (equal), Writing – original draft (equal), Writing – review & editing (equal)

Appendix. Comparison of SMEFT constraints with a simplified likelihood procedure

The SMEFT constraints presented in section 11 are derived from the full combination likelihood. This likelihood function, constructed following the methodology described in section 5, encodes the signal yields in each analysis region as functions of the SMEFT parameters. Formulating the statistical model in this way ensures the most accurate representation of the data. For example, all non-Gaussian effects are preserved, and correlations between NPs across different analysis regions are fully incorporated. However, the full combination likelihood is complex and computationally intensive to evaluate. In this appendix, a simplified likelihood procedure is used to extract the SMEFT constraints, and the results are compared with those obtained using the full combination likelihood.

The simplified likelihood procedure follows the prescription detailed in [126]. Its inputs are taken from the 97 POI fit to the parameters representing the products of cross sections and branching fractions (figure 12). The best fit values $\hat{\mu}_I$ and 68% CL intervals σ_I, defined relative to the SM predictions, are used to construct the simplified likelihood, where the index I runs over the 97 POIs. The numerical values are those listed in table 8. In addition, the correlations between the POIs ρ_{IJ} are required, which are provided as supplementary material to this paper [41].

A linear approximation of the likelihood is constructed as follows. The measurements μ_I are transformed into new variables χ_I that follow a standard normal distribution,

$$\mu_I = a_I + b_I \chi_I, \tag{A.1}$$

such that

$$\chi_I = (\mu_I - a_I)/b_I. \tag{A.2}$$

From this, the parameters a_I and b_I can be identified as $\hat{\mu}_I$ and σ_I, respectively, to arrive at the usual definition of a χ^2 test statistic,

$$\chi^2(\vec{\mu}) = \tilde{\rho}_{IJ}^{-1}\chi_I\chi_J, \tag{A.3}$$

where a summation over repeated indices is implied. Note this expression is equivalent, up to an additive constant, to $-2\ln\mathcal{L}(\vec{\mu})$ obtained under the assumption that the measurements are distributed according to a multivariate Gaussian distribution. The term $\tilde{\rho}_{IJ}$ represents the correlation matrix between the χ_I variables, which in the linear approximation is equal to the correlation matrix between the μ_I parameters, ρ_{IJ}. Furthermore, in the linear approximation, the 68% CL intervals are symmetric and are obtained by averaging the upper and lower 68% CL intervals in the measurements, $\sigma_I = (\sigma_I^+ + \sigma_I^-)/2$. The parameters from the $H \to \gamma\gamma$ and $H \to ZZ$ decay channels are restricted to nonnegative values in the 97 POI fit, and therefore the lower 68% CL intervals can be truncated. In these cases, the upper 68% CL interval is used as a conservative estimate of the symmetrized uncertainty, $\sigma_I = \sigma_I^+$.

As a result of the limited size of the data set and the asymmetric impact of systematic uncertainties, many of the measurements in the 97 POI fit deviate from the asymptotic Gaussian limit. This can be seen by considering the asymmetric 68% CL intervals in table 8, where $\sigma_I^+ \neq \sigma_I^-$. The asymmetry can be accounted for in the simplified likelihood by including the nonlinear next-to-leading term in the expansion,

$$\mu_I = a_I + b_I\chi_I + c_I\chi_I^2, \tag{A.4}$$

where the additional parameter c_I incorporates skewness, and is typically much smaller than b_I. Inverting this expression yields a quadratic formula for χ_I,

$$\chi_I = \frac{\sqrt{b_I^2 - 4c_I(a_I - \mu_I)} - b_I}{2c_I}. \tag{A.5}$$

This formula can then be used in equation (A.3) to construct a test statistic that accounts for asymmetric uncertainties. Because of the presence of a square root, χ_I can become imaginary. However, as long as c_I is small compared to the linear term b_I, this only occurs far from the best fit value, where $\chi^2(\vec{\mu}) \gg 1$, which does not cause a problem for the extraction of results. In the asymmetric uncertainty model, the correlation matrix $\tilde{\rho}_{IJ}$ is no longer equal to ρ_{IJ}, because of the small contribution from the quadratic terms.

The values of a_I, b_I, c_I, and $\tilde{\rho}_{IJ}$ are determined using the moment matching procedure described in [126]. The first three moments for measurement μ_I can be estimated using $\hat{\mu}_I$, σ_I^+, and σ_I^-, from the dimidated Gaussian distribution [127], which joins two half-Gaussian distributions at the mode, with different widths σ_I^+ and σ_I^- on either side. The code used to extract the third moment is taken from the SPEY tool, and is described in appendix C of [128]. As with the symmetric case, for the $H \to \gamma\gamma$ and $H \to ZZ$ parameters with truncated lower 68% CL intervals, the upper 68% CL interval is used for both σ_I^+ and σ_I^- to provide a conservative symmetric approximation. The calculated moments, along with ρ_{IJ}, are then used to solve for a_I, b_I, c_I, and $\tilde{\rho}_{IJ}$ using formulae 2.9–2.12 from [126]. As an example, the a_I, b_I, and c_I parameters for the (ggH 0J $p_T^H < 10\,\text{GeV}$, $H \to \gamma\gamma$) POI are determined to be 0.58, 0.32, and 0.06, respectively. This demonstrates that even for a measurement with a significant asymmetry ($0.58^{+0.38}_{-0.26}$), the quadratic term c_I is small compared to the linear term b_I.

Constraints on the SMEFT eigenvectors $\vec{\text{EV}}$ are then extracted by replacing μ_I in the χ_I formulae with the corresponding SMEFT parametrization $\mu_I(\vec{\text{EV}})$ (shown in figures 26–27), thereby obtaining χ^2 as a function of $\vec{\text{EV}}$. A new test statistic is defined as,

$$q_{\text{simplified}}\left(\vec{\text{EV}}\right) = \chi^2\left(\vec{\text{EV}}\right) - \chi^2\left(\hat{\vec{\text{EV}}}\right), \tag{A.6}$$

where $\hat{\vec{\text{EV}}}$ are the best fit values that minimize $\chi^2(\vec{\text{EV}})$. This test statistic provides an approximation of the profile likelihood ratio test statistic q used in the full combined likelihood fit, and can likewise be used to derive SMEFT constraints following the procedure outlined in section 5.3. All results for the simplified likelihood fits were extracted using a single 8-core CPU in approximately five minutes. In contrast, the full likelihood fit required $\mathcal{O}(50000)$ CPU hours on a batch system.



Figure A1. Comparison of the constraints on the linear combinations of SMEFT WCs for different fit strategies. The results shown in black are obtained using the full combination likelihood, and are the same as those shown in figure 28. The results in orange and red are obtained with a simplified likelihood procedure, constructed from the 97 POI fit to STXS measurements (result shown in figure 12). The results shown in orange are based on a symmetric uncertainty model, while the results shown in red account for asymmetric uncertainties. The left panel shows the observed best fit values, and 68% and 95% CL intervals, as well as the expected 95% CL intervals. In the right panel, the results are translated into a 95% CL lower limit on the BSM physics energy scale, assuming $\mathrm{EV}_j = 1$. The eigenvectors are listed in order of the expected excluded energy scale from the full likelihood fit. The fits use a linear SMEFT parametrization, in which terms up to $\mathcal{O}(\mathrm{EV}/\Lambda^2)$ are included.

Figure A1 compares the SMEFT constraints obtained using the simplified likelihood procedure with those obtained using the full combination likelihood (black). The simplified likelihood results are shown for both the symmetric (orange) and asymmetric (red) uncertainty models, obtained with the linear and nonlinear approximations of the likelihood, respectively. The left panel shows the observed best fit values, and 68% and 95% CL intervals, in addition to the expected 95% CL intervals. The right panel shows the results translated into 95% CL lower limits on the energy scale of new physics Λ_j, assuming $\mathrm{EV}_j = 1$.

Overall, the results obtained with the simplified likelihood show good agreement with those from the full combination likelihood. Between the two implementations, the asymmetric uncertainty model provides a better approximation of the full likelihood results, motivating the inclusion of asymmetric uncertainties in future simplified likelihood fits. In general, the simplified likelihood provides a reliable

estimate of the sensitivity, although a few notable shifts are observed in the best fit values. For example, the best fit value for EV_2 shifts from 0.036 using the full likelihood, to -0.022 (-0.024) using the simplified likelihood with the symmetric (asymmetric) uncertainty model. This eigenvector is linked to the qqH, VH, and ttH STXS bin measurements in the $H \to ZZ$ decay channel (shown in figure 26). These parameters exhibit highly asymmetric (and sometimes truncated) 68% CL intervals, which the simplified likelihood does not accurately capture. In most cases, the shifts in the best fit values are within the 68% CL intervals of the full likelihood. However, with increased sensitivity, such shifts could lead to false conclusions about deviations from the SM.

The differences between the simplified likelihood results and full likelihood results can be attributed to several factors. First, the measured parameters μ_I do not follow an exact (dimidated) Gaussian distribution. In addition, the parameters from the 97 POI fit are defined according to a scheme in which some of the STXS bins are merged, whereas the full combination likelihood parametrizes the signal according to the full STXS stage 1.2 binning. The simplified likelihood also neglects the effect of NPs, which can impact the constraints obtained with the full likelihood. Finally, special treatment is required in the simplified likelihood procedure for parameters with truncated 68% CL intervals; a complication that is absent in the full likelihood fit. Therefore, while the simplified likelihood procedure provides a useful approximation, the full combination likelihood is the most accurate representation of the data. The caveats discussed above should be considered when using the simplified likelihood procedure in future SMEFT interpretations.

The CMS Collaboration

C Yang
State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, People's Republic of China

A Hayrapetyan, V Makarenko, A Tumasyan[1]
Yerevan Physics Institute, Yerevan, Armenia

W Adam, L Benato, T Bergauer, M Dragicevic, P S Hussain, M Jeitler[2], N Krammer, A Li, D Liko, M Matthewman, J Schieck[2], R Schöfbeck[2], M Shooshtari, M Sonawane, W Waltenberger, C-E Wulz[2]
Institut für Hochenergiephysik, Vienna, Austria

T Janssen, H Kwon, D Ocampo Henao, T Van Laer, P Van Mechelen
Universiteit Antwerpen, Antwerpen, Belgium

J Bierkens, N Breugelmans, J D'Hondt, S Dansana, A De Moor, M Delcourt, F Heyen, Y Hong, P Kashko, S Lowette, I Makarenko, D Müller, S Tavernier, M Tytgat[3], G P Van Onsem, S Van Putte, D Vannerom
Vrije Universiteit Brussel, Brussel, Belgium

B Bilin, B Clerbaux, A K Das, I De Bruyn, G De Lentdecker, H Evard, L Favart, P Gianneios, A Khalilzadeh, F A Khan, A Malara, M A Shahzad, A Sharma, L Thomas, M Vanden Bemden, C Vander Velde, P Vanlaer, F Zhang
Université Libre de Bruxelles, Bruxelles, Belgium

M De Coen, D Dobur, C Giordano, G Gokbulut, K Kaspar, D Kavtaradze, D Marckx, K Skovpen, A M Tomaru, N Van Den Bossche, J van der Linden, J Vandenbroeck
Ghent University, Ghent, Belgium

H Aarup Petersen, S Bein, A Benecke, A Bethani, G Bruno, A Cappati, J De Favereau De Jeneret, C Delaere, F Gameiro Casalinho, A Giammanco, A O Guzel, V Lemaitre, J Lidrych, P Malek, P Mastrapasqua, S Turkcapar
Université Catholique de Louvain, Louvain-la-Neuve, Belgium

G A Alves, M Barroso Ferreira Filho, E Coelho, C Hensel, D Matos Figueiredo, T Menezes De Oliveira, C Mora Herrera, P Rebello Teles, M Soeiro, E J Tonelli Manganote[4], A Vilela Pereira
Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

W L Aldá Júnior, H Brandao Malbouisson, W Carvalho, J Chinellato[5], M Costa Reis, E M Da Costa, G G Da Silveira[6], D De Jesus Damiao, S Fonseca De Souza, R Gomes De Souza, S S Jesus, T Laux Kuhn[6], M Macedo, K Mota Amarilo, L Mundim, H Nogima, J P Pinheiro, A Santoro, A Sznajder, M Thiel, F Torres Da Silva De Araujo[7]
Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

C A Bernardes[6], L Calligaris, F Damas, T R Fernandez Perez Tomei, E M Gregores, B Lopes Da Costa, I Maietto Silverio, P G Mercadante, S F Novaes, Sandra S Padula, V Scheurer
Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

A Aleksandrov, G Antchev, P Danev, R Hadjiiska, P Iaydjiev, M Shopova, G Sultanov
Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A Dimitrov, L Litov, B Pavlov, P Petkov, A Petrov
University of Sofia, Sofia, Bulgaria

S Keshri, D Laroze, S Thakur
Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

W Brooks
Universidad Tecnica Federico Santa Maria, Valparaiso, Chile

T Cheng, T Javaid, L Wang, L Yuan
Beihang University, Beijing, People's Republic of China

Z Hu, Z Liang, J Liu, X Wang, H Yang
Department of Physics, Tsinghua University, Beijing, People's Republic of China

G M Chen[8], H S Chen[8], M Chen[8], Y Chen, Q Hou, X Hou, F Iemmi, C H Jiang, H Liao, G Liu, Z-A Liu[9], J N Song[9], S Song, J Tao, C Wang[8], J Wang, H Zhang, J Zhao
Institute of High Energy Physics, Beijing, People's Republic of China

A Agapitos, Y Ban, A Carvalho Antunes De Oliveira, S Deng, B Guo, Q Guo, C Jiang, A Levin, C Li, Q Li, Y Mao, S Qian, S J Qian, X Qin, C Quaranta, X Sun, D Wang, J Wang, C Yang, M Zhang, Y Zhao, C Zhou
State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, People's Republic of China

S Yang
State Key Laboratory of Nuclear Physics and Technology, Institute of Quantum Matter, South China Normal University, Guangzhou, People's Republic of China

Z You
Sun Yat-Sen University, Guangzhou, People's Republic of China

N Lu
University of Science and Technology of China, Hefei, People's Republic of China

G Bauer[10,11], Z Cui[11], B Li[12], H Wang, K Yi[13], J Zhang
Nanjing Normal University, Nanjing, People's Republic of China

Y Li, Y Zhou[14]
Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE)—Fudan University, Shanghai, People's Republic of China

Z Lin, C Lu, M Xiao[15]
Zhejiang University, Hangzhou, Zhejiang, People's Republic of China

C Avila, D A Barbosa Trujillo, A Cabrera, C Florez, J Fraga, J A Reyes Vega
Universidad de Los Andes, Bogota, Colombia

C Rendón, M Rodriguez, A A Ruales Barbosa, J D Ruiz Alvarez
Universidad de Antioquia, Medellin, Colombia

N Godinovic, D Lelas, A Sculac
University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

M Kovac, A Petkovic, T Sculac
University of Split, Faculty of Science, Split, Croatia

P Bargassa, V Brigljevic, B K Chitroda, D Ferencek, K Jakovcic, A Starodumov, T Susa
Institute Rudjer Boskovic, Zagreb, Croatia

A Attikis, K Christoforou, S Konstantinou, C Leonidou, L Paizanos, F Ptochos, P A Razis, H Rykaczewski, H Saka, A Stepennov
University of Cyprus, Nicosia, Cyprus

M Finger[†], M Finger Jr
Charles University, Prague, Czech Republic

E Ayala
Escuela Politecnica Nacional, Quito, Ecuador

E Carrera Jarrin
Universidad San Francisco de Quito, Quito, Ecuador

A A Abdelalim[16,17], R Aly[18,16]
Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

A Hussein, H Mohammed
Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

K Jaffel, M Kadastik, T Lange, C Nielsen, J Pata, M Raidal, N Seeba, L Tani
National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

E Brücken, A Milieva, K Osterberg, M Voutilainen
Department of Physics, University of Helsinki, Helsinki, Finland

F Garcia, P Inkaew, K T S Kallonen, R Kumar Verma, T Lampén, K Lassila-Perini, B Lehtela, S Lehti, T Lindén, N R Mancilla Xinto, M Myllymäki, M M Rantanen, S Saariokari, N T Toikka, J Tuominiemi
Helsinki Institute of Physics, Helsinki, Finland

N Bin Norjoharuddeen, H Kirschenmann, P Luukka, H Petrow
Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

M Besancon, F Couderc, M Dejardin, D Denegri, P Devouge, J L Faure, F Ferri, P Gaigne, S Ganjour, P Gras, F Guilloux, G Hamel de Monchenault, M Kumar, V Lohezic, Y Maidannyk, J Malcles, F Orlandi, L Portales, S Ronchi, M Ö Sahin, A Savoy-Navarro[19], P Simkina, M Titov, M Tornago
IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

R Amella Ranz, F Beaudette, G Boldrini, P Busson, C Charlot, M Chiusi, T D Cuisset, O Davignon, A De Wit, T Debnath, I T Ehle, S Ghosh, A Gilbert, R Granier de Cassagnac, L Kalipoliti, M Manoni, M Nguyen, S Obraztsov, C Ochando, R Salerno, J B Sauvan, Y Sirois, G Sokmen, Y Song, L Urda Gómez, A Zabi, A Zghiche

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

J-L Agram[20], J Andrea, D Bloch, J-M Brom, E C Chabert, C Collard, G Coulon, S Falke, U Goerlach, R Haeberle, A-C Le Bihan, M Meena, O Poncet, G Saha, P Vaucelle
Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

A Di Florio, B Orzari
Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

D Amram, S Beauceron, B Blancon, G Boudoul, N Chanon, D Contardo, P Depasse, H El Mamouni, J Fay, E Fillaudeau, S Gascon, M Gouzevitch, C Greenberg, G Grenier, B Ille, E Jourd'Huy, M Lethuillier, B Massoteau, L Mirabito, A Purohit, M Vander Donckt, J Xiao
Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D Chokheli, I Lomidze, Z Tsamalaidze[21]
Georgian Technical University, Tbilisi, Georgia

V Botta, S Consuegra Rodríguez, L Feld, K Klein, M Lipinski, P Nattland, V Oppenländer, A Pauls, D Pérez Adán, N Röwert
RWTH Aachen University, Idot Physikalisches Institut, Aachen, Germany

C Daumann, S Diekmann, N Eich, D Eliseev, F Engelke, J Erdmann, M Erdmann, B Fischer, T Hebbeker, K Hoepfner, F Ivone, A Jung, N Kumar, M Y Lee, F Mausolf, M Merschmeyer, A Meyer, A Pozdnyakov, W Redjeb, H Reithler, U Sarkar, V Sarkisovi, A Schmidt, C Seth, A Sharma, J L Spah, V Vaulin, S Zaleski
RWTH Aachen University, IIIdot Physikalisches Institut A, Aachen, Germany

M R Beckers, C Dziwok, G Flügge, N Hoeflich, T Kress, A Nowack, O Pooth, A Stahl, A Zotz
RWTH Aachen University, IIIdot Physikalisches Institut B, Aachen, Germany

A Abel, M Aldaya Martin, J Alimena, S Amoroso, Y An, I Andreev, J Bach, S Baxter, H Becerril Gonzalez, O Behnke, A Belvedere, F Blekman[22], K Borras[23], A Campbell, S Chatterjee, L X Coll Saravia, G Eckerlin, D Eckstein, E Gallo[22], A Geiser, M Guthoff, A Hinzmann, L Jeppe, M Kasemann, C Kleinwort, R Kogler, M Komm, D Krücker, W Lange, D Leyva Pernia, K-Y Lin, K Lipka[24], W Lohmann[25], J Malvaso, R Mankel, I-A Melzer-Pellmann, M Mendizabal Morentin, A B Meyer, G Milella, K Moral Figueroa, A Mussgiller, L P Nair, J Niedziela, A Nürnberg, J Park, E Ranken, A Raspereza, D Rastorguev, L Rygaard, M Scham[26,23], S Schnake[23], P Schütze, C Schwanenberger[22], D Schwarz, D Selivanova, K Sharko, M Shchedrolosiev, D Stafford, M Torkian, A Ventura Barroso, R Walsh, D Wang, Q Wang, K Wichmann, L Wiens[23], C Wissing, Y Yang, S Zakharov, A Zimermmane Castro Santos
Deutsches Elektronen-Synchrotron, Hamburg, Germany

A R Alves Andrade, M Antonello, S Bollweg, M Bonanomi, L Ebeling, K El Morabit, Y Fischer, M Frahm, E Garutti, A Grohsjean, A A Guvenli, J Haller, D Hundhausen, G Kasieczka, P Keicher, R Klanner, W Korcari, T Kramer, C C Kuo, F Labe, J Lange, A Lobanov, J Matthiesen, L Moureaux, K Nikolopoulos, A Paasch, K J Pena Rodriguez, N Prouvost, B Raciti, M Rieger, D Savoiu, P Schleper, M Schröder, J Schwandt, M Sommerhalder, H Stadie, G Steinbrück, R Ward, B Wiederspan, M Wolf, C Yede
University of Hamburg, Hamburg, Germany

S Brommer, A Brusamolino, E Butz, Y M Chen, T Chwalek, A Dierlamm, G G Dincer, D Druzhkin, U Elicabuk, N Faltermann, M Giffels, A Gottmann, F Hartmann[27], M Horzela, F Hummer, U Husemann, J Kieseler, M Klute, J Knolle, R Kunnilan Muhammed Rafeek, O Lavoryk, J M Lawhorn, S Maier, A A Monsch, M Mormile, Th Müller, E Pfeffer, M Presilla, G Quast, K Rabbertz, B Regnery, R Schmieder, N Shadskiy,

I Shvetsov, H J Simonis, L Sowa, L Stockmeier, K Tauqeer, M Toms, B Topko, N Trevisani, C Verstege, T Voigtländer, R F Von Cube, J Von Den Driesch, C Winter, R Wolf, W D Zeuner, X Zuo
Karlsruher Institut fuer Technologie, Karlsruhe, Germany

G Anagnostou, G Daskalakis, A Kyriakis
Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G Melachroinos, Z Painesis, I Paraskevas, N Saoulidou, K Theofilatos, E Tziaferi, E Tzovara, K Vellidis, I Zisopoulos
National and Kapodistrian University of Athens, Athens, Greece

T Chatzistavrou, G Karapostoli, K Kousouris, E Siamarkou, G Tsipolitis
National Technical University of Athens, Athens, Greece

I Bestintzanos, I Evangelou, C Foudas, P Katsoulis, P Kokkas, P G Kosmoglou Kioseoglou, N Manthos, I Papadopoulos, J Strologas
University of Ioánnina, Ioánnina, Greece

C Hajdu, D Horvath[28,29], Á Kadlecsik, K Márton, A J Rádl[30], F Sikler, V Veszpremi
HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

M Csanád, K Farkas, A Fehérkuti[31], M M A Gadallah[32], M León Coello, G Pásztor, G I Veres
MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

B Ujvari, G Zilizi
Faculty of Informatics, University of Debrecen, Debrecen, Hungary

G Bencze, S Czellar, J Molnar, Z Szillasi
HUN-REN ATOMKI—Institute of Nuclear Research, Debrecen, Hungary

T Csorgo[31], F Nemes[31], T Novak, I Szanyi[33]
Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

S Bahinipati, S Nayak, R Raturi
IIT Bhubaneswar, Bhubaneswar, India

S Bansal, S B Beri, V Bhatnagar, S Chauhan, N Dhingra[34], A Kaur, H Kaur, M Kaur, S Kumar, T Sheokand, J B Singh, A Singla
Panjab University, Chandigarh, India

A Bhardwaj, A Chhetri, B C Choudhary, A Kumar, A Kumar, M Naimuddin, S Phor, K Ranjan, M K Saini
University of Delhi, Delhi, India

P Palni
Indian Institute of Technology Mandi (IIT-Mandi), Himachal Pradesh, India

S Acharya[35], B Gomber
University of Hyderabad, Hyderabad, India

S Mukherjee
Indian Institute of Technology Kanpur, Kanpur, India

S Bhattacharya, S Das Gupta, S Dutta, S Dutta, S Sarkar
Saha Institute of Nuclear Physics, HBNI, Kolkata, India

M M Ameen, P K Behera, S Chatterjee, G Dash, A Dattamunsi, P Jana, P Kalbhor, S Kamble, J R Komaragiri[36], T Mishra, P R Pujahari, A K Sikdar, R K Singh, P Verma, S Verma, A Vijay
Indian Institute of Technology Madras, Madras, India

B K Sirasva
IISER Mohali, India, Mohali, India

L Bhatt, S Dugad, G B Mohanty, M Shelake, P Suryadevara
Tata Institute of Fundamental Research-A, Mumbai, India

A Bala, S Banerjee, S Barman[37], R M Chatterjee, M Guchait, Sh Jain, A Jaiswal, S Kumar, M Maity[37], G Majumder, K Mazumdar, S Parolia, R Saxena, A Thachayath
Tata Institute of Fundamental Research-B, Mumbai, India

D Maity[38], P Mal, K Naskar[38], A Nayak[38], K Pal, P Sadangi, S K Swain, S Varghese[38], D Vats[38]
National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S Dube, P Hazarika, B Kansal, A Laha, R Sharma, S Sharma, K Y Vaish
Indian Institute of Science Education and Research (IISER), Pune, India

S Ghosh
Indian Institute of Technology Hyderabad, Telangana, India

H Bakhshiansohi[39], A Jafari[40], V Sedighzadeh Dalavi, M Zeinali[41]
Isfahan University of Technology, Isfahan, Iran

S Bashiri, S Chenarani[42], S M Etesami, Y Hosseini, M Khakzad, E Khazaie, M Mohammadi Najafabadi, S Tizchang[43]
Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

M Felcini, M Grunewald
University College Dublin, Dublin, Ireland

M Abbrescia[a,b], M Barbieri[a,b], M Buonsante[a,b], A Colaleo[a,b], D Creanza[a,c], N De Filippis[a,c], M De Palma[a,b], W Elmetenawee[a,b,16], N Ferrara[a,c], L Fiore[a], L Generoso[a,b], L Longo[a], M Louka[a,b], G Maggi[a,c], M Maggi[a], I Margjeka[a], V Mastrapasqua[a,b], S My[a,b], F Nenna[a,b], S Nuzzo[a,b], A Pellecchia[a,b], A Pompili[a,b], G Pugliese[a,c], R Radogna[a,b], D Ramos[a], A Ranieri[a], L Silvestris[a], F M Simone[a,c], Ü Sözbilir[a], A Stamerra[a,b], D Troiano[a,b], R Venditti[a,b], P Verwilligen[a], A Zaza[a,b]
INFN Sezione di Bari[a], Università di Bari[b], Politecnico di Bari[c], Bari, Italy

G Abbiendi[a], C Battilana[a,b], D Bonacorsi[a,b], P Capiluppi[a,b], F R Cavallo[a], M Cuffiani[a,b], G M Dallavalle[a], T Diotalevi[a,b], F Fabbri[a], A Fanfani[a,b], R Farinelli[a], D Fasanella[a], P Giacomelli[a], C Grandi[a], L Guiducci[a,b], S Lo Meo[a,44], M Lorusso[a,b], L Lunerti[a], S Marcellini[a], G Masetti[a], F L Navarria[a,b], G Paggi[a,b], A Perrotta[a], A M Rossi[a,b], S Rossi Tisbeni[a,b]
INFN Sezione di Bologna[a], Università di Bologna[b], Bologna, Italy

S Costa[a,b,45], A Di Mattia[a], A Lapertosa[a], R Potenza[a,b], A Tricomi[a,b,45]
INFN Sezione di Catania[a], Università di Catania[b], Catania, Italy

J Altork[a,b], P Assiouras[a], G Barbagli[a], G Bardelli[a], M Bartolini[a,b], A Calandri[a,b], B Camaiani[a,b], A Cassese[a], R Ceccarelli[a], V Ciulli[a,b], C Civinini[a], R D'Alessandro[a,b], L Damenti[a,b], E Focardi[a,b], T Kello[a], G Latino[a,b], P Lenzi[a,b], M Lizzo[a], M Meschini[a], S Paoletti[a], A Papanastassiou[a,b], G Sguazzoni[a], L Viliani[a]
INFN Sezione di Firenze[a], Università di Firenze[b], Firenze, Italy

L Benussi, S Colafranceschi[46], S Meola[47], D Piccolo
INFN Laboratori Nazionali di Frascati, Frascati, Italy

M Alves Gallo Pereira[a], F Ferro[a], E Robutti[a], S Tosi[a,b]
INFN Sezione di Genova[a], Università di Genova[b], Genova, Italy

A Benaglia[a], F Brivio[a], V Camagni[a,b], F Cetorelli[a,b], F De Guio[a,b], M E Dinardo[a,b], P Dini[a], S Gennai[a], R Gerosa[a,b], A Ghezzi[a,b], P Govoni[a,b], L Guzzi[a], M R Kim[a], G Lavizzari[a,b], M T Lucchini[a,b], M Malberti[a], S Malvezzi[a], A Massironi[a], D Menasce[a], L Moroni[a], M Paganoni[a,b], S Palluotto[a,b], D Pedrini[a], A Perego[a,b], G Pizzati[a,b], T Tabarelli de Fatis[a,b]
INFN Sezione di Milano-Bicocca[a], Università di Milano-Bicocca[b], Milano, Italy

S Buontempo[a], C Di Fraia[a,b], F Fabozzi[a,c], L Favilla[a,d], A O M Iorio[a,b], L Lista[a,b,48], P Paolucci[a,27], B Rossi[a]
INFN Sezione di Napoli[a], Università di Napoli 'Federico II'[b], Napoli, Italy; Università della Basilicata[c], Potenza, Italy; Scuola Superiore Meridionale (SSM)[d], Napoli, Italy

P Azzi[a], N Bacchetta[a,49], D Bisello[a,b], L Borella[a], P Bortignon[a,c], G Bortolato[a,b], A C M Bulla[a,c], R Carlin[a,b], T Dorigo[a,50], S Fantinel[a], F Gasparini[a,b], U Gasparini[a,b], S Giorgetti[a], E Lusiani[a], M Margoni[a,b], A T Meneguzzo[a,b], J Pazzini[a,b], F Primavera[a,b], P Ronchese[a,b], R Rossin[a,b], F Simonetto[a,b], M Tosi[a,b], A Triossi[a,b], S Ventura[a], P Zotto[a,b], A Zucchetta[a,b], G Zumerle[a,b]
INFN Sezione di Padova[a], Università di Padova[b], Padova, Italy; Universita degli Studi di Cagliari[c], Cagliari, Italy

A Braghieri[a], M Brunoldi[a,b], S Calzaferri[a,b], P Montagna[a,b], M Pelliccioni[a,b], V Re[a], C Riccardi[a,b], P Salvini[a], I Vai[a,b], P Vitulo[a,b]
INFN Sezione di Pavia[a], Università di Pavia[b], Pavia, Italy

S Ajmal[a,b], M E Ascioti[a,b], G M Bilei[†,a], C Carrivale[a,b], D Ciangottini[a,b], L Della Penna[a,b], L Fanò[a,b], V Mariani[a,b], M Menichelli[a], F Moscatelli[a,51], A Rossi[a,b], A Santocchia[a,b], D Spiga[a], T Tedeschi[a,b]
INFN Sezione di Perugia[a], Università di Perugia[b], Perugia, Italy

C Aimè[a,b], C A Alexe[a,c], P Asenov[a,b], P Azzurri[a], G Bagliesi[a], L Bianchini[a,b], T Boccali[a], E Bossini[a], D Bruschini[a,c], R Castaldi[a], F Cattafesta[a,c], M A Ciocci[a,d], M Cipriani[a,b], R Dell'Orso[a], S Donato[a,b], R Forti[a,b], A Giassi[a], F Ligabue[a,c], A C Marini[a,b], A Messineo[a,b], S Mishra[a], V K Muraleedharan Nair Bindhu[a,b], S Nandan[a], F Palla[a], M Riggirello[a,c], A Rizzi[a,b], G Rolandi[a,c], S Roy Chowdhury[a,52], T Sarkar[a], A Scribano[a], P Solanki[a,b], P Spagnolo[a], F Tenchini[a,b], R Tenchini[a], G Tonelli[a,b], N Turini[a,d], F Vaselli[a,c], A Venturi[a], P G Verdini[a]
INFN Sezione di Pisa[a], Università di Pisa[b], Scuola Normale Superiore di Pisa[c], Pisa, Italy; Università di Siena[d], Siena, Italy

P Akrap[a,b], C Basile[a,b], S C Behera[a], F Cavallari[a], L Cunqueiro Mendez[a,b], F De Riggi[a,b], D Del Re[a,b], M Del Vecchio[a,b], E Di Marco[a], M Diemoz[a], F Errico[a], L Frosina[a,b], R Gargiulo[a,b], B Harikrishnan[a,b], F Lombardi[a,b], E Longo[a,b], L Martikainen[a,b], J Mijuskovic[a,b], G Organtini[a,b], N Palmeri[a,b], R Paramatti[a,b], T Pauletto[a,b], S Rahatlou[a,b], C Rovelli[a], F Santanastasio[a,b], L Soffi[a], V Vladimirov[a,b]
INFN Sezione di Roma[a], Sapienza Università di Roma[b], Roma, Italy

N Amapane[a,b], R Arcidiacono[a,c], S Argiro[a,b], M Arneodo[†,a,c], N Bartosik[a,c], R Bellan[a,b], A Bellora[a,b], C Biino[a], C Borca[a,b], N Cartiglia[a], M Costa[a,b], R Covarelli[a,b], P De Remigis[a], N Demaria[a], M Ferrero[a], L Finco[a], M Grippo[a,b], B Kiani[a,b], L Lanteri[a,b], F Legger[a], F Luongo[a,b], C Mariotti[a], S Maselli[a], A Mecca[a,b], L Menzio[a,b], P Meridiani[a], E Migliore[a,b], M Monteno[a], M M Obertino[a,b], G Ortona[a], L Pacher[a,b], N Pastrone[a], M Ruspa[a,c], F Siviero[a,b], V Sola[a,b], A Solano[a,b], C Tarricone[a,b], D Trocino[a], G Umoret[a,b], R White[a,b]

INFN Sezione di Torino[a], Università di Torino[b], Torino, Italy; Università del Piemonte Orientale[c], Novara, Italy

J Babbar[a,b,52], S Belforte[a], V Candelise[a,b], M Casarsa[a], F Cossutti[a], K De Leo[a], G Della Ricca[a,b], R Delli Gatti[a,b]
INFN Sezione di Trieste[a], Università di Trieste[b], Trieste, Italy

S Dogra, J Hong, J Kim, T Kim, D Lee, H Lee, J Lee, S W Lee, C S Moon, Y D Oh, S Sekmen, B Tae, Y C Yang
Kyungpook National University, Daegu, Republic of Korea

M S Kim
Department of Mathematics and Physics—GWNU, Gangneung, Republic of Korea

G Bak, P Gwak, H Kim, D H Moon, J Seo
Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Republic of Korea

E Asilar, F Carnevali, J Choi[53], T J Kim, Y Ryou, J Song
Hanyang University, Seoul, Republic of Korea

S Ha, S Han, B Hong, J Kim, K Lee, K S Lee, S Lee, J Yoo
Korea University, Seoul, Republic of Korea

J Goh, J Shin, S Yang
Kyung Hee University, Department of Physics, Seoul, Republic of Korea

Y Kang, H S Kim, Y Kim, B Ko, S Lee
Sejong University, Seoul, Republic of Korea

J Almond, J H Bhyun, J Choi, J Choi, W Jun, H Kim, J Kim, T Kim, Y Kim, Y W Kim, S Ko, H Lee, J Lee, J Lee, B H Oh, J Shin, U K Yang, I Yoon
Seoul National University, Seoul, Republic of Korea

W Jang, D Kim, S Kim, J S H Lee, Y Lee, I C Park, Y Roh, I J Watson
University of Seoul, Seoul, Republic of Korea

G Cho, K Hwang, B Kim, S Kim, K Lee, H D Yoo
Yonsei University, Department of Physics, Seoul, Republic of Korea

Y Lee, I Yu
Sungkyunkwan University, Suwon, Republic of Korea

T Beyrouthy, Y Gharbia
College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

F Alazemi
Kuwait University—College of Science—Department of Physics, Safat, Kuwait

K Dreimanis, O M Eberlins, A Gaile, C Munoz Diaz, D Osite, G Pikurs, R Plese, A Potrebko, M Seidel, D Sidiropoulos Kontos
Riga Technical University, Riga, Latvia

N R Strautnieks
University of Latvia (LU), Riga, Latvia

M Ambrozas, A Juodagalvis, S Nargelas, A Rinkevicius, G Tamulaitis
Vilnius University, Vilnius, Lithuania

I Yusuff[54], Z Zolkapli
National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

J F Benitez, A Castaneda Hernandez, A Cota Rodriguez, L E Cuevas Picos, H A Encinas Acosta, L G Gallegos Maríñez, J A Murillo Quijada, L Valencia Palomo
Universidad de Sonora (UNISON), Hermosillo, Mexico

G Ayala, H Castilla-Valdez, H Crotte Ledesma, R Lopez-Fernandez, J Mejia Guisao, R Reyes-Almanza, A Sánchez Hernández
Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

C Oropeza Barrera, D L Ramirez Guadarrama, M Ramírez García
Universidad Iberoamericana, Mexico City, Mexico

I Bautista, F E Neri Huerta, I Pedraza, H A Salazar Ibarguen, C Uribe Estrada
Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I Bubanja, N Raicevic
University of Montenegro, Podgorica, Montenegro

P H Butler
University of Canterbury, Christchurch, New Zealand

A Ahmad, M I Asghar, A Awais, M I M Awan, W A Khan
National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

V Avati, L Forthomme, L Grzanka, M Malawski, K Piotrzkowski
AGH University of Krakow, Krakow, Poland

M Bluj, M Górski, M Kazana, M Szleper, P Zalewski
National Centre for Nuclear Research, Swierk, Poland

K Bunkowski, K Doroba, A Kalinowski, M Konecki, J Krolikowski, A Muhammad
Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

P Fokow, K Pozniak, W Zabolotny
Warsaw University of Technology, Warsaw, Poland

M Araujo, D Bastos, C Beirão Da Cruz E Silva, A Boletti, M Bozzo, T Camporesi, G Da Molin, M Gallinaro, J Hollar, N Leonardo, G B Marozzo, A Petrilli, M Pisano, J Seixas, J Varela, J W Wulff
Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

P Adzic, L Markovic, P Milenovic, V Milosevic
Faculty of Physics, University of Belgrade, Belgrade, Serbia

D Devetak, M Dordevic, J Milosevic, L Nadderd, V Rekovic, M Stojanovic
VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

M Alcalde Martinez, J Alcaraz Maestre, Cristina F Bedoya, J A Brochero Cifuentes, Oliver M Carretero, M Cepeda, M Cerrada, N Colino, B De La Cruz, A Delgado Peris, A Escalante Del Valle, D Fernández Del Val, J P Fernández Ramos, J Flix, M C Fouz, M Gonzalez Hernandez, O Gonzalez Lopez, S Goy Lopez, J M Hernandez, M I Josa, J Llorente Merino, C Martin Perez, E Martin Viscasillas, D Moran, C M Morcillo Perez, Á Navarro Tobar, R Paz Herrera, A Pérez-Calero Yzquierdo, J Puerta Pelayo, I Redondo, J Vazquez Escobar
Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

J F de Trocóniz
Universidad Autónoma de Madrid, Madrid, Spain

B Alvarez Gonzalez, J Ayllon Torresano, A Cardini, J Cuevas, J Del Riego Badas, D Estrada Acevedo, J Fernandez Menendez, S Folgueras, I Gonzalez Caballero, P Leguina, M Obeso Menendez, E Palencia Cortezon, J Prado Pico, A Soto Rodríguez, P Vischia
Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

S Blanco Fernández, I J Cabrillo, A Calderon, J Duarte Campderros, M Fernandez, G Gomez, C Lasaosa García, R Lopez Ruiz, C Martinez Rivero, P Martinez Ruiz del Arbol, F Matorras, P Matorras Cuevas, E Navarrete Ramos, J Piedra Gomez, C Quintana San Emeterio, L Scodellaro, I Vila, R Vilar Cortabitarte, J M Vizan Garcia
Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

B Kailasapathy[55], D D C Wickramarathna
University of Colombo, Colombo, Sri Lanka

W G D Dharmaratna[56], K Liyanage, N Perera
University of Ruhuna, Department of Physics, Matara, Sri Lanka

D Abbaneo, C Amendola, R Ardino, E Auffray, J Baechler, D Barney, J Bendavid, M Bianco, A Bocci, L Borgonovi, C Botta, A Bragagnolo, C E Brown, C Caillol, G Cerminara, P Connor, K Cormier, D d'Enterria, A Dabrowski, A David, A De Roeck, M M Defranchis, M Deile, M Dobson, P J Fernández Manteca, B A Fontana Santos Alves, E Fontanesi, W Funk, A Gaddi, S Giani, D Gigi, K Gill, F Glege, M Glowacki, A Gruber, J Hegeman, J K Heikkilä, R Hofsaess, B Huber, T James, P Janot, O Kaluzinska, O Karacheban[25], G Karathanasis, S Laurila, P Lecoq, E Leutgeb, C Lourenço, A-M Lyon, M Magherini, L Malgeri, M Mannelli, A Mehta, F Meijers, J A Merlin, S Mersi, E Meschi, M Migliorini, F Monti, F Moortgat, M Mulders, M Musich, I Neutelings, S Orfanelli, F Pantaleo, M Pari, G Petrucciani, A Pfeiffer, M Pierini, M Pitt, H Qu, D Rabady, A Reimers, B Ribeiro Lopes, F Riti, P Rosado, M Rovere, H Sakulin, R Salvatico, S Sanchez Cruz, S Scarfi, M Selvaggi, K Shchelina, P Silva, P Sphicas[57], A G Stahl Leiton, A Steen, S Summers, D Treille, P Tropea, E Vernazza, J Wanczyk[58], S Wuchterl, M Zarucki, P Zehetner, P Zejdl, G Zevi Della Porta
CERN, European Organization for Nuclear Research, Geneva, Switzerland

L Caminada[59], W Erdmann, R Horisberger, Q Ingram, H C Kaestli, D Kotlinski, C Lange, U Langenegger, A Nigamova, L Noehte[59], T Rohe, A Samalan
PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T K Aarrestad, M Backhaus, T Bevilacqua[59], G Bonomelli, C Cazzaniga, K Datta, P De Bryas Dexmiers D'Archiacchiac[58], A De Cosa, G Dissertori, M Dittmar, M Donegà, F Glessgen, C Grab, N Härringer, T G Harte, W Lustermann, M Malucchi, R A Manzoni, L Marchese, A Mascellani[58], F Nessi-Tedaldi, F Pauss, B Ristic, R Seidita, J Steggemann[58], A Tarabini, D Valsecchi, R Wallny
ETH Zurich—Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

C Amsler[60], P Bärtschi, F Bilandzija, M F Canelli, G Celotto, V Guglielmi, A Jofrehei, B Kilminster, T H Kwok, S Leontsinis, V Lukashenko, A Macchiolo, F Meng, M Missiroli, J Motta, P Robmann, E Shokr, F Stäger, R Tramontano, P Viscone
Universität Zürich, Zurich, Switzerland

D Bhowmik, C M Kuo, P K Rout, S Taj, P C Tiwari[36]
National Central University, Chung-Li, Taiwan

L Ceard, K F Chen, Z G Chen, A De Iorio, W-S Hou, T H Hsu, Y W Kao, S Karmakar, G Kole, Y Y Li, R-S Lu, E Paganis, X F Su, J Thomas-Wilsker, L S Tsai, D Tsionou, H Y Wu, E Yazgan
National Taiwan University (NTU), Taipei, Taiwan

C Asawatangtrakuldee, N Srimanobhas
High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

Y Maghrbi
Tunis El Manar University, Tunis, Tunisia

D Agyel, F Dolek, I Dumanoglu[61], Y Guler[62], E Gurpinar Guler[62], C Isik, O Kara[63], A Kayis Topaksu, Y Komurcu, G Onengut, K Ozdemir[64], B Tali[65], U G Tok, E Uslan, I S Zorbakir
Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

S Sen
Hacettepe University, Ankara, Turkey

M Yalvac[66]
Middle East Technical University, Physics Department, Ankara, Turkey

B Akgun, I O Atakisi[67], E Gülmez, M Kaya[68], O Kaya[69], M A Sarkisla[70], S Tekten[71]
Bogazici University, Istanbul, Turkey

D Boncukcu, A Cakir, K Cankocak[61,72]
Istanbul Technical University, Istanbul, Turkey

B Hacisahinoglu, I Hos[73], B Kaynak, S Ozkorucuklu, O Potok, H Sert, C Simsek, C Zorbilmez
Istanbul University, Istanbul, Turkey

S Cerci, C Dozen[74], B Isildak, E Simsek, D Sunar Cerci, T Yetkin[74]
Yildiz Technical University, Istanbul, Turkey

A Boyaryntsev, O Dadazhanova, B Grynyov
Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

L Levchuk
National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

J J Brooke, A Bundock, F Bury, E Clement, D Cussans, D Dharmender, H Flacher, J Goldstein, H F Heath, M-L Holmberg, L Kreczko, S Paramesvaran, L Robertshaw, M S Sanjrani[39], J Segal, V J Smith
University of Bristol, Bristol, United Kingdom

A H Ball, K W Bell, A Belyaev[75], C Brew, R M Brown, D J A Cockerill, A Elliot, K V Ellis, J Gajownik, K Harder, S Harper, J Linacre, K Manolopoulos, M Moallemi, D M Newbold, E Olaiya, D Petyt, T Reis, A R Sahasransu, G Salvi, T Schuh, C H Shepherd-Themistocleous, I R Tomalin, K C Whalen, T Williams
Rutherford Appleton Laboratory, Didcot, United Kingdom

I Andreou, R Bainbridge, P Bloch, O Buchmuller, C A Carrillo Montoya, D Colling, I Das, P Dauncey, G Davies, M Della Negra, S Fayer, G Fedi, G Hall, H R Hoorani, A Howard, G Iles, C R Knight, P Krueper, J Langford, K H Law, J León Holgado, L Lyons, A-M Magnan, B Maier, S Mallios, A Mastronikolis, M Mieskolainen, J Nash[76], M Pesaresi, P B Pradeep, B C Radburn-Smith, A Richards, A Rose, L Russell, K Savva, R Schmitz, C Seez, R Shukla, A Tapper, K Uchida, G P Uttley, T Virdee[27], M Vojinovic, N Wardle, D Winterbottom
Imperial College, London, United Kingdom

J E Cole, A Khan, P Kyberd, I D Reid
Brunel University, Uxbridge, United Kingdom

S Abdullin, A Brinkerhoff, E Collins, M R Darwish, J Dittmann, K Hatakeyama, V Hegde, J Hiltbrand, B McMaster, J Samudio, S Sawant, C Sutantawibul, J Wilson
Baylor University, Waco, TX, United States of America

J M Hogan
Bethel University, St. Paul, MN, United States of America

R Bartek, A Dominguez, S Raj, B Sahu, A E Simsek, S S Yu
Catholic University of America, Washington, DC, United States of America

B Bam, A Buchot Perraguin, S Campbell, R Chudasama, S I Cooper, C Crovella, G Fidalgo, S V Gleyzer, A Khukhunaishvili, K Matchev, E Pearson, P Rumerio[77], E Usai, R Yi
The University of Alabama, Tuscaloosa, AL, United States of America

S Cholak, G De Castro, Z Demiragli, C Erice, C Fangmeier, C Fernandez Madrazo, J Fulcher, F Golf, S Jeon, J O'Cain, I Reed, J Rohlf, K Salyer, D Sperka, D Spitzbart, I Suarez, A Tsatsos, E Wurtz, A G Zecchinelli
Boston University, Boston, MA, United States of America

G Barone, G Benelli, D Cutts, S Ellis, L Gouskos, M Hadley, U Heintz, K W Ho, T Kwon, L Lambrecht, G Landsberg, K T Lau, J Luo, S Mondal, J Roloff, T Russell, S Sagir[78], X Shen, M Stamenkovic, N Venkatasubramanian
Brown University, Providence, RI, United States of America

S Abbott, S Baradia, B Barton, R Breedon, H Cai, M Calderon De La Barca Sanchez, E Cannaert, M Chertok, M Citron, J Conway, P T Cox, F Eble, R Erbacher, O Kukral, G Mocellin, S Ostrom, I Salazar Segovia, J S Tafoya Vargas, W Wei, S Yoo
University of California, Davis, Davis, CA, United States of America

K Adamidis, M Bachtis, D Campos, R Cousins, S Crossley, G Flores Avila, J Hauser, M Ignatenko, M A Iqbal, T Lam, Y F Lo, E Manca, A Nunez Del Prado, D Saltzberg, V Valuev
University of California, Los Angeles, CA, United States of America

R Clare, J W Gary, G Hanson
University of California, Riverside, Riverside, CA, United States of America

A Aportela, A Arora, J G Branson, S Cittolin, S Cooperstein, B D'Anzi, D Diaz, J Duarte, L Giannini, Y Gu, J Guiang, V Krutelyov, R Lee, J Letts, H Li, M Masciovecchio, F Mokhtar, S Mukherjee, M Pieri, D Primosch, M Quinnan, V Sharma, M Tadel, E Vourliotis, F Würthwein, A Yagil, Z Zhao
University of California, San Diego, La Jolla, CA, United States of America

A Barzdukas, L Brennan, C Campagnari, S Carron Montero[79], K Downham, C Grieco, M M Hussain, J Incandela, M W K Lai, A J Li, P Masterson, J Richman, S N Santpur, D Stuart, T Á Vámi, X Yan, D Zhang
University of California, Santa Barbara—Department of Physics, Santa Barbara, CA, United States of America

A Albert, S Bhattacharya, A Bornheim, O Cerri, R Kansal, H B Newman, G Reales Gutiérrez, T Sievert, M Spiropulu, J R Vlimant, R A Wynne, S Xie
California Institute of Technology, Pasadena, CA, United States of America

J Alison, S An, M Cremonesi, V Dutta, E Y Ertorer, T Ferguson, T A Gómez Espinosa, A Harilal, A Kallil Tharayil, M Kanemura, C Liu, M Marchegiani, P Meiring, S Murthy, P Palit, K Park, M Paulini, A Roberts, A Sanchez, W Terrill
Carnegie Mellon University, Pittsburgh, PA, United States of America

J P Cumalat, W T Ford, A Hart, S Kwan, J Pearkes, C Savard, N Schonbeck, K Stenson, K A Ulmer, S R Wagner, N Zipper, D Zuolo
University of Colorado Boulder, Boulder, CO, United States of America

J Alexander, X Chen, J Dickinson, A Duquette, J Fan, X Fan, J Grassi, S Hogan, P Kotamnives, J Monroy, G Niendorf, M Oshiro, J R Patterson, A Ryd, J Thom, P Wittich, R Zou, L Zygala
Cornell University, Ithaca, NY, United States of America

M Albrow, M Alyari, O Amram, G Apollinari, A Apresyan, L A T Bauerdick, D Berry, J Berryhill, P C Bhat, K Burkett, J N Butler, A Canepa, G B Cerati, H W K Cheung, F Chlebana, C Cosby, G Cummings, I Dutta, V D Elvira, J Freeman, A Gandrakota, Z Gecse, L Gray, D Green, A Grummer, S Grünendahl, D Guerrero, O Gutsche, R M Harris, J Hirschauer, V Innocente, B Jayatilaka, S Jindariani, M Johnson, U Joshi, B Klima, S Lammel, C Lee, D Lincoln, R Lipton, T Liu, K Maeshima, D Mason, P McBride, P Merkel, S Mrenna, S Nahn, J Ngadiuba, D Noonan, S Norberg, V Papadimitriou, N Pastika, K Pedro, C Pena[80], C E Perez Lara, V Perovic, F Ravera, A Reinsvold Hall[81], L Ristori, M Safdari, E Sexton-Kennedy, E Smith, N Smith, A Soha, L Spiegel, S Stoynev, J Strait, L Taylor, S Tkaczyk, N V Tran, L Uplegger, E W Vaandering, C Wang, I Zoi
Fermi National Accelerator Laboratory, Batavia, IL, United States of America

C Aruta, P Avery, D Bourilkov, P Chang, V Cherepanov, R D Field, C Huh, E Koenig, M Kolosova, J Konigsberg, A Korytov, G Mitselmakher, K Mohrman, A Muthirakalayil Madhu, N Rawal, S Rosenzweig, V Sulimov, Y Takahashi, J Wang
University of Florida, Gainesville, FL, United States of America

T Adams, A Al Kadhim, A Askew, S Bower, R Goff, R Hashmi, A Hassani, R S Kim, T Kolberg, G Martinez, M Mazza, H Prosper, P R Prova, R Yohay
Florida State University, Tallahassee, FL, United States of America

B Alsufyani, S Butalla, S Das, M Hohlmann, M Lavinsky, E Yanes
Florida Institute of Technology, Melbourne, FL, United States of America

M R Adams, N Barnett, A Baty, C Bennett, R Cavanaugh, R Escobar Franco, O Evdokimov, C E Gerber, H Gupta, M Hawksworth, A Hingrajiya, D J Hofman, Z Huang, J H Lee, C Mills, S Nanda, G Nigmatkulov, B Ozek, T Phan, D Pilipovic, R Pradhan, E Prifti, P Roy, T Roy, D Shekar, N Singh, A Thielen, M B Tonjes, N Varelas, M A Wadud, J Yoo
University of Illinois Chicago, Chicago, IL, United States of America

M Alhusseini, D Blend, K Dilsiz[82], O K Köseyan, A Mestvirishvili[83], O Neogi, H Ogul[84], Y Onel, A Penzo, C Snyder, E Tiras[85]
The University of Iowa, Iowa City, IA, United States of America

B Blumenfeld, J Davis, A V Gritsan, L Kang, S Kyriacou, P Maksimovic, M Roguljic, S Sekhar, M V Srivastav, M Swartz
Johns Hopkins University, Baltimore, MD, United States of America

A Abreu, L F Alcerro Alcerro, J Anguiano, S Arteaga Escatel, P Baringer, A Bean, R Bhattacharya, Z Flowers, D Grove, J King, G Krintiras, M Lazarovits, C Le Mahieu, J Marquez, M Murray, M Nickel, S Popescu[86], C Rogan, C Royon, S Rudrabhatla, S Sanders, C Smith, G Wilson
The University of Kansas, Lawrence, KS, United States of America

B Allmond, N Islam, A Ivanov, K Kaadze, Y Maravin, J Natoli, G G Reddy, D Roy, G Sorrentino
Kansas State University, Manhattan, KS, United States of America

A Baden, A Belloni, J Bistany-riebman, S C Eno, N J Hadley, S Jabeen, R G Kellogg, T Koeth, B Kronheim, S Lascio, P Major, A C Mignerey, C Palmer, C Papageorgakis, M M Paranjpe, E Popova[87], A Shevelev, L Zhang
University of Maryland, College Park, MD, United States of America

C Baldenegro Barrera, H Bossi, S Bright-Thonney, I A Cali, Y C Chen, P C Chou, M D'Alfonso, J Eysermans, C Freer, G Gomez-Ceballos, M Goncharov, G Grosso, P Harris, D Hoang, G M Innocenti, K Ivanov, G Kopp, D Kovalskyi, J Krupa, L Lavezzo, Y-J Lee, K Long, C Mcginn, A Novak, M I Park, C Paus, C Reissel, C Roland, G Roland, S Rothman, T A Sheng, G S F Stephans, P Van Steenweghen, D Walter, J Wang, Z Wang, B Wyslouch, T J Yang
Massachusetts Institute of Technology, Cambridge, MA, United States of America

A Alpana, B Crossman, W J Jackson, C Kapsiak, M Krohn, D Mahon, J Mans, B Marzocchi, R Rusack, O Sancar, R Saradhy, N Strobbe
University of Minnesota, Minneapolis, MN, United States of America

K Bloom, D R Claes, G Haza, J Hossain, C Joo, I Kravchenko, K H M Kwok, A Rohilla, J E Siado, W Tabb, A Vagnerini, A Wightman, F Yan
University of Nebraska-Lincoln, Lincoln, NE, United States of America

H Bandyopadhyay, L Hay, H W Hsia, I Iashvili, A Kalogeropoulos, A Kharchilava, A Mandal, M Morris, D Nguyen, S Rappoccio, H Rejeb Sfar, A Williams, D Yu
State University of New York at Buffalo, Buffalo, NY, United States of America

A Aarif, G Alverson, E Barberis, J Bonilla, B Bylsma, M Campana, J Dervan, Y Haddad, Y Han, I Israr, A Krishna, M Lu, N Manganelli, R Mccarthy, D M Morse, T Orimoto, L Skinnari, C S Thoreson, E Tsai, D Wood
Northeastern University, Boston, MA, United States of America

S Dittmer, K A Hahn, M Mcginnis, Y Miao, D G Monk, M H Schmitt, A Taliercio, M Velasco, J Wang
Northwestern University, Evanston, IL, United States of America

G Agarwal, R Band, R Bucci, S Castells, A Das, A Datta, A Ehnis, R Goldouzian, M Hildreth, K Hurtado Anampa, T Ivanov, C Jessop, A Karneyeu, K Lannon, J Lawrence, N Loukas, L Lutton, J Mariano, N Marinelli, T McCauley, C Mcgrady, C Moore, Y Musienko[21], H Nelson, M Osherson, A Piccinelli, R Ruchti, A Townsend, Y Wan, M Wayne, H Yockey
University of Notre Dame, Notre Dame, IN, United States of America

M Carrigan, R De Los Santos, L S Durkin, C Hill, M Joyce, D A Wenzl, B L Winer, B R Yates
The Ohio State University, Columbus, OH, United States of America

H Bouchamaoui, G Dezoort, P Elmer, A Frankenthal, M Galli, B Greenberg, N Haubrich, K Kennedy, Y Lai, D Lange, A Loeliger, D Marlow, I Ojalvo, J Olsen, F Simpson, D Stickland, C Tully
Princeton University, Princeton, NJ, United States of America

S Malik, R Sharma
University of Puerto Rico, Mayaguez, PR, United States of America

S Chandra, A Gu, L Gutay, M Huwiler, M Jones, A W Jung, D Kondratyev, J Li, M Liu, G Negro, N Neumeister, G Paspalaki, S Piperov, N R Saha, J F Schulte, F Wang, A Wildridge, W Xie, Y Yao, Y Zhong
Purdue University, West Lafayette, IN, United States of America

N Parashar, A Pathak, E Shumka
Purdue University Northwest, Hammond, IN, United States of America

D Acosta, A Agrawal, C Arbour, T Carnahan, P Das, K M Ecklund, F J M Geurts, T Huang, I Krommydas, N Lewis, W Li, J Lin, O Miguel Colin, B P Padley, R Redjimi, J Rotter, C Vico Villalba, M Wulansatiti, E Yigitbasi, Y Zhang
Rice University, Houston, TX, United States of America

O Bessidskaia Bylund, A Bodek, P de Barbaro[†], R Demina, A Garcia-Bellido, H S Hare, O Hindrichs, N Parmar, P Parygin[87], H Seo, R Taus, Y H Yu
University of Rochester, Rochester, NY, United States of America

B Chiarito, J P Chou, S V Clark, S Donnelly, D Gadkari, Y Gershtein, E Halkiadakis, C Houghton, D Jaroslawski, A Kobert, I Laflotte, A Lath, J Martins, M Perez Prada, B Rand, J Reichert, P Saha, S Salur, S Somalwar, R Stone, S A Thayil, S Thomas, J Vora
Rutgers, The State University of New Jersey, Piscataway, NJ, United States of America

D Ally, A G Delannoy, S Fiorendi, J Harris, T Holmes, A R Kanuganti, N Karunarathna, J Lawless, L Lee, E Nibigira, B Skipworth, S Spanier
University of Tennessee, Knoxville, TN, United States of America

D Aebi, M Ahmad, T Akhter, K Androsov, A Basnet, A Bolshov, O Bouhali[88], A Cagnotta, V D'Amante, R Eusebi, P Flanagan, J Gilmore, Y Guo, T Kamon, S Luo, R Mueller, A Safonov
Texas A&M University, College Station, TX, United States of America

N Akchurin, J Damgov, Y Feng, N Gogate, W Jin, S W Lee, C Madrid, A Mankel, T Peltola, I Volobouev
Texas Tech University, Lubbock, TX, United States of America

E Appelt, Y Chen, S Greene, A Gurrola, W Johns, R Kunnawalkam Elayavalli, A Melo, D Rathjens, F Romeo, P Sheldon, S Tuo, J Velkovska, J Viinikainen, J Zhang
Vanderbilt University, Nashville, TN, United States of America

B Cardwell, H Chung, B Cox, J Hakala, G Hamilton Ilha Machado, R Hirosky, M Jose, A Ledovskoy, C Mantilla, C Neu, C Ramón Álvarez, Z Wu
University of Virginia, Charlottesville, VA, United States of America

S Bhattacharya, P E Karchin
Wayne State University, Detroit, MI, United States of America

A Aravind, S Banerjee, K Black, T Bose, E Chavez, S Dasu, P Everaerts, C Galloni, H He, M Herndon, A Herve, C K Koraka, S Lomte, R Loveless, A Mallampalli, A Mohammadi, S Mondal, T Nelson, G Parida, L Pétré, D Pinna, A Savin, V Shang, V Sharma, W H Smith, D Teague, A Warden
University of Wisconsin—Madison, Madison, WI, United States of America

S Afanasiev, V Alexakhin, Yu Andreev, T Aushev, D Budkouski, R Chistov, M Danilov, T Dimova, A Ershov, S Gninenko, I Gorbunov, A Kamenev, V Karjavine, M Kirsanov, V Klyukhin, O Kodolova[89], V Korenkov, I Korsakov, A Kozyrev, N Krasnikov, A Lanev, A Malakhov, V Matveev, A Nikitenko[90,89], V Palichik, V Perelygin, S Petrushanko, O Radchenko, M Savina, V Shalaev, S Shmatov, S Shulha, Y Skovpen, K Slizhevskiy, V Smirnov, O Teryaev, I Tlisova, A Toropin, N Voytishin, A Zarubin, I Zhizhin
Authors affiliated with an international laboratory covered by a cooperation agreement with CERN

E Boos, V Bunichev, M Dubinin[80], A Gribushin, V Savrin, A Snigirev, L Dudko, V Kim[21], V Murzin, V Oreshkin, D Sosnov
Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

[†]Deceased

[1]Also at Yerevan State University, Yerevan, Armenia
[2]Also at TU Wien, Vienna, Austria
[3]Also at Ghent University, Ghent, Belgium
[4]Also at FACAMP—Faculdades de Campinas, Sao Paulo, Brazil
[5]Also at Universidade Estadual de Campinas, Campinas, Brazil
[6]Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil
[7]Also at The University of the State of Amazonas, Manaus, Brazil
[8]Also at University of Chinese Academy of Sciences, Beijing, People's Republic of China
[9]Also at University of Chinese Academy of Sciences, Beijing, People's Republic of China
[10]Also at School of Physics, Zhengzhou University, Zhengzhou, People's Republic of China
[11]Now at Henan Normal University, Xinxiang, People's Republic of China
[12]Also at University of Shanghai for Science and Technology, Shanghai, People's Republic of China
[13]Also at The University of Iowa, Iowa City, IA, United States of America
[14]Also at Nanjing Normal University, Nanjing, People's Republic of China
[15]Also at Center for High Energy Physics, Peking University, Beijing, People's Republic of China
[16]Also at Helwan University, Cairo, Egypt
[17]Now at Zewail City of Science and Technology, Zewail, Egypt
[18]Also at British University in Egypt, Cairo, Egypt
[19]Also at Purdue University, West Lafayette, IN, United States of America
[20]Also at Université de Haute Alsace, Mulhouse, France
[21]Also at an institute formerly covered by a cooperation agreement with CERN
[22]Also at University of Hamburg, Hamburg, Germany
[23]Also at RWTH Aachen University, IIIdot Physikalisches Institut A, Aachen, Germany
[24]Also at Bergische University Wuppertal (BUW), Wuppertal, Germany
[25]Also at Brandenburg University of Technology, Cottbus, Germany
[26]Also at Forschungszentrum Jülich, Juelich, Germany
[27]Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
[28]Also at HUN-REN ATOMKI—Institute of Nuclear Research, Debrecen, Hungary
[29]Now at Universitatea Babes-Bolyai—Facultatea de Fizica, Cluj-Napoca, Romania
[30]Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary
[31]Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
[32]Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
[33]Also at The University of Kansas, Lawrence, KS, United States of America
[34]Also at Punjab Agricultural University, Ludhiana, India
[35]Also at University of Hyderabad, Hyderabad, India
[36]Also at Indian Institute of Science (IISc), Bangalore, India
[37]Also at University of Visva-Bharati, Santiniketan, India
[38]Also at Institute of Physics, Bhubaneswar, India
[39]Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
[40]Also at Isfahan University of Technology, Isfahan, Iran
[41]Also at Sharif University of Technology, Tehran, Iran
[42]Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
[43]Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
[44]Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
[45]Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
[46]Also at James Madison University, Harrisonburg, MD, United States of America
[47]Also at Università degli Studi Guglielmo Marconi, Roma, Italy
[48]Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
[49]Also at Fermi National Accelerator Laboratory, Batavia, IL, United States of America
[50]Also at Lulea University of Technology, Lulea, Sweden
[51]Also at Consiglio Nazionale delle Ricerche—Istituto Officina dei Materiali, Perugia, Italy
[52]Also at UPES—University of Petroleum and Energy Studies, Dehradun, India
[53]Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
[54]Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
[55]Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
[56]Also at Saegis Campus, Nugegoda, Sri Lanka

[57]Also at National and Kapodistrian University of Athens, Athens, Greece
[58]Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
[59]Also at Universität Zürich, Zurich, Switzerland
[60]Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
[61]Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
[62]Also at Konya Technical University, Konya, Turkey
[63]Also at Istanbul Topkapi University, Istanbul, Turkey
[64]Also at Izmir Bakircay University, Izmir, Turkey
[65]Also at Adiyaman University, Adiyaman, Turkey
[66]Also at Bozok Universitetesi Rektörlügü, Yozgat, Turkey
[67]Also at Istanbul Sabahattin Zaim University, Istanbul, Turkey
[68]Also at Marmara University, Istanbul, Turkey
[69]Also at Milli Savunma University, Istanbul, Turkey
[70]Also at Informatics and Information Security Research Center, Gebze/Kocaeli, Turkey
[71]Also at Kafkas University, Kars, Turkey
[72]Now at Istanbul Okan University, Istanbul, Turkey
[73]Also at Istanbul University—Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
[74]Also at Istinye University, Istanbul, Turkey
[75]Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
[76]Also at Monash University, Faculty of Science, Clayton, Australia
[77]Also at Università di Torino, Torino, Italy
[78]Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
[79]Also at California Lutheran University, Thousand Oaks, CA, United States of America
[80]Also at California Institute of Technology, Pasadena, CA, United States of America
[81]Also at United States Naval Academy, Annapolis, MD, United States of America
[82]Also at Bingol University, Bingol, Turkey
[83]Also at Georgian Technical University, Tbilisi, Georgia
[84]Also at Sinop University, Sinop, Turkey
[85]Also at Erciyes University, Kayseri, Turkey
[86]Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
[87]Now at another institute formerly covered by a cooperation agreement with CERN
[88]Also at Hamad Bin Khalifa University (HBKU), Doha, Qatar
[89]Also at Yerevan Physics Institute, Yerevan, Armenia
[90]Also at Imperial College, London, United Kingdom

References

[1] ATLAS Collaboration 2008 The ATLAS experiment at the CERN large Hadron collider *J. Instrum.* **3** S08003
[2] CMS Collaboration 2008 The CMS experiment at the CERN LHC *J. Instrum.* **3** S08004
[3] ATLAS Collaboration 2012 Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC *Phys. Lett. B* **716** 1
[4] CMS Collaboration 2012 Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC *Phys. Lett. B* **716** 30
[5] CMS Collaboration 2013 Observation of a new boson with mass near 125 GeV in pp collisions at $\sqrt{s} = 7$ and 8 TeV *J. High Energy Phys.* JHEP06(2013)081
[6] Englert F and Brout R 1964 Broken symmetry and the mass of gauge vector mesons *Phys. Rev. Lett.* **13** 321
[7] Higgs P W 1964 Broken symmetries, massless particles and gauge fields *Phys. Lett.* **12** 132
[8] Higgs P W 1964 Broken symmetries and the masses of gauge bosons *Phys. Rev. Lett.* **13** 508
[9] Guralnik G S, Hagen C R and Kibble T W B 1964 Global conservation laws and massless particles *Phys. Rev. Lett.* **13** 585
[10] Higgs P W 1966 Spontaneous symmetry breakdown without massless bosons *Phys. Rev.* **145** 1156
[11] Kibble T W B 1967 Symmetry breaking in non-abelian gauge theories *Phys. Rev.* **155** 1554
[12] Evans L and Bryant P 2008 LHC machine *J. Instrum.* **3** S08001
[13] LHC Higgs Cross section Working Group 2016 *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*
[14] Anastasiou C *et al* 2016 High precision determination of the gluon fusion Higgs boson cross-section at the LHC *J. High Energy Phys.* JHEP05(2016)058
[15] Cacciari M *et al* 2015 Fully differential Vector-boson-fusion Higgs production at next-to-next-to-leading order *Phys. Rev. Lett.* **115** 082002
Cacciari M *et al* 2018 Fully differential Vector-boson-fusion Higgs production at next-to-next-to-leading order *Phys. Rev. Lett.* **120** 139901 (erratum)
[16] Brein O, Djouadi A and Harlander R 2004 NNLO QCD corrections to the Higgs-strahlung processes at hadron colliders *Phys. Lett. B* **579** 149

[17] Brein O, Harlander R V and Zirke T J E 2013 vh@nnlo—Higgs strahlung at hadron colliders *Comput. Phys. Commun.* **184** 998
[18] Harlander R V, Liebler S and Zirke T 2014 Higgs strahlung at the Large Hadron Collider in the 2-Higgs-doublet model *J. High Energy Phys.* JHEP02(2014)023
[19] Altenkamp L *et al* 2013 Gluon-induced Higgs-strahlung at next-to-leading order QCD *J. High Energy Phys.* JHEP02(2013)078
[20] Frixione S *et al* 2015 Electroweak and QCD corrections to top-pair hadroproduction in association with heavy bosons *J. High Energy Phys.* JHEP06(2015)184
[21] Demartin F, Maltoni F, Mawatari K and Zaro M 2015 Higgs production in association with a single top quark at the LHC *Eur. Phys. J. C* **75** 267
[22] Berger N et al 2019 Simplified template cross sections – Stage 1.1 and 1.2 (arXiv:1906.02754)
[23] LHC Higgs Cross section Working Group 2013 Handbook of LHC Higgs cross sections: 3. Higgs properties (arXiv:1307.1347)
[24] Brivio I and Trott M 2019 The standard model as an effective field theory *Phys. Rept.* **793** 1
[25] ATLAS Collaboration 2016 Measurements of the Higgs boson production and decay rates and coupling strengths using pp collision data at $\sqrt{s} = 7$ and 8 TeV in the ATLAS experiment *Eur. Phys. J. C* **76** 6
[26] CMS Collaboration 2015 Precise determination of the mass of the Higgs boson and tests of compatibility of its couplings with the standard model predictions using proton collisions at 7 and 8 TeV *Eur. Phys. J. C* **75** 212
[27] ATLAS and CMS Collaborations 2016 Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV *J. High Energy Phys.* JHEP08(2016)045
[28] ATLAS Collaboration 2020 Combined measurements of Higgs boson production and decay using up to 80 fb^{-1} of proton-proton collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS experiment *Phys. Rev. D* **101** 012002
[29] ATLAS Collaboration 2022 A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery *Nature* **607** 52
ATLAS Collaboration 2023 A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery *Nature* **623** E5 (erratum)
[30] ATLAS Collaboration 2024 Interpretations of the ATLAS measurements of Higgs boson production and decay rates and differential cross-sections in pp collisions at $\sqrt{s} = 13$ TeV *J. High Energy Phys.* JHEP11(2024)097
[31] CMS Collaboration 2019 Combined measurements of Higgs boson couplings in proton-proton collisions at $\sqrt{s} = 13$ TeV *Eur. Phys. J. C* **79** 421
[32] CMS Collaboration 2022 A portrait of the Higgs boson by the CMS experiment ten years after the discovery *Nature* **607** 60
CMS Collaboration 2023 A portrait of the Higgs boson by the CMS experiment ten years after the discovery *Nature* **18** E4 (corrigendum)
[33] CMS Collaboration 2024 Measurement of the Higgs boson production via vector boson fusion and its decay into bottom quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV *J. High Energy Phys.* JHEP01(2024)173
[34] CMS Collaboration 2024 Measurement of boosted Higgs bosons produced via vector boson fusion or gluon fusion in the $\text{H} \to \text{b}\bar{\text{b}}$ decay mode using LHC proton-proton collision data at $\sqrt{s} = 13$ TeV *J. High Energy Phys.* JHEP12(2024)035
[35] CMS Collaboration 2025 Measurement of the Higgs boson mass and width using the four-lepton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV *Phys. Rev. D* **111** 092014
[36] CMS Collaboration 2023 A search for decays of the Higgs boson to invisible particles in events with a top-antitop quark pair or a vector boson in proton-proton collisions at $\sqrt{s} = 13$ TeV *Eur. Phys. J. C* **83** 933
[37] CMS Collaboration 2024 Measurement of simplified template cross sections of the Higgs boson produced in association with W or Z bosons in the $\text{H} \to \text{b}\bar{\text{b}}$ decay channel in proton-proton collisions at $\sqrt{s} = 13$ TeV *Phys. Rev. D* **109** 092011
[38] CMS Collaboration 2025 Measurement of the $\text{t}\bar{\text{t}}\text{H}$ and tH production rates in the $\text{H} \to \text{b}\bar{\text{b}}$ decay channel using proton-proton collision data at $\sqrt{s} = 13$ TeV *J. High Energy Phys.* JHEP02(2025)097
[39] CMS Collaboration 2020 A measurement of the Higgs boson mass in the diphoton decay channel *Phys. Lett. B* **805** 135425
[40] ATLAS Collaboration 2023 Combined measurement of the Higgs boson mass from the $\text{H} \to \gamma\gamma$ and $\text{H} \to ZZ^* \to 4\ell$ decay channels with the ATLAS detector using $\sqrt{s} = 7$, 8 and 13 TeV pp collision data *Phys. Rev. Lett.* **131** 251802
[41] 2026 HEPData record for this analysis (https://doi.org/10.17182/hepdata.168509)
[42] CMS Collaboration 2024 Development of the CMS detector for the CERN LHC Run 3 *J. Instrum.* **19** 05064
[43] CMS Collaboration 2020 Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV *J. Instrum.* **15** 10017
[44] CMS Collaboration 2017 The CMS trigger system *J. Instrum.* **12** 01020
[45] CMS Collaboration 2024 Performance of the CMS high-level trigger during LHC Run 2 *J. Instrum.* **19** 11021
[46] CMS Collaboration 2021 Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC *J. Instrum.* **16** 05014
[47] CMS Collaboration 2018 Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV *J. Instrum.* **13** 06015
[48] CMS Collaboration 2014 Description and performance of track and primary-vertex reconstruction with the CMS tracker *J. Instrum.* **9** 10009
[49] CMS Collaboration 2021 Measurements of Higgs boson production cross sections and couplings in the diphoton decay channel at $\sqrt{s} = 13$ TeV *J. High Energy Phys.* JHEP07(2021)027
[50] CMS Collaboration 2021 Measurements of production cross sections of the Higgs boson in the four-lepton final state in proton–proton collisions at $\sqrt{s} = 13$ TeV *Eur. Phys. J. C* **81** 488
[51] CMS Collaboration 2023 Measurements of the Higgs boson production cross section and couplings in the W boson pair decay channel in proton-proton collisions at $\sqrt{s} = 13$ TeV *Eur. Phys. J. C* **83** 667
[52] CMS Collaboration 2023 Measurements of Higgs boson production in the decay channel with a pair of τ leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV *Eur. Phys. J. C* **83** 562
[53] CMS Collaboration 2021 Measurement of the Higgs boson production rate in association with top quarks in final states with electrons, muons and hadronically decaying tau leptons at $\sqrt{s} = 13$ TeV *Eur. Phys. J. C* **81** 378
[54] CMS Collaboration 2021 Evidence for Higgs boson decay to a pair of muons *J. High Energy Phys.* JHEP01(2021)148
[55] CMS Collaboration 2023 Search for Higgs boson decays to a Z boson and a photon in proton-proton collisions at $\sqrt{s} = 13$ TeV *J. High Energy Phys.* JHEP05(2023)233
[56] CMS Collaboration 2021 Search for new particles in events with energetic jets and large missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV *J. High Energy Phys.* JHEP11(2021)153

[57] CMS Collaboration 2022 Search for invisible decays of the Higgs boson produced via vector boson fusion in proton-proton collisions at $\sqrt{s} = 13$ TeV *Phys. Rev. D* **105** 092007

[58] CMS Collaboration 2021 Search for dark matter produced in association with a leptonically decaying Z boson in proton-proton collisions at $\sqrt{s} = 13$ TeV *Eur. Phys. J. C* **81** 13
CMS Collaboration 2020 Search for dark matter produced in association with a leptonically decaying Z boson in proton-proton collisions at $\sqrt{s} = 13$ TeV *Eur. Phys. J. C* **81** 333 (erratum)

[59] CMS Collaboration 2024 Performance of the CMS electromagnetic calorimeter in pp collisions at $\sqrt{s} = 13$ TeV *J. Instrum.* **19** 09004

[60] CMS Collaboration 2022 Identification of hadronic tau lepton decays using a deep neural network *J. Instrum.* **17** 07023

[61] CMS Collaboration 2017 Particle-flow reconstruction and global event description with the CMS detector *J. Instrum.* **12** 10003

[62] CMS Collaboration 2025 Performance of heavy-flavour jet identification in Lorentz-boosted topologies in proton-proton collisions at $\sqrt{s} = 13$ TeV *J. Instrum.* **20** 11006

[63] CMS Collaboration 2023 Search for Higgs boson decay to a charm quark-antiquark pair in proton-proton collisions at $\sqrt{s} = 13$ TeV *Phys. Rev. Lett.* **131** 061801

[64] CMS Collaboration 2023 Search for Higgs boson and observation of Z boson through their decay into a charm quark-antiquark pair in boosted topologies in proton-proton collisions at $\sqrt{s} = 13$ TeV *Phys. Rev. Lett.* **131** 041801

[65] CMS Collaboration 2026 Simultaneous probe of the charm and bottom quark Yukawa couplings using $\mathrm{t\bar{t}H}$ events *Phys. Rev. Lett.* **136** 011801

[66] Alwall J *et al* 2014 The automated computation of tree-level and next-to-leading order differential cross sections and their matching to parton shower simulations *J. High Energy Phys.* JHEP07(2014)079

[67] Hamilton K, Nason P, Re E and Zanderighi G 2013 NNLOPS simulation of Higgs boson production *J. High Energy Phys.* JHEP10(2013)222

[68] Hamilton K, Nason P and Zanderighi G 2012 MINLO: multi-scale improved NLO *J. High Energy Phys.* JHEP10(2012)155

[69] Luisoni G, Nason P, Oleari C and Tramontano F 2013 $HW^{\pm}/HZ + 0$ and 1 jet at NLO with the POWHEG BOX interfaced to GoSam and their merging within MiNLO *J. High Energy Phys.* JHEP10(2013)083

[70] Sjöstrand T *et al* 2015 An introduction to PYTHIA 8.2 *Comput. Phys. Commun.* **191** 159

[71] CMS Collaboration 2020 Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements *Eur. Phys. J. C* **80** 4

[72] Collaboration N N P D F 2017 Parton distributions from high-precision collider data *Eur. Phys. J. C* **77** 663

[73] Becker K *et al* 2024 Precise predictions for boosted Higgs production *SciPost Phys. Core* **7** 001

[74] ATLAS and CMS Collaborations and LHC Higgs Combination Group 2011 Procedure for the LHC Higgs boson search combination in Summer 2011 *Technical Report* CMS-NOTE-2011-005, ATL-PHYS-PUB-2011-11

[75] CMS Collaboration 2012 Combined results of searches for the standard model Higgs boson in pp collisions at $\sqrt{s} = 7$ TeV *Phys. Lett. B* **710** 26

[76] CMS Collaboration 2024 The CMS statistical analysis and combination tool: combine *Comput. Softw. Big Sci.* **8** 19

[77] Verkerke W and Kirkby D P 2003 The RooFit toolkit for data modeling *eConf* **C0303241** MOLT007

[78] Moneta L *et al* 2010 The RooStats project *Proc. Sci.* **ACAT2010** 057

[79] Bernlochner F U, Fry D C, Menary S B and Persson E 2023 Cover your bases: asymptotic distributions of the profile likelihood ratio when constraining effective field theories in high-energy physics *SciPost Phys. Core* **6** 013

[80] Cowan G, Cranmer K, Gross E and Vitells O 2011 Asymptotic formulae for likelihood-based tests of new physics *Eur. Phys. J. C* **71** 1554
(https://doi.org/10.1140/epjc/s10052-013-2501-z) (erratum)

[81] CMS Collaboration 2021 Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS *Eur. Phys. J. C* **81** 800

[82] CMS Collaboration 2018 CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV *CMS Physics Analysis Summary* CMS-PAS-LUM-17-004

[83] CMS Collaboration 2019 CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV *CMS Physics Analysis Summary* CMS-PAS-LUM-18-002

[84] Barlow R J and Beeston C 1993 Fitting using finite Monte Carlo samples *Comput. Phys. Commun.* **77** 219

[85] Cacciari M, Salam G P and Soyez G 2008 The anti-k_{T} jet clustering algorithm *J. High Energy Phys.* JHEP04(2008)063

[86] Cacciari M, Salam G P and Soyez G 2012 FastJet user manual *Eur. Phys. J. C* **72** 1896

[87] Davis J *et al* 2022 Constraining anomalous Higgs boson couplings to virtual photons *Phys. Rev. D* **105** 096027

[88] Gao Y *et al* 2010 Spin determination of single-produced resonances at Hadron colliders *Phys. Rev. D* **81** 075022

[89] Bolognesi S *et al* 2012 On the spin and parity of a single-produced resonance at the LHC *Phys. Rev. D* **86** 095031

[90] Anderson I *et al* 2014 Constraining anomalous HVV interactions at proton and lepton colliders *Phys. Rev. D* **89** 035007

[91] Gritsan A V, Röntsch R, Schulze M and Xiao M 2016 Constraining anomalous Higgs boson couplings to the heavy flavor fermions using matrix element techniques *Phys. Rev. D* **94** 055023

[92] Campbell J M and Ellis R K 2010 MCFM for the Tevatron and the LHC *Nucl. Phys. B* **205-206** 10

[93] Degrassi G, Giardino P P, Maltoni F and Pagani D 2016 Probing the Higgs self coupling via single Higgs production at the LHC *J. High Energy Phys.* JHEP12(2016)080

[94] Maltoni F, Pagani D, Shivaji A and Zhao X 2017 Trilinear Higgs coupling determination via single-Higgs differential measurements at the LHC *Eur. Phys. J. C* **77** 887

[95] CMS Collaboration 2025 Constraints on the Higgs boson self-coupling from the combination of single and double Higgs boson production in proton-proton collisions at $\sqrt{s} = 13$ TeV *Phys. Lett. B* **861** 139210

[96] CMS Collaboration 2021 Search for nonresonant Higgs boson pair production in final states with two bottom quarks and two photons in proton-proton collisions at $\sqrt{s} = 13$ TeV *J. High Energy Phys.* JHEP03(2021)257

[97] Monti F et al 2022 Modelling of the single-Higgs simplified template cross-sections (STXS 1.2) for the determination of the Higgs boson trilinear self-coupling *LHC Higgs Working Group Note* LHCHWG-2022-002 (CERN)

[98] Branco G C *et al* 2012 Theory and phenomenology of two-Higgs-doublet models *Phys. Rept.* **516** 1

[99] Gunion J F and Haber H E 2003 The CP conserving two Higgs doublet model: the approach to the decoupling limit *Phys. Rev. D* **67** 075019

[100] Maiani L, Polosa A D and Riquer V 2013 Bounds to the Higgs sector masses in minimal supersymmetry from LHC data *Phys. Lett. B* **724** 274

[101] Glashow S L and Weinberg S 1977 Natural conservation laws for neutral currents *Phys. Rev. D* **15** 1958
[102] Paschos E A 1977 Diagonal neutral currents *Phys. Rev. D* **15** 1966
[103] Djouadi A 2008 The anatomy of electro-weak symmetry breaking tome II. the Higgs bosons in the minimal supersymmetric model *Phys. Rept.* **459** 1
[104] Gunion J F and Haber H E 1986 Higgs bosons in supersymmetric models (I) *Nucl. Phys. B* **272** 1
[105] Haber H and Kane G 1985 The search for supersymmetry: probing physics beyond the standard model *Phys. Rept.* **117** 75
[106] Nilles H 1984 Supersymmetry, supergravity and particle physics *Phys. Rept.* **110** 1
[107] Djouadi A *et al* 2013 The post-Higgs MSSM scenario: Habemus MSSM? *Eur. Phys. J. C* **73** 2650
[108] Djouadi A *et al* 2015 Fully covering the MSSM Higgs sector at the LHC *J. High Energy Phys.* JHEP06(2015)168
[109] ATLAS Collaboration 2015 Constraints on new phenomena via Higgs boson couplings and invisible decays with the ATLAS detector *J. High Energy Phys.* JHEP11(2015)206
[110] Carena M *et al* 2013 MSSM Higgs boson searches at the LHC: benchmark scenarios after the discovery of a Higgs-like particle *Eur. Phys. J. C* **73** 2552
[111] Bahl H, Liebler S and Stefaniak T 2019 MSSM Higgs benchmark scenarios for Run 2 and beyond: the low tanβ region *Eur. Phys. J. C* **79** 279
[112] CMS Collaboration 2023 Searches for additional Higgs bosons and for vector leptoquarks in ττ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV *J. High Energy Phys.* JHEP07(2023)073
[113] CMS Collaboration 2025 Search for heavy pseudoscalar and scalar bosons decaying to a top quark pair in proton-proton collisions at $\sqrt{s} = 13$ TeV *Rept. Prog. Phys.* **88** 127801
[114] CMS Collaboration 2025 Searches for Higgs boson production through decays of heavy resonances *Phys. Rept.* **1115** 368
[115] Grzadkowski B, Iskrzynski M, Misiak M and Rosiek J 2010 Dimension-six terms in the standard model Lagrangian *J. High Energy Phys.* JHEP10(2010)085
[116] Brivio I 2021 SMEFTsim 3.0 — a practical guide *J. High Energy Phys.* JHEP04(2021)073
[117] Dawson S and Giardino P P 2018 Electroweak corrections to Higgs boson decays to γγ and W^+W^- in standard model EFT *Phys. Rev. D* **98** 095005
[118] Dawson S and Giardino P P 2018 Higgs decays to *ZZ* and *Z*γ in the standard model effective field theory: an NLO analysis *Phys. Rev. D* **97** 093003
[119] Degrande C *et al* 2021 Automated one-loop computations in the standard model effective field theory *Phys. Rev. D* **103** 096024
[120] Particle Data Group 2024 Review of particle physics *Phys. Rev. D* **110** 030001
[121] Ellis J *et al* 2021 Top, Higgs, diboson and electroweak fit to the standard model effective field theory *J. High Energy Phys.* JHEP04(2021)279
[122] Brivio I, Corbett T and Trott M 2019 The Higgs width in the SMEFT *J. High Energy Phys.* JHEP10(2019)056
[123] Brooijmans G *et al* 2020 Les Houches 2019 physics at TeV colliders: new physics working group report *11th Les Houches Workshop on Physics at TeV Colliders: PhysTeV Les Houches*
[124] Buckley A *et al* 2013 Rivet user manual *Comput. Phys. Commun.* **184** 2803
[125] Brivio I et al 2025 POPxf: an exchange format for polynomial observable predictions *Joint LHC EFT Working Group and LHC Higgs Working Group note* LHCHWG-2025-050, (CERN)
[126] Buckley A *et al* 2019 The simplified likelihood framework *J. High Energy Phys.* JHEP04(2019)064
[127] Barlow R 2003 Asymmetric errors *Proc: PHYSTAT (2003): Statistical Problems in Particle Physics, Astrophysics and Cosmology*
[128] Araz J Y 2024 Spey: smooth inference for reinterpretation studies *SciPost Phys.* **16** 032